

**ACETAMINOPHEN FOR CHILDREN- acetaminophen suppository**  
**Padagis Israel Pharmaceuticals Ltd**

-----  
**Perrigo Acetaminophen Suppositories USP, 120 mg Drug Facts**

**Active ingredient (in each rectal suppository)**

Acetaminophen 120 mg

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily

- reduces fever
- relieves minor aches, pains, and headache

**Warnings**

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**For rectal use only**

**Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- your child has liver disease
- your child is taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- fever lasts more than 3 days (72 hours), or recurs.
- pain lasts more than 5 days or gets worse.
- new symptoms occur.
- redness or swelling is present in the painful area. These may be signs of a serious condition.

**Keep out of reach of children.**

If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or warnings for adult use**
- **do not use more than directed**
- remove wrapper
- carefully insert suppository well up into the rectum

<b>Dosing Chart</b>	
<b>Age</b>	<b>Dose</b>
under 3 years	Do not use unless directed by a doctor
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Other information**

- store at 20-25°C (68-77°F) or in a cool place
- tamper-evident: suppositories are individually wrapped. Do not use if printed wrapper is opened or damaged.

**Inactive ingredient**

hydrogenated vegetable oil

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to FeverAll® active ingredient

For Children

Acetaminophen Suppositories USP, 120 mg

Pain Reliever/Fever Reducer

Ages 3-6 years

See New Warning

12 RECTAL SUPPOSITORIES, 120 mg EACH



**Drug Facts** (continued)**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**For rectal use only****Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

**Ask a doctor before use if**

- the child has liver disease
- the child is taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- fever lasts more than 3 days (72 hours), or recurs
- pain lasts more than 5 days or gets worse
- new symptoms occur
- redness or swelling is present in the painful area.

These may be signs of a serious condition. ▶

**Drug Facts** (continued)

**Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or warnings for adult use**
- **do not use more than directed**
- remove wrapper
- carefully insert suppository well up into the rectum

**Dosing Chart**

Age	Dose
under 3 years	Do not use unless directed by a doctor
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Other information**

- store at 20-25°C (68-77°F) or in a cool place
- tamper-evident: suppositories are individually wrapped. Do not use if printed wrapper is opened or damaged.

**Inactive ingredient** hydrogenated vegetable oil

**Questions or comments?** 1-800-719-9260

NDC 45802-732-30

**Acetaminophen  
Suppositories USP, 120 mg**  
Pain Reliever/Fever Reducer

**ACETAMINOPHEN FOR CHILDREN**

acetaminophen suppository

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:45802-732
<b>Route of Administration</b>	RECTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	120 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>COTTONSEED OIL</b> (UNII: H3E878020N)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-732-33	100 in 1 CARTON	12/14/2010	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:45802-732-30	12 in 1 CARTON	03/09/2011	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA070607	12/14/2010	

**Labeler** - Padagis Israel Pharmaceuticals Ltd (600093611)

Revised: 11/2021

Padagis Israel Pharmaceuticals Ltd