#### LANSOPRAZOLE - lansoprazole capsule, delayed release Gericare Pharmaceuticals

-----

#### Lansoprazole Delayed Release Capsules USP

#### Active ingredient(s)

Lansoprazole USP, 15 mg

#### Purpose

Acid reducer

## Use(s)

- treats frequent heartburn (occurs **<u>2 or more</u>** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

## Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

#### Do not use

• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

## Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

## Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

# Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

## <u>14-Day Course of Treatment</u>

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a days
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

## **<u>Repeated 14-Day Courses (if needed)</u>**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

## Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20° 25°C (68° 77° F)
- keep product out of high heat and humidity
- protect product from moisture

#### **Inactive ingredients**

FD&C Blue No. 2, gelatin, hydroxypropyl cellulose, iron oxide black, iron oxide red, iron oxide yellow, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol 6000, polysorbate 80, sodium lauryl sulphate, starch (corn), sucrose, sugar spheres, talc, titanium dioxide

## Questions or comments?

call 1-888-375-3784

Manufactured by:

## Dr. Reddy's Laboratories Limited

Bachupally - 500 090 INDIA

## Principal Display Panel - container carton

#### **Container carton : 14's count**

	LEGEL MODIA YOU THE MENO		GERI (ARE) Lansoprazole Magai Malan Dynas Rift Franz (Malanari		
Characterization Construction	Even (a second sec	Progr Facher Scottering ( The Control of the Contr	<text></text>	b british bank coreasis trong mentangan an ana ang	
	447452		5 - 57896-74162 -		

lansoprazole capsule, delayed rele	ase				
r · · · · · · · · · · · · · · · · · · ·					
Product Information					
Product T ype	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-761(NDC:55111-739)		
Route of Administration	ORAL				
Active Ingredient/Active Moi	ety				
Ing	<b>Basis of Strength</b>	Strength			
- Lansoprazole (UNII: 0K5C5T2QPG) (L	ansoprazole - UNII:0K5C	5T2QPG)	Lansoprazole	15 mg	
Inactive Ingredients					
Inactive Ingredients					
Inactive Ingredients	Ingredient N	ame		Strength	
, i i i i i i i i i i i i i i i i i i i	•	ame		Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DC	2 (K)	ame		Strengtl	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6	2K) 75)	ame		Strengtl	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43	2K) 75)	ame		Strength	
Inactive Ingredients FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UNI	2K) 75) 802MRT)	ame		Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UI	2K) 75) 802MRT) NII: RFW2ET671P)	āme		Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UNI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI			Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0E)	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA)	I: 2165RE0K14)		Strength	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UNI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0E) METHACRYLIC ACID - ETHYL ACRY	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) <b>(LATE COPOLYMER (1</b> )	I: 2165RE0K14)	5T8J)	Strengt	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0 E METHACRYLIC ACID - ETHYL ACRY POLYETHYLENE GLYCOL 6000 (UI	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) 7LATE COPOLYMER (1: NII: 30IQX730WE)	I: 2165RE0K14)	5T8J)	Strengtl 	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UN FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0 E) METHACRYLIC ACID - ETHYL ACRY POLYETHYLENE GLYCOL 6000 (UN POLYSORBATE 80 (UNII: 60ZP39ZG	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) <b>7LATE COPOLYMER (1:</b> NII: 30IQX730WE) 8H)	I: 2165RE0K14)	5T8J)	Strengtl 	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0 E METHACRYLIC ACID - ETHYL ACRY POLYETHYLENE GLYCOL 6000 (UI POLYSORBATE 80 (UNII: 60ZP39ZG SODIUM LAURYL SULFATE (UNII: 36	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) <b>7LATE COPOLYMER (1:</b> NII: 30IQX730WE) 8H)	I: 2165RE0K14)	5T8J)	Strengtl Strengtl	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC OXIDE RED (UNII: 1K09F3G6 FERRIC OXIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE (UNI FERROSOFERRIC OXIDE (UNII: XM0 HYDROXYPROPYL CELLULOSE, LC MAGNESIUM CARBONATE (UNII: 0 E METHACRYLIC ACID - ETHYL ACRY POLYETHYLENE GLYCOL 6000 (UN POLYSORBATE 80 (UNII: 60ZP39ZG SODIUM LAURYL SULFATE (UNII: 30 STARCH, CORN (UNII: 08232NY3SJ)	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) <b>7LATE COPOLYMER (1:</b> NII: 30IQX730WE) 8H)	I: 2165RE0K14)	5T8J)	Strengtl Strengtl	
FD&C BLUE NO. 2 (UNII: L06K8R7DC FERRIC O XIDE RED (UNII: 1K09F3G6 FERRIC O XIDE YELLOW (UNII: EX43 GELATIN (UNII: 2G86QN327L)	2K) 75) 802MRT) NII: RFW2ET671P) M87F357) <b>DW SUBSTITUTED</b> (UNI 53J927NA) <b>7LATE COPOLYMER (1:</b> NII: 30IQX730WE) 8H)	I: 2165RE0K14)	5T8J)	Strengt Strengt	

Product Characteristics									
Color	PINK (opa	aque pink colored cap) , GREEN (opaque green colored t	Score		no score				
Shape	CAPSULE	2	Size		3mm				
Flavor					int Code	RDY;398			
Contains									
Packaging									
# Item	Item Code Package Description		Marketing Start	Date	Marketing	g End Date			
1 NDC:5789	57896-761-42 3 in 1 PACKAGE, COMBINATION		04/25/2013						
1		14 in 1 BOTTLE; Type 0: Not a Combination Product							
Marketing Information									
Marketing	Marketing Category Application Number or Monograph C		Marketing Start	Date	Marketing	g End Date			
ANDA		ANDA202194	04/25/2013						

Labeler - Gericare Pharmaceuticals (611196254)

Revised: 12/2017

Gericare Pharmaceuticals