

IBUPROFEN 200- ibuprofen tablet, film coated
Walgreen Company

Walgreens 44-393

Active ingredient (in each orange caplet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - minor pain of arthritis
 - backache
 - menstrual cramps
 - toothache
 - muscular aches
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

NDC 0363-0393-20

Walgreens

**Compare to the active ingredient
in Motrin® IB Caplets††**

• WALGREENS •
PHARMACIST RECOMMENDED†

EASY OPEN

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg/
PAIN RELIEVER/FEVER REDUCER (**NSAID**)

225

CAPLETS

ACTUAL SIZE

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

50844 ORG122139320

†Walgreens Pharmacist Recommended.

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Caplets.

DISTRIBUTED BY: **WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015**
100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co.

NDC 0363-0393-20

Walgreens WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredient in Motrin® IB Caplets†

EASY OPEN

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER/ FEVER REDUCER (NSAID)

225 CAPLETS

ACTUAL SIZE

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each orange caplet) Purpose
Ibuprofen USP (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ toothache
■ backache ■ menstrual cramps ■ the common cold
■ minor pain of arthritis
■ Temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ rash ■ hives ■ facial swelling ■ shock
■ asthma (wheezing) ■ skin reddening. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

50844 ORG122139320 W30RG0122-F
REV0422
Walgreens Pharmacist Recommended
Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Caplets.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co.
ITEM 546335 W00000-0000-0

3 11917108981 2

Discard Seal, Empty & Replace Cap
PLASTIC BOTTLE
how2recycle.info

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if ■ you are taking a diuretic
■ you have a history of stomach problems, such as heartburn ■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint
■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ weakness in one part or side of body
■ slurred speech ■ trouble breathing ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
■ If pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information ■ avoid excessive heat 40°C (104°F)
■ store between 20°-25°C (68°-77°F)
■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

(Questions or comments?) Call 1-800-426-9391
8:30 AM-4:00 PM ET, Monday-Friday

STOP PEELING

Walgreens 44-393

IBUPROFEN 200

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0393
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	44;393
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0393-20	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	
2	NDC:0363-0393-15	1 in 1 CARTON	03/01/1999	01/08/2022
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:0363-0393-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/01/1999	04/22/2023
4	NDC:0363-0393-37	1 in 1 CARTON	03/01/1999	02/07/2021
4		75 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

5	NDC:0363-0393-12	1 in 1 CARTON	03/01/1999	04/07/2023
5		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

Labeler - Walgreen Company (008965063)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0393)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0393)