IBUPROFEN- ibuprofen tablet, film coated Kroger Company

Kroger Co. Ibuprofen Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

• are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

• take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used

Adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- Children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-632-6900

Principal Display Panel

COMPARE TO the active ingredient of ADVIL® IBUPROFEN TABLETS

See back panel

Kroger_®

Read label for current directions and warnings

OUR PHARMACIST RECOMMENDED

Ibuprofen Tablets 200 mg

Pain Reliever/Fever Reducer (NSAID)

actual size

500 COATED TABLETS



IBUPROFEN

ibuprofen tablet, film coated

Product Information							
Product Type	HUMAN OTC DRUG	Item Code (Source) ND		NDC:30	DC:30142-604		
Route of Administration	ORAL						
Active Ingredient/Active Moiety							
Ingredient Name Basis of Strengt					Strength		
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) IBUPROFEN				200 mg			
Inactive Ingredients							
Inactive Ingredients	Ingredient Name				Strength		
Inactive Ingredients SILICON DIOXIDE (UNII: ETJ7Z6X	-				Strength		
-	BU4)				Strength		
SILICON DIOXIDE (UNII: ETJ7Z6X	BU4) SJ)				Strength		
SILICON DIOXIDE (UNII: ETJ7Z6X STARCH, CORN (UNII: 08232NY3	BU4) SJ) NII: M28OL1HH48)				Strength		

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	12
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:30142-604- 62	1 in 1 CARTON	01/15/1988				
1		24 in 1 BOTTLE; Type 0: Not a Combination Product					
2	NDC:30142-604- 71	1 in 1 CARTON	02/15/1988	03/31/2023			
2		50 in 1 BOTTLE; Type 0: Not a Combination Product					
3	NDC:30142-604- 78	1 in 1 CARTON	09/15/1988				
3		100 in 1 BOTTLE; Type 0: Not a Combination Product					
4	NDC:30142-604- 85	250 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2005	03/31/2023			
5	NDC:30142-604- 90	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/04/1991				
6	NDC:30142-604- 93	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/08/2011	12/29/2017			
7	NDC:30142-604- 76	1 in 1 CARTON	01/14/1992	03/31/2020			
7		120 in 1 BOTTLE; Type 0: Not a Combination Product					
Marketing Information							
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			

01/15/1988

Labeler - Kroger Company (006999528)

ANDA072096

ANDA