

STAFAC 50- virginiamycin powder
Phibro Animal Health

Stafac® 50
(virginiamycin)
TYPE A MEDICATED ARTICLE

Active Drug Ingredient

Virginiamycin 11%

(Contains 50 g virginiamycin activity per lb)

Ingredients:

Processed grain by-products, roughage products, calcium carbonate,
carboxymethylcellulose, mineral oil

Caution:

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive drug to use by or on the order of a licensed veterinarian.

For further manufacture only.

Follow Directions on Back Panel

Stafac is registered trademark of Phibro Animal Health Corporation

Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.6 kg)

Distributed by:

Phibro Animal Health Corporation

Teaneck, NJ 07666

NADA #91-467, Approved by FDA

9910000

101-9107-09A

(50 g virginiamycin activity per lb)

For use in swine and poultry feeds as specified below

Directions for Use

Mixing Directions—Swine

Stafac 50 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of Stafac 50 required from the table below. Dilute Stafac 50 with a portion of one of the feed ingredients (enough to make a quantity equal to 10 lb per ton) before final mixing.

Swine	Required level of virginiamycin activity	Required amount of Stafac 50 per ton of	Amount of virginiamycin
--------------	---	--	--------------------------------

	per ton of Type C Medicated Feed (complete feed)	Type C Medicated Feed (complete feed)	activity provided to animal per lb of body weight per day
Treatment of swine dysentery in nonbreeding swine weighing over 120 lb	100 g for 2 weeks	2 lb	2.27 mg
Treatment and control of swine dysentery in swine weighing up to 120 lb	100 g for 2 weeks followed by 50 g	2 lb 1 lb	2.27 mg 1.14 mg
Aid in control of swine dysentery in swine weighing up to 120 lb For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred	25 g	8 oz	0.57 mg
CAUTION: Not for use in breeding swine over 120 lbs.			

Mixing Directions—Poultry

Mix 1 lb of Stafac 50 with 9 lb of feed ingredients to make 10 lb of an intermediate mix to contain 5 g/lb virginiamycin. Add 1 lb of the intermediate feed premix for each 5 g of virginiamycin needed to medicate a ton of feed.

Poultry	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)
For use in broiler chickens		
Prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to virginiamycin	20 g	180 g
Not for Use in Laying Chickens		

Storage and Handling

Feed Continuously as Sole Ration

Store At Or Below 25°C (77°F), Excursions Permitted Up to 40°C (104°F)

Close Container After Use

Not for Human Use

Restricted Drug (California) – Use Only As Directed

Package/Label Principal Display Panel



0211 1 03 64104 99100 0

Stafac[®] 50 (virginiamycin)

TYPE A
MEDICATED ARTICLE

(50 g virginiamycin activity per lb)
For use in swine and poultry feeds as specified below



Directions for Use
Mixing Directions—Swine
Stafac 50 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of Stafac 50 required from the table below. Place Stafac 50 with a portion of one of the feed ingredients through to make a quantity equal to 10 lb per ton before final mixing.

Swine	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)	Amount of virginiamycin activity provided per animal per lb of body weight per day
Treatment and control of swine in non-lactating sows weighing over 120 lb	100 g for 2 weeks	2 lb	2.27 mg
Treatment and control of swine in sows in lactation weighing up to 120 lb	100 g for 2 weeks	2 lb	2.27 mg
Ad as control of swine dysentery in sows weighing up to 120 lb	25 g	0.5 lb	0.57 mg

For use in sows or on piglets with a history of one or more outbreaks but where symptoms have not yet occurred.

CAUTION: Not for use in breeding sows over 120 lbs.

Mixing Directions—Poultry
Mix 1 lb of Stafac 50 with 3 lb of feed ingredients to make 10 lb of an intermediate mix to contain 5 g/lb virginiamycin. Add 1 lb of the intermediate feed premix for each 5 g of virginiamycin needed to medicate a ton of feed.

Poultry	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)
For use in broiler chickens	25 g	0.5 lb
Prevention of necrotic enteritis caused by <i>Campylobacter parvorum</i> susceptible to virginiamycin	25 g	0.5 lb

Not for Use in Layer Chickens
Feed Contaminants as Side Effect
Store At or Below 20°C (77°F). Excursions Permitted Up to 40°C (104°F)
Close Container After Use
Not For Human Use
Restricted Drug (California) - Use Only as Directed

9910000

101-9107-00A

Phibro Animal Health Corporation
Teaneck, NJ 07666

Stafac[®] 50
(virginiamycin)



Stafac[®] 50 (virginiamycin)

TYPE A
MEDICATED ARTICLE

Active Drug Ingredient
Virginiamycin 11%
(Contains 50 g virginiamycin activity per lb)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Ingredients:
Processed grain by-products, roughage products, calcium carbonate, carboxymethylcellulose, mineral oil

For further manufacture only.

Follow Directions on Back Panel

Stafac is a registered trademark of Phibro Animal Health Corporation
Phibro Animal Health Corporation, Teaneck, NJ 07666

NADA #91-467, Approved by FDA
9910000
101-9107-00A

Net Weight 50 lb (22.6 kg)



Stafac[®] 50
(virginiamycin)

STAFAC 50

virginiamycin powder

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66 104-88 15
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VIRGINIAMYCIN (UNII: C49WS9N75L) (VIRGINIAMYCIN - UNII:C49WS9N75L)	VIRGINIAMYCIN	50 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
RICE BRAN (UNII: R60QEP13IC)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66 104-88 15-0	22.6 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA091467	03/24/2010	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 12/2018

Phibro Animal Health