PEPTO-BISMOL ULTRA- bismuth subsalicylate tablet Procter & Gamble Manufactura, S. de R.L. de C.V.

Pepto[®] BISMOL[™] ULTRA

Drug Facts

Active ingredient (in each caplet)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - \circ indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem

bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
- 1 caplet (1 dose) every ¹/₂ hour or 2 caplets (2 doses) every hour as needed for diarrhea
- 1 caplet (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (8 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

• each caplet contains: calcium 54 mg

- salicylate 199 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

calcium carbonate, D&C Red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

Questions?

1-800-717-3786

TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.

MADE IN MEXICO DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 12 Caplet Carton

Pepto

Bismol® ULTRA

Bismuth Subsalicylate Upset Stomach Reliever/ Antidiarrheal

2x CONCENTRATED FORMULA*

5 SYMPTOM RELIEF

NAUSEA HEARTBURN INDIGESTION UPSET STOMACH DIARRHEA

12 CAPLETS

TAMPER EVIDENT: Do not use if inner printed seal on bottl is broken or missing.

OfOSG Agrulomsia



PEPTO-BISMOL ULTRA

bismuth subsalicylate tablet

UNII:O414PZ4LPZ)

| Product Information | | | | | |
|-------------------------------------|--------------------------|----------------|------------|----------|----------|
| Product Type | HUMAN OTC DRUG | Item Code (Sou | rce) | NDC:5893 | 3-478 |
| Route of Administration | ORAL | | | | |
| | | | | | |
| • • • • • • • • | | | | | |
| Active Ingredient/Active | Molety | | | | |
| Ingr | edient Name | | Basis of S | trength | Strength |
| BISMUTH SUBSALICYLATE (UNII: | 62TEY51RR1) (SALICYLIC A | CID - | | TE | 525 mg |

SUBSALICYLATE

| - | | | | | | | | |
|------------------|---|--|--|-----------------|---------------------------|-----------------------|--|--|
| In | active Ingre | dient | ts | | | | | |
| | | | Ingredient | Name | | Strength | | |
| MA | MAGNESIUM STEARATE (UNII: 70097M6I30) | | | | | | | |
| MA | ANNITOL (UNII: 30 | OWL53 | L36A) | | | | | |
| CE | LLULOSE, MICR | OCRY | STALLINE (UNII: OP1R32D6 | 1U) | | | | |
| РО | LYSORBATE 80 | (UNII: | 6OZP39ZG8H) | | | | | |
| СА | LCIUM CARBON | ATE (l | JNII: HOG9379FGK) | | | | | |
| PO | VIDONE (UNII: FZ | Z 989G | H94E) | | | | | |
| D۵ | C RED NO. 27 (| UNII: 2 | LRS185U6K) | | | | | |
| SO | DIUM STARCH C | GLYCO | LATE TYPE A POTATO (U | NII: 5856J3G2A2 |) | | | |
| SIL | | (UNII: E | TJ7Z6XBU4) | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| Pr | roduct Chara | icter | istics | | | | | |
| Color | | | pink | Score | | no score | | |
| Shape | | | OVAL (Caplet) | Size | | 18mm | | |
| Flavor | | | | Imprint | Code | Pepto | | |
| Co | ontains | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| Pa | ackaging | | | | | | | |
| Pa # | ackaging Item Code | | Package Descript | tion | Marketing Start Date | Marketing End Date | | |
| # | | 1 in 1 | Package Descript | | - | - | | |
| # | Item Code NDC:58933-478- | | CARTON 1 BOTTLE; Type 0: Not a Co | | Date | - | | |
| # 1 1 | Item Code NDC:58933-478- | 24 in Produ | CARTON 1 BOTTLE; Type 0: Not a Co | ombination | Date | - | | |
| # 1 1 | Item Code NDC:58933-478- 24 NDC:58933-478- | 24 in Produ 1 in 1 | CARTON 1 BOTTLE; Type 0: Not a Co ct CARTON 1 BOTTLE; Type 0: Not a Co | ombination | Date 02/01/2018 | - | | |
| # 1 1 2 | Item Code NDC:58933-478- 24 NDC:58933-478- | 24 in Produ 1 in 1 12 in | CARTON 1 BOTTLE; Type 0: Not a Co ct CARTON 1 BOTTLE; Type 0: Not a Co | ombination | Date 02/01/2018 | - | | |
| # 1 2 2 | Item Code NDC:58933-478- 24 NDC:58933-478- | 24 in Produ 1 in 1 12 in Produ | CARTON 1 BOTTLE; Type 0: Not a Co ct CARTON 1 BOTTLE; Type 0: Not a Co ct | ombination | Date 02/01/2018 | - | | |
| # 1 2 2 | Item Code NDC:58933-478- 24 NDC:58933-478- 12 | 24 in Produ 1 in 1 12 in Produ | CARTON 1 BOTTLE; Type 0: Not a Co ct CARTON 1 BOTTLE; Type 0: Not a Co ct | ombination | Date 02/01/2018 | - | | |

Labeler - Procter & Gamble Manufactura, S. de R.L. de C.V. (812807550)

Revised: 10/2023

Procter & Gamble Manufactura, S. de R.L. de C.V.