

**PEPTO-BISMOL ULTRA- bismuth subsalicylate tablet**  
**Procter & Gamble Manufactura, S. de R.L. de C.V.**

-----  
**Pepto<sup>®</sup> BISMOL<sup>™</sup> ULTRA**

***Drug Facts***

**Active ingredient (in each caplet)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

**Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem

- bloody or black stool

### **Ask a doctor before use if you have**

- fever
- mucus in the stool

### **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
  - 1 caplet (1 dose) every ½ hour or 2 caplets (2 doses) every hour as needed for diarrhea
  - 1 caplet (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (8 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each caplet contains:**calcium 54 mg

- salicylate 199 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

calcium carbonate, D&C Red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

### **Questions?**

**1-800-717-3786**

**TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.**

**MADE IN MEXICO**

**DIST. BY PROCTER & GAMBLE,**

**CINCINNATI OH 45202**

### **PRINCIPAL DISPLAY PANEL - 12 Caplet Carton**

Pepto

Bismol® ULTRA

Bismuth Subsalicylate

Upset Stomach Reliever/

Antidiarrheal

2x CONCENTRATED FORMULA\*

5 SYMPTOM RELIEF

NAUSEA

HEARTBURN

INDIGESTION

UPSET STOMACH

DIARRHEA

12 CAPLETS



**Drug Facts**

Active ingredient (in each caplet)	Purpose
Bismuth subsalicylate ..... 525 mg	Upset stomach reliever and antidiarrheal

**Uses** relieves

- upset stomach due to overindulgence in food and drink, including:
- nausea
- travelers' diarrhea
- heartburn
- gas
- indigestion
- belching
- fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**Drug Facts (continued)**

If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
  - 1 caplet (1 dose) every ½ hour or 2 caplets (2 doses) every hour as needed for diarrhea
  - 1 caplet (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (8 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years; ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- each caplet contains: calcium 54 mg
- salicylate 199 mg
- very low sodium
- avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** calcium carbonate, D&C Red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

**Questions? 1-800-717-3786**

\*vs Original Pepto Bismol®

**MADE IN MEXICO**  
 DIST. BY PROCTER & GAMBLE,  
 CINCINNATI OH 45202  
 Patents: www.pg.com/patents  
 www.pepto-bismol.com

**P&G**  
 www.pg.com  
 91169459



<b>PEPTO-BISMOL ULTRA</b>			
bismuth subsalicylate tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58933-478
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL (Caplet)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	Pepto
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-478-24	1 in 1 CARTON	02/01/2018	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:58933-478-12	1 in 1 CARTON	02/01/2018	
2		12 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/01/2018	

**Labeler** - Procter & Gamble Manufactura, S. de R.L. de C.V. (812807550)

Revised: 10/2025

Procter & Gamble Manufactura, S. de R.L. de C.V.