

NOREPINEPHRINE BITARTRATE- norepinephrine bitartrate injection, solution

Baxter Healthcare Corporation

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **NOREPINEPHRINE BITARTRATE IN DEXTROSE injection** and **NOREPINEPHRINE BITARTRATE IN SODIUM CHLORIDE injection** safely and effectively. See full prescribing information for **NOREPINEPHRINE BITARTRATE IN DEXTROSE injection** and **NOREPINEPHRINE BITARTRATE IN SODIUM CHLORIDE injection**.

NOREPINEPHRINE BITARTRATE IN DEXTROSE injection, for intravenous use

NOREPINEPHRINE BITARTRATE IN SODIUM CHLORIDE injection, for intravenous use

Initial U.S. Approval: 1950

INDICATIONS AND USAGE

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection is a catecholamine indicated for restoration of blood pressure in adult patients with acute hypotensive states. (1)

DOSAGE AND ADMINISTRATION

- No further dilution prior to infusion is required. (2.1)
- Initiate at 8 mcg/min to 12 mcg/min, and adjust the rate to maintain blood pressure sufficient to maintain the circulation of vital organs. (2.2)
- The average maintenance dose ranges from 2 mcg/min to 4 mcg/min. (2.2)

DOSAGE FORMS AND STRENGTHS

Injection: 250-mL single dose-containers with (3)

- 4 mg equivalent of norepinephrine (16 mcg/mL) in 5% Dextrose or 0.9% Sodium Chloride.
- 8 mg equivalent of norepinephrine (32 mcg/mL) in 5% Dextrose or 0.9% Sodium Chloride.
- 16 mg equivalent of norepinephrine (64 mcg/mL) in 5% Dextrose or 0.9% Sodium Chloride.

CONTRAINDICATIONS

- None. (4)

WARNINGS AND PRECAUTIONS

- **Tissue Ischemia:** Infuse Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection into a large vein. To prevent sloughing and necrosis in areas in with extravasation, infiltrate the area with an adrenergic blocking agent in saline. (5.1)
- **Hypotension After Abrupt Discontinuation:** Gradually taper a norepinephrine infusion to prevent hypotension. (5.2)
- **Cardiac Arrhythmias:** Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection may cause arrhythmias. Monitor cardiac function in patients with underlying heart disease. (5.3)
- **Allergic Reactions with Sulfite:** Norepinephrine Bitartrate in Sodium Chloride Injection contains sodium metabisulfite: Sulfite may cause allergic-type reactions. (5.4)

ADVERSE REACTIONS

Most common adverse reactions are hypertension, bradycardia, ischemic injury, anxiety, headache, respiratory difficulty, pulmonary edema, and extravasation necrosis at injection site. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Monoamine oxidase inhibitors (MAOI) or tricyclic antidepressants of the triptyline or imipramine types may result in hypertension. (7.1, 7.2)
- Antidiabetics: Norepinephrine can decrease insulin sensitivity and raise blood glucose (7.3)
- Cyclopropane and halothane anesthetics increase cardiac autonomic irritability. (7.4)

-----**USE IN SPECIFIC POPULATIONS**-----

- Elderly patients may be at greater risk of developing adverse reactions (8.5)

See 17 for **PATIENT COUNSELING INFORMATION**.

Revised: 2/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Instructions

2.2 Dosage

2.3 Drug Incompatibilities

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Tissue Ischemia

5.2 Hypotension after Abrupt Discontinuation

5.3 Cardiac Arrhythmias

5.4 Allergic Reactions Associated with Sulfite

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

7.1 MAO-Inhibiting Drugs

7.2 Tricyclic Antidepressants

7.3 Antidiabetics

7.4 Halogenated Anesthetics

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection is indicated to raise blood pressure in adult patients with severe, acute hypotension.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Instructions

Correct Hypovolemia

Address hypovolemia before initiation of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection therapy. If the patient does not respond to therapy, suspect occult hypovolemia [see *Warnings and Precautions (5.1)*].

Administration

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection is a ready to administer product that requires no further dilution prior to infusion. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use the solution if its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Infuse Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection into a large vein. Avoid infusions into the veins of the leg in the elderly or in patients with occlusive vascular disease of the legs [see *Warnings and Precautions (5.1)*]. Avoid using a catheter-tie-in technique.

The choice of appropriate concentration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection depends on clinical fluid volume requirements. Use higher concentration solutions in patients requiring fluid restriction.

Discontinuation

When discontinuing the infusion, reduce the flow rate gradually. Avoid abrupt withdrawal.

2.2 Dosage

After an initial dosage of 8 mcg per minute to 12 mcg per minute via intravenous infusion, assess patient response and adjust dosage to maintain desired hemodynamic effect. Monitor blood pressure every two minutes until the desired hemodynamic effect is achieved, and then monitor blood pressure every five minutes for the duration of the infusion.

Typical maintenance intravenous dosage is 2 mcg per minute to 4 mcg per minute.

2.3 Drug Incompatibilities

Avoid contact with iron salts, alkalis, or oxidizing agents.

Whole blood or plasma, if indicated to increase blood volume, should be administered separately.

3 DOSAGE FORMS AND STRENGTHS

Injection: Norepinephrine bitartrate in either 5% Dextrose or 0.9% Sodium Chloride is a colorless to slightly yellow solution for intravenous infusion, supplied in 250-mL single dose containers as:

- 4 mg equivalent of norepinephrine (16 mcg/mL).
- 8 mg equivalent of norepinephrine (32 mcg/mL).
- 16 mg equivalent of norepinephrine (64 mcg/mL).

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Tissue Ischemia

Administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection to patients who are hypotensive from hypovolemia can result in severe peripheral and visceral vasoconstriction, decreased renal perfusion and reduced urine output, tissue hypoxia, lactic acidosis, and reduced systemic blood flow despite “normal” blood pressure. Address hypovolemia prior to initiating Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection [see *Dosage and Administration (2.1)*]. Avoid Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection in patients with mesenteric or peripheral vascular thrombosis, as this may increase ischemia and extend the area of infarction.

Gangrene of the extremities has occurred in patients with occlusive or thrombotic vascular disease or who received prolonged or high dose infusions. Monitor for changes to the skin of the extremities in susceptible patients.

Extravasation of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection may cause necrosis and sloughing of surrounding tissue. To reduce the risk of extravasation, infuse into a large vein, check the infusion site frequently for free flow, and monitor for signs of extravasation [see *Dosage and Administration (2.1)*].

Emergency Treatment of Extravasation

To prevent sloughing and necrosis in areas in which extravasation has occurred, infiltrate the ischemic area as soon as possible, using a syringe with a fine hypodermic needle with 5 mg to 10 mg of phentolamine mesylate in 10 mL to 15 mL of 0.9% Sodium Chloride Injection in adults.

Sympathetic blockade with phentolamine causes immediate and conspicuous local hyperemic changes if the area is infiltrated within 12 hours.

5.2 Hypotension after Abrupt Discontinuation

Sudden cessation of the infusion rate may result in marked hypotension. When discontinuing the infusion, gradually reduce the Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection infusion rate while expanding blood volume with intravenous fluids.

5.3 Cardiac Arrhythmias

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection elevates intracellular calcium concentrations and may cause arrhythmias, particularly in the setting of hypoxia or hypercarbia. Perform continuous cardiac monitoring of patients with arrhythmias.

5.4 Allergic Reactions Associated with Sulfite

This product contains sodium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in non-asthmatic people.

6 ADVERSE REACTIONS

The following serious adverse reactions are described in greater detail in other sections:

- Tissue Ischemia [see *Warnings and Precautions (5.1)*]
- Hypotension [see *Warnings and Precautions (5.2)*]
- Cardiac Arrhythmias [see *Warnings and Precautions (5.3)*]
- Allergic Reactions associated with Sulfite [see *Warnings and Precautions (5.4)*]

The most common adverse reactions are hypertension and bradycardia.

The following adverse reactions can occur:

Nervous system disorders: Anxiety, headache

Respiratory disorders: Respiratory difficulty, pulmonary edema

General disorders and administration site conditions: Extravasation, injection site necrosis [see *Warnings and Precautions (5.1)*].

7 DRUG INTERACTIONS

7.1 MAO-Inhibiting Drugs

Co-administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection with monoamine oxidase (MAO) inhibitors or other drugs with MAO-inhibiting properties (e.g., linezolid) can cause severe, prolonged hypertension.

If administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection cannot be avoided in patients who recently have

received any of these drugs and in whom, after discontinuation, MAO activity has not yet sufficiently recovered, monitor for hypertension.

7.2 Tricyclic Antidepressants

Co-administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection with tricyclic antidepressants (including amitriptyline, nortriptyline, protriptyline, clomipramine, desipramine, imipramine) can cause severe, prolonged hypertension. If administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection cannot be avoided in these patients, monitor for hypertension.

7.3 Antidiabetics

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection can decrease insulin sensitivity and raise blood glucose. Monitor glucose and consider dosage adjustment of antidiabetic drugs.

7.4 Halogenated Anesthetics

Concomitant use of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection with halogenated anesthetics (e.g., cyclopropane, desflurane, enflurane, isoflurane, and sevoflurane) may lead to ventricular tachycardia or ventricular fibrillation. Monitor cardiac rhythm in patients receiving concomitant halogenated anesthetics.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited published data consisting of a small number of case reports and multiple small trials involving the use of norepinephrine in pregnant women at the time of delivery have not identified an increased risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. There are risks to the mother and fetus from hypotension associated with septic shock, myocardial infarction and stroke which are medical emergencies in pregnancy and can be fatal if left untreated. (*see Clinical Considerations*). In animal reproduction studies, using high doses of intravenous norepinephrine resulted in lowered maternal placental blood flow. Clinical relevance to changes in the human fetus is unknown since the average maintenance dose is ten times lower (*see Data*). Increased fetal reabsorptions were observed in pregnant hamsters after receiving daily injections at approximately 2 times the maximum recommended dose on a mg/m³ basis for four days during organogenesis (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2%-4% and 15%-20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Hypotension associated with septic shock, myocardial infarction, and stroke are medical emergencies in pregnancy which can be fatal if left untreated. Delaying treatment in pregnant women with hypotension associated with septic shock, myocardial infarction and stroke may increase the risk of maternal and fetal morbidity and mortality. Life-sustaining therapy for the pregnant woman should not be withheld due to potential concerns regarding the effects of norepinephrine on the fetus.

Data

Animal Data

A study in pregnant sheep receiving high doses of intravenous norepinephrine (40 mcg/min, at approximately 10 times the average maintenance dose of 2 mcg/min -4 mcg/min in human, on a mg/kg basis) exhibited a significant decrease in maternal placental blood flow. Decreases in fetal oxygenation, urine and lung liquid flow were also observed.

Norepinephrine administration to pregnant rats on Gestation Day 16 or 17 resulted in cataract production in rat fetuses.

In hamsters, an increased number of resorptions (29.1% in study group vs. 3.4% in control group), fetal microscopic liver abnormalities and delayed skeletal ossification were observed at approximately 2 times the maximum recommended intramuscular or subcutaneous dose (on a mg/m² basis at a maternal subcutaneous dose of 0.5 mg/kg/day from Gestation Day 7-10).

8.2 Lactation

Risk Summary

There are no data on the presence of norepinephrine in either human or animal milk, the effects on the breastfed infant, or the effects on milk production. Clinically relevant exposure to the infant is not expected based on the short half-life and poor oral bioavailability of norepinephrine.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Clinical studies of norepinephrine did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

Avoid administration of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection into the veins in the leg in elderly patients [*see Warnings and Precautions (5.1)*].

10 OVERDOSAGE


Overdosage with Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection may result in headache, severe hypertension, reflex bradycardia, marked increase in peripheral resistance, and decreased cardiac output.

In case of overdosage, discontinue Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection until the condition of the patient stabilizes.

11 DESCRIPTION

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection contains norepinephrine, a sympathomimetic amine. Norepinephrine is sometimes referred to as l-arterenol/Levarterenol or l-norepinephrine which differs from epinephrine by the absence of a methyl group on the nitrogen atom.

Chemically, norepinephrine bitartrate monohydrate is (-)- α -(aminomethyl)-3,4-dihydroxybenzyl alcohol tartrate (1:1) (salt) monohydrate and has the following structural formula:



Norepinephrine is sparingly soluble in water, very slightly soluble in alcohol and ether, and readily soluble in acids.

Norepinephrine Bitartrate in 5% Dextrose Injection is supplied as a sterile aqueous solution administered by intravenous infusion. Each mL contains the equivalent of 16 mcg, 32 mcg, or 64 micrograms of norepinephrine base supplied as 31.9 mcg, 63.8 mcg, and 127.6 micrograms per mL of norepinephrine bitartrate monohydrate. It contains dextrose monohydrate (50 mg/mL) and may contain hydrochloric acid and/or sodium hydroxide for pH adjustment. It has a target pH of 3.7. The air in the containers has been displaced by nitrogen gas.

Norepinephrine Bitartrate in 0.9% Sodium Chloride Injection is supplied as a sterile aqueous solution administered by intravenous infusion. Each mL contains the equivalent of 16 mcg, 32 mcg, or 64 micrograms of norepinephrine base supplied as 31.9 mcg, 63.8 mcg, and 127.6 micrograms per mL of norepinephrine bitartrate monohydrate. It contains sodium chloride (9 mg/mL), sodium metabisulfite (3.2 mcg/mL, 6.4 mcg/mL and

12.8 mcg/mL, respectively) and may contain hydrochloric acid and/or sodium hydroxide for pH adjustment. It has a target pH of 3.8. The air in the containers has been displaced by nitrogen gas.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Norepinephrine is a peripheral vasoconstrictor (alpha-adrenergic action) and an inotropic stimulator of the heart and dilator of coronary arteries (beta-adrenergic action).

12.2 Pharmacodynamics

The primary pharmacodynamic effects of norepinephrine are cardiac stimulation and vasoconstriction. Cardiac output is generally unaffected, although it can be decreased, and total peripheral resistance is also elevated. The elevation in resistance and pressure result in reflex vagal activity, which slows the heart rate and increases stroke volume. The elevation in vascular tone or resistance reduces blood flow to the major abdominal organs as well as to skeletal muscle. Coronary blood flow is substantially increased secondary to the indirect effects of alpha stimulation. After intravenous administration, a pressor response occurs rapidly and reaches steady state within 5 minutes. The pharmacologic actions of norepinephrine are terminated primarily by uptake and metabolism in sympathetic nerve endings. The pressor action stops within 1 minutes-2 minutes after the infusion is discontinued.

12.3 Pharmacokinetics

Absorption

Following initiation of intravenous infusion, the steady state plasma concentration is achieved in 5 min.

Distribution

Plasma protein binding of norepinephrine is approximately 25%. It is mainly bound to plasma albumin and to a smaller extent to prealbumin and alpha 1-acid glycoprotein. The volume of distribution is 8.8 L. Norepinephrine localizes mainly in sympathetic nervous tissue. It crosses the placenta but not the blood-brain barrier.

Elimination

The mean half-life of norepinephrine is approximately 2.4 min. The average metabolic clearance is 3.1 L/min.

Metabolism

Norepinephrine is metabolized in the liver and other tissues by a combination of reactions involving the enzymes catechol-O-methyltransferase (COMT) and monoamine oxidase (MAO). The major metabolites are normetanephrine and 3-methoxy-4-hydroxy mandelic acid (vanillylmandelic acid, VMA), both of which are inactive. Other inactive metabolites include 3-methoxy-4-hydroxyphenylglycol, 3,4-dihydroxymandelic acid, and 3,4-dihydroxyphenylglycol.

Excretion

Norepinephrine metabolites are excreted in urine primarily as the sulfate conjugates and, to a lesser extent, as the glucuronide conjugates. Only small quantities of norepinephrine are excreted unchanged.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis, mutagenesis, and fertility studies have not been performed.

16 HOW SUPPLIED/STORAGE AND HANDLING

Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection for intravenous infusion is a colorless to slightly yellow solution available in single-dose, ready-to-use containers in an amber/foil overwrap. Each 250 mL of Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection, 4 mg/250 mL, 8 mg/250 mL, and 16 mg/250 mL contains the equivalent of 4 mg, 8 mg, and 16 mg of norepinephrine, respectively (provided as norepinephrine bitartrate monohydrate). Norepinephrine Bitartrate in Dextrose Injection or Norepinephrine Bitartrate in Sodium Chloride Injection is available in the following:

Norepinephrine Bitartrate in 5% Dextrose Injection	
Product Description	NDC Number
Twenty containers of 4 mg/250 mL (16 mcg/mL)	NDC 0338-0112-20
Twenty containers of 8 mg/250 mL (32 mcg/mL)	NDC 0338-0108-20
Twenty containers of 16 mg/250 mL (64 mcg/mL)	NDC 0338-0116-20

Norepinephrine Bitartrate in 0.9% Sodium Chloride Injection	
Product Description	NDC Number
Twenty containers of 4 mg/250 mL (16 mcg/mL)	NDC 0338-0040-20
Twenty containers of 8 mg/250 mL (32 mcg/mL)	NDC 0338-0042-20
Twenty containers of 16 mg/250 mL (64 mcg/mL)	NDC 0338-0128-20

Store at room temperature [20°C to 25°C (68°F to 77°F)], excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Once the overwrap is removed, the bag can be stored at room temperature for up to 30 days.

17 PATIENT COUNSELING INFORMATION

Risk of Tissue Damage

Advise the patient, family, or caregiver to report signs of extravasation urgently [see *Warnings and Precautions (5.1)*].

Manufactured by:
Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Baxter and Viaflo are trademarks of Baxter International, Inc. or its subsidiaries.

CB-30-03-122

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

EZPE7748

NDC 0338-0112-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

4 mg / 250 mL
(16 mcg / mL)

For Intravenous Infusion Only

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 1.6 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only – Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.

Rx Only

Baxter

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

Do not use
this port



CB-35-05-133

LOT

EXP

Container Label

EZPE7748

NDC 0338-0112-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

4 mg / 250 mL

(16 mg / mL)

For Intravenous Infusion Only

Each 100mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 1.6 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only – Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15° C to 30°C (59°F to 86° F) [see USP Controlled Room Temperature.] Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

VIAFLO container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

CB-35-05-133

VIAFLO container

Symbol

07

0

Symbol

Do not use

this port

Barcode

(01)00303380112203

LOT

EXP

EZPE7788

NDC 0338-0108-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

8 mg / 250 mL
(32 mcg / mL)

For Intravenous Infusion Only

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 3.2 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

Viaflo container is not made with natural rubber latex, DEHP, or PVC. Rx Only

Baxter

Baxter Healthcare Corporation
 Deerfield, IL 60015 USA
 Made in Ireland

Viaflo container

Do not use this port



(01)00303380108206

08-35-05-134

LOT

EXP

Container Label

EZPE7788

NDC 0338-0108-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

8 mg / 250 mL

(32 mcg / mL)

For Intravenous Infusion Only

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 3.2 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to

30°C (59°F to 86°F). [See USP Controlled Room Temperature.] Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

VIAFLO container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland

VIAFLO container

Symbol
Do not use
this port

Symbol
07
0

CB-35-05-134
Symbol
1

Barcode
(01)00303380108206

LOT

EXP

EZPE7758

NDC 0338-0116-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

16 mg / 250 mL
(64 mcg / mL)

For Intravenous Infusion Only

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 6.4 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland

Viaflo container

Do not use
this port



(0 1) 0 0 3 0 3 3 8 0 1 1 6 2 0 1

CB-35-05-144

LOT

EXP

Container Label

EZPE7758

NDC 0338-0116-20

250 mL

Norepinephrine

Bitartrate in 5% Dextrose Injection

16 mg / 250 mL

(64 mcg / mL)

For Intravenous Infusion Only

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 6.4 mg norepinephrine and 5 g Dextrose Monohydrate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to

30°C (59°F to 86°F). [See USP Controlled Room Temperature.] Protect from light. Keep in overwrap until ready to use. Once removed from overwrap, bag can be stored at room temperature and should be used within 30 days.

VIAFLO container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland

VIAFLO container

Symbol
Do not use
this port

Symbol
07
0

CB-35-05-144
Symbol
1

Barcode
(01)00303380116201

LOT

EXP

EZPE7764

NDC 0338-0040-01

250 mL

Norepinephrine

Bitartrate in 0.9% Sodium Chloride Injection

4 mg / 250 mL (16 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 1.6 mg norepinephrine, 0.9 g Sodium Chloride and 0.32 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container – Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

CB-35-05-683

Do not use
this port
LOT



(01)00303380040018

EXP

Container Label

EZPE7764

NDC 0338-0040-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

4 mg / 250 mL (16 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each **100** mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to **1.6** mg norepinephrine, **0.9** g Sodium Chloride and **0.32** mg Sodium Metabisulfate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only – Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

**Do not use
this port
Symbol**

Symbol

07

0

Barcode

(01)00303380040018

CB-35-05-683

LOT

EXP

EZPE7765

NDC 0338-0042-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

8 mg / 250 mL (32 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 3.2 mg norepinephrine, 0.9 g Sodium Chloride, 0.64 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container - Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container

CB-35-05-684

Do not use
this port



(0 1) 0 0 3 0 3 3 8 0 0 4 2 0 1 2

LOT

EXP

Container Label

EZPE7765

NDC 0338-0042-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

8 mg / 250 mL (32 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each **100** mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to **3.2** mg norepinephrine, **0.9** g Sodium Chloride, **0.64** mg Sodium Metabisulfate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.

Rx Only

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

**Do not use
this port
Symbol**

Symbol

07

0

CB-35-05-684

Barcode

(01)00303380042012

LOT

EXP

EZPE7766

NDC 0338-0128-01

250 mL

Norepinephrine

Bitartrate in 0.9% Sodium Chloride Injection

16 mg / 250 mL (64 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to 6.4 mg norepinephrine, 0.9 g Sodium Chloride, 1.28 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container - Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container

CB-35-05-685

Do not use
this port 
LOT



EXP

Container Label

EZPE7766

NDC 0338-0128-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

16 mg / 250 mL (64 mcg / mL)

For Intravenous Infusion Only

WARNING: CONTAINS SULFITES

Each **100** mL of sterile, nonpyrogenic solution contains: Norepinephrine Bitartrate Monohydrate USP equivalent to **6.4** mg norepinephrine, **0.9** g Sodium Chloride, **1.28** mg Sodium Metabisulfate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

**Do not use
this port
Symbol**

Symbol

07

0

CB-35-05-685

Barcode

(01)003033800128013

LOT

EXP

EZPE7764

NDC 0338-0040-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

4 mg / 250 mL (16 mcg / mL)



For Intravenous Infusion Only

(91)CB1001265

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 1.6 mg norepinephrine, 0.9 g Sodium Chloride, 0.32 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container - Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information. Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container



(0 1) 0 0 3 0 3 3 8 0 0 4 0 0 1 8

See solution container for lot and expiry.

CB-10-01-265

Un-varnished area (No text)

Overwrap Label

EZPE7764

NDC 0338-0040-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

4 mg / 250 mL (16 mcg / mL)

QR code

For Intravenous Infusion Only

(91)CB1001265

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 1.6 mg
norepinephrine, 0.9 g Sodium Chloride, 0.32 mg Sodium Metabisulfate
in Water for Injection, USP. May contain hydrochloric acid and/or
sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.

Recommended dosage: See prescribing information.

Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to
30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect
from light. Use within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.

Rx Only

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Made in Ireland

Viaflo container

Symbol

07

0

Barcode

(01)00303380040018

See solution container for lot and expiry.

CB-10-01-265

Un-varnished area (No text)

EZPE7765

NDC 0338-0042-01

250 mL

Norepinephrine

Bitartrate in 0.9% Sodium Chloride Injection

8 mg / 250 mL (32 mcg / mL)



For Intravenous Infusion Only

(91)CB1001266

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 3.2 mg norepinephrine, 0.9 g Sodium Chloride, 0.64 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container – Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information. Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC. Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container



(01)00303380042012

See solution container for lot and expiry.

CB-10-01-266

Un-varnished area (No text)

Overwrap Label

EZPE7765

NDC 0338-0042-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

8 mg / 250 mL (32 mcg / mL)

QR code

For Intravenous Infusion Only

(91)CB1001266

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 3.2 mg
norepinephrine, 0.9 g Sodium Chloride, 0.64 mg Sodium Metabisulfate
in Water for Injection, USP. May contain hydrochloric acid and/or
sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.
Recommended dosage: See prescribing information.
Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to
30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect
from light. Use within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland

Viaflo container

Symbol

07
0

Barcode
(01)00303380042012

See solution container for lot and expiry.

CB-10-01-266

Un-varnished area (No text)

EZPE7766

NDC 0338-0128-01

250 mL

Norepinephrine

Bitartrate in 0.9% Sodium Chloride Injection

16 mg / 250 mL (64 mcg / mL)



For Intravenous Infusion Only

(91)CB1001267

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 6.4 mg norepinephrine, 0.9 g Sodium Chloride, 1.28 mg Sodium Metabisulfite in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Container – Discard unused portion. For Intravenous Use. Recommended dosage: See prescribing information. Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use product within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container



(01)00303380128013

See solution container for lot and expiry.

CB-10-01-267

Un-varnished area (No text)

Overwrap Label

EZPE7766

NDC 0338-0128-01

250 mL

Norepinephrine

Bitartrate in **0.9% Sodium Chloride** Injection

16 mg / 250 mL (64 mcg / mL)

QR code

For Intravenous Infusion Only

(91)CB1001267

WARNING: CONTAINS SULFITES

Each 100 mL of sterile, nonpyrogenic solution contains:
Norepinephrine Bitartrate Monohydrate USP equivalent to 6.4 mg norepinephrine, 0.9 g Sodium Chloride, 1.28 mg Sodium Metabisulfate in Water for Injection, USP. May contain hydrochloric acid and/or sodium hydroxide for pH adjustment.

Single Dose Only - Discard unused portion. For Intravenous Use.
Recommended dosage: See prescribing information.
Store at 20°C to 25°C (68°F to 77°F), excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light. Use within 30 days of removal from overwrap.

Viaflo container is not made with natural rubber latex, DEHP, or PVC.
Rx Only

Baxter Logo

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Made in Ireland
Viaflo container

Symbol

07
0

Barcode
(01)00303380128013

See solution container for lot and expiry.

CB-10-01-267

Un-varnished area (No text)

NOREPINEPHRINE BITARTRATE			
norepinephrine bitartrate injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0112
Route of Administration	INTRAVENOUS		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)		NOREPINEPHRINE	4 mg in 250 mL

Inactive Ingredients

Ingredient Name	Strength
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	12.5 g in 250 mL
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0112-20	20 in 1 CARTON	01/15/2021	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214313	01/15/2021	

NOREPINEPHRINE BITARTRATE

norepinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0108
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)	NOREPINEPHRINE	8 mg in 250 mL

Inactive Ingredients

Ingredient Name	Strength
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	12.5 g in 250 mL
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0112-20	20 in 1 CARTON	01/15/2021	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

1	NDC:0338-0108-20	20 in 1 CARTON	01/15/2021	
1		250 mL in 1 CONTAINER; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NDA	NDA214313		01/15/2021	

NOREPINEPHRINE BITARTRATE				
norepinephrine bitartrate injection, solution				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0116	
Route of Administration	INTRAVENOUS			
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)			NOREPINEPHRINE	16 mg in 250 mL
Inactive Ingredients				
Ingredient Name			Strength	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)			12.5 g in 250 mL	
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
HYDROCHLORIC ACID (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0116-20	20 in 1 CARTON	11/21/2023	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NDA	NDA214313		11/21/2023	

NOREPINEPHRINE BITARTRATE

norepinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0040
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)	NOREPINEPHRINE	4 mg in 250 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	2.25 g in 250 mL
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	0.8 mg in 250 mL
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0040-20	20 in 1 CARTON	02/06/2026	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214313	11/21/2023	

NOREPINEPHRINE BITARTRATE

norepinephrine bitartrate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0042
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)			NOREPINEPHRINE	8 mg in 250 mL
Inactive Ingredients				
Ingredient Name			Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X)			2.25 g in 250 mL	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)			1.6 mg in 250 mL	
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
HYDROCHLORIC ACID (UNII: QTT17582CB)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0042-20	20 in 1 CARTON	02/06/2026	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NDA	NDA214313		11/21/2023	

NOREPINEPHRINE BITARTRATE			
norepinephrine bitartrate injection, solution			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-0128
Route of Administration	INTRAVENOUS		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NOREPINEPHRINE BITARTRATE (UNII: IFY5PE3ZRW) (NOREPINEPHRINE - UNII:X4W3ENH1CV)		NOREPINEPHRINE	16 mg in 250 mL
Inactive Ingredients			
Ingredient Name		Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X)		2.25 g in 250 mL	

SODIUM METABISULFITE (UNII: 4VON5FNS3C)	3.2 mg in 250 mL
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0128-20	20 in 1 CARTON	02/06/2026	
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214313	11/21/2023	

Labeler - Baxter Healthcare Corporation (005083209)

Revised: 2/2026

Baxter Healthcare Corporation