

IBUPROFEN- ibuprofen tablet, coated

Teeter

1112C-HTE-2025-0520

Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The

chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
 - take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, FD&C yellow #6, polyethylene glycol, polyvinyl alcohol, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Harris Teeter™

NDC 72036-212-02

Ibuprofen

†Compare to the Active Ingredient in Motrin® IB

Tablets, 200 mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

50 COATED CAPLETS**

**Capsule-Shaped Tablets

Actual Size

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Motrin® IB.

Drug Facts

Active ingredient (in each caplet) Purposes
Ibuprofen 200 mg (NSAID) ... Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ muscular aches ■ headache
 ■ toothache ■ minor pain of arthritis
 ■ the common cold ■ backache
 ■ menstrual cramps
 ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin redness ■ rash ■ dizziness
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs combining prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack.

Drug Facts (continued)

heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product: ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used
 adults and children 12 years and older ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years ■ ask a doctor

Other information



■ read all warnings and directions before use ■ store between 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

cellulose, silicon dioxide, corn starch, FD&C yellow #6, polyethylene glycol, polyvinyl alcohol, polydome K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

yourpharmsguarantee.
 Proudly Distributed By:
 HARRIS TETTER, LLC MATTHEWS, NC 28105
 Made in India
 1-800-432-6111 or harristetter.com

Important: Read all product information before using. Keep this box for important information.
Drug Facts (continued)
Questions or comments? 1-844-705-4384



IBUPROFEN ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72036-212-02	1 in 1 CARTON	12/13/2019	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:72036-212-03	1 in 1 CARTON	12/13/2019	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:72036-212-08	225 in 1 PACKAGE; Type 0: Not a Combination Product	12/13/2019	12/31/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202312	12/13/2019	

Labeler - Teeter (047279351)

