

CAPMIST DM- dextromethorphan hydrobromide, guaifenesin and pseudoephedrine hydrochloride tablet
Capital Pharmaceutical, LLC

Capmist DM Tablets

Drug Facts

Active ingredients

(in each tablet)

Dextromethorphan HBr 15mg

Purpose

Cough Suppressant

Active ingredients

(in each tablet)

Guaifenesin 400mg

Purpose

Expectorant

Active ingredients

(in each tablet)

Pseudoephedrine HCl 60mg

Purpose

Nasal Decongestant

Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- nasal congestion

- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson’s disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlargement of the prostate gland

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 4 tablets in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 2 tablets in 24 hours, or as directed by a doctor.
Children 2 to under 6 years of	Consult a doctor

age:

CONSULT A DOCTOR.

Other information

Store at 15°-30°C (59°-86°F).

Capmist DM is a light green, oblong, capsule shaped, scored tablet, debossed "CAP" bisect "DM" on one side and blank on the other.

Contains color additives including FD&C Yellow No. 5 (tartrazine).

Supplied in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients

FD&C Blue #1, FD&C Yellow #5, Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate.

PRINCIPAL DISPLAY PANEL


NDC 29978-601-90

CAPMIST DM Tablets

Cough Suppressant ● Expectorant

Nasal Decongestant

90 tablets



Drug Facts (continued)

Other information
 Store at 15°-30°C (59°-86°F).

Capmist DM is a light green, oblong, capsule shaped, scored tablet, debossed "CAP" bisect "DM" on one side and blank on the other.

Contains color additives including FD&C Yellow No. 5 (tartrazine).

Supplied in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients
 FD&C Blue # 1, FD&C Yellow # 5, Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate.

Drug Facts

Active ingredients (in each tablet)	Purposes
Dextromethorphan HBr 15 mg ..	Cough Suppressant
Guaifenesin 400 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

Uses
 Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- nasal congestion
- reduces swelling of nasal passages

▲ Lift Here

Drug Facts (continued)

Warnings
Do not exceed recommended dosage.
Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- heart disease ■ high blood pressure ■ thyroid disease
- diabetes ■ trouble urinating due to enlargement of the prostate gland

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Do not exceed recommended dosage.

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 4 tablets in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 2 tablets in 24 hours, or as directed by a doctor.
Children 2 to under 6 years of age:	Consult a doctor.

4.3125"

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	green (light green)	Score	2 pieces
Shape	CAPSULE (oblong)	Size	19mm
Flavor		Imprint Code	CAP;DM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29978-601-30	2 in 1 BOX	06/20/2013	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:29978-601-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	06/20/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/20/2013	

Labeler - Capital Pharmaceutical, LLC (831545541)