

VIJOICE- alpelisib tablet
VIJOICE- alpelisib
VIJOICE- alpelisib granule
Novartis Pharmaceuticals Corporation

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VIJOICE safely and effectively. See full prescribing information for VIJOICE.

VIJOICE® (alpelisib) tablets, for oral use

VIJOICE® (alpelisib) oral granules

Initial U.S. Approval: 2019

RECENT MAJOR CHANGES

Dosage and Administration, VIJOICE Preparation and Administration Instructions (2.4)	7/2025
Warnings and Precautions, Severe Hypersensitivity (5.1)	7/2025

INDICATIONS AND USAGE

VIJOICE is a kinase inhibitor indicated for the treatment of adult and pediatric patients 2 years of age and older with severe manifestations of PIK3CA-Related Overgrowth Spectrum (PROS) who require systemic therapy.

This indication is approved under accelerated approval based on response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s). (1)

DOSAGE AND ADMINISTRATION

Recommended Dose:

- Pediatric patients (2 to less than 18 years of age): 50 mg taken orally once daily with food. (2.1)
- Adult patients: 250 mg taken orally once daily with food. (2.1)

Administration:

- Swallow tablets whole. Do not chew, divide, or crush. (2.4)
- For patients who cannot swallow tablets whole, use tablets or oral granules to create a suspension or a mixture. (2.4)
- VIJOICE suspension made with water can be administered orally or via a nasogastric or gastric tube. (2.4)

See full prescribing information for preparation and administration instructions.

DOSAGE FORMS AND STRENGTHS

Tablets: 50 mg, 125 mg, and 200 mg (3)

Oral Granules: 50 mg (3)

CONTRAINDICATIONS

Severe hypersensitivity to VIJOICE or to any of its ingredients. (4)

WARNINGS AND PRECAUTIONS

- **Severe Hypersensitivity:** Permanently discontinue VIJOICE. Promptly initiate appropriate treatment. (5.1)
- **Severe Cutaneous Adverse Reactions (SCARs):** VIJOICE can cause SCARs, including Stevens-Johnson syndrome (SJS), erythema multiforme (EM), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS). Interrupt VIJOICE for signs or symptoms of SCARs. Permanently discontinue VIJOICE if SCARs are confirmed. (2.5, 5.2)
- **Hyperglycemia:** VIJOICE can cause severe hyperglycemia, in some cases associated with hyperglycemic hyperosmolar non-ketotic syndrome (HHNKS) or ketoacidosis. The safety of VIJOICE in patients with Type 1 or uncontrolled Type 2 diabetes has not been established. Before initiating treatment with VIJOICE, test fasting plasma glucose (FPG), HbA1c, and optimize blood glucose. After initiating treatment, monitor periodically. Initiate or optimize anti-hyperglycemic medications as clinically indicated. Interrupt, reduce dose, or discontinue VIJOICE if severe hyperglycemia occurs. (2.5, 5.3)
- **Pneumonitis:** VIJOICE can cause severe pneumonitis and interstitial lung disease. Monitor for clinical symptoms or radiological changes. Permanently discontinue VIJOICE if pneumonitis occurs. (2.5, 5.4)

- **Diarrhea or Colitis:** VIJOICE can cause severe diarrhea, resulting in dehydration and in some cases in acute kidney injury, as well as colitis. Monitor for diarrhea and additional symptoms of colitis, including abdominal pain and mucus or blood in stool. Interrupt, reduce dose, or permanently discontinue VIJOICE based on severity of diarrhea or colitis. Patients with colitis may require additional treatment, such as enteric-acting and/or systemic steroids. (2.5, 5.5)
- **Embryo-Fetal Toxicity:** VIJOICE can cause fetal harm. Advise patients of potential risk to a fetus and to use effective contraception. (5.6, 8.1, 8.3)

-----**ADVERSE REACTIONS**-----

Most common adverse reactions (Grades 1 to 4, incidence ≥ 10%) were diarrhea, stomatitis, and hyperglycemia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Novartis Pharmaceuticals Corporation at 1-888-669-6682 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----**DRUG INTERACTIONS**-----

- **CYP3A4 Inducers:** Avoid coadministration of VIJOICE with a strong CYP3A4 inducer. Consider an alternative concomitant drug with no or minimal potential to induce CYP3A4. (7.1)
- **Breast Cancer Resistance Protein (BCRP) Inhibitors:** Avoid the use of BCRP inhibitors in patients treated with VIJOICE. If unable to use alternative drugs, closely monitor for increased adverse reactions. (7.1)

-----**USE IN SPECIFIC POPULATIONS**-----

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 7/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 VIJOICE Dosage Form Overview
- 2.3 VIJOICE Administration Overview
- 2.4 VIJOICE Preparation and Administration Instructions
- 2.5 Dosage Modifications for Adverse Reactions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Severe Hypersensitivity
- 5.2 Severe Cutaneous Adverse Reactions
- 5.3 Hyperglycemia
- 5.4 Pneumonitis
- 5.5 Diarrhea or Colitis
- 5.6 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience
- 6.2 Postmarketing Experience and Other Spontaneous Adverse Reaction Reports

7 DRUG INTERACTIONS

- 7.1 Effect of Other Drugs on VIJOICE

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation

- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 10 OVERDOSAGE**
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

VIJOICE is indicated for the treatment of adult and pediatric patients 2 years of age and older with severe manifestations of PIK3CA-Related Overgrowth Spectrum (PROS) who require systemic therapy.

This indication is approved under accelerated approval based on response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Adult Patients

The recommended dosage of VIJOICE in adult patients is 250 mg orally, once daily, administered as recommended [see *Dosage and Administration* (2.2, 2.3, 2.4)] until disease progression or unacceptable toxicity.

Pediatric Patients (2 to less than 18 years of age)

The recommended initial dosage of VIJOICE in pediatric patients is 50 mg orally, once daily, administered as recommended [see *Dosage and Administration* (2.2, 2.3, 2.4)] until disease progression or unacceptable toxicity.

Consider a dose increase to 125 mg once daily in pediatric patients \geq 6 years old for response optimization (clinical/radiological) after 24 weeks of treatment with VIJOICE at 50 mg once daily. When a pediatric patient turns 18 years old, consider a gradual dose increase up to 250 mg. Recommended dose increases by age group are listed in Table

1.

Table 1: Recommended Daily VIJOICE Dose Levels for Pediatric Patients (2 to less than 18 years of age)

Patient age (years)	Initial dose	Dose increase
2 to < 6	50 mg ^a	Not applicable ^b
6 to < 18	50 mg ^a	125 mg

^aDose can be administered as VIJOICE tablets or VIJOICE oral granules.

^bA recommended increased dose has not been established.

2.2 VIJOICE Dosage Form Overview

VIJOICE is available in two dosage forms: tablets and oral granules.

Prescribe the most appropriate dosage form of VIJOICE according to the dose required and patient needs.

VIJOICE Tablets (50 mg, 125 mg, and 200 mg) may be administered as:

- Whole tablets: For patients who can swallow whole tablets.
- Tablets prepared as an oral suspension: For patients who have difficulty swallowing whole tablets [see *Dosage and Administration (2.4)*].

VIJOICE Oral Granules (50 mg per packet):

- For patients who are prescribed a 50 mg daily dose only [see *Dosage and Administration (2.1)*].
- Do not use multiple 50 mg packets or a partial packet of oral granules for patients prescribed a 125 mg or a 250 mg dose [see *Clinical Pharmacology (12.3)*].
- Do not combine VIJOICE tablets and VIJOICE oral granules to achieve the prescribed dose.

2.3 VIJOICE Administration Overview

Take VIJOICE with food at approximately the same time each day [see *Clinical Pharmacology (12.3)*].

If a dose of VIJOICE is missed, it can be taken with food within 9 hours after the time it is usually taken. After more than 9 hours, skip the dose for that day. The next day, take VIJOICE at the usual time.

If the patient vomits after taking the dose, advise the patient not to take an additional dose on that day, and to resume the dosing schedule the next day at the usual time.

2.4 VIJOICE Preparation and Administration Instructions

VIJOICE Tablets

[Swallow VIJOICE tablets whole, or prepare as a suspension to administer orally, or via

feeding tubes.

VIJOICE Tablets, Whole

- Swallow VIJOICE tablets whole and take with food. Do not chew, divide, or crush.
- Do not use broken, cracked, or damaged tablets.

VIJOICE Tablets Prepared as a Suspension for Oral use, or Feeding Tubes (Nasogastric or Gastric Tube) Administration

- For patients who are not able to swallow tablets, or who are using a feeding tube, prepare and administer VIJOICE as a suspension and take with food [see *Clinical Pharmacology (12.3)*].
 - To prepare VIJOICE tablets as a **suspension for oral use**, place VIJOICE tablets in a cup containing 2 to 4 ounces of water. To prepare VIJOICE tablets as a **suspension for feeding tubes administration**, place VIJOICE tablets in a cup containing 1 to 2 ounces of water. Make the suspension with water only.
 - Let tablets stand in water for approximately 5 minutes.
 - Crush the tablets with a spoon and stir until a suspension is obtained.
 - Immediately after preparation, administer the suspension as directed below. If not administered immediately after preparation, stir the suspension with the same spoon to re-suspend any particles before administration.
 - Discard the suspension if it is not administered within 60 minutes after preparation.

Oral Administration

- Administer the suspension from the cup. After administration of the suspension, add approximately 1 to 2 ounces of water to the same cup. Stir with the same spoon to re-suspend any remaining particles and administer the entire contents of the cup orally. Repeat if particles remain.

Feeding Tubes Administration

- Administer VIJOICE tablets prepared as a suspension via French size 5 to 12 diameter silicone or polyurethane nasogastric tubes, or via French size 12 to 24 diameter silicone gastric tubes.
- Withdraw VIJOICE suspension from the cup into an enteral syringe and administer it via the nasogastric or gastric tube.
- After administration, add approximately 1 to 2 ounces (approximately 30 to 60 mL) of water to the same cup. Stir with the same spoon to re-suspend any remaining particles. Withdraw the contents of the cup into the same enteral syringe and administer it via the nasogastric or gastric tube. Repeat if particles remain.

VIJOICE Oral Granules

Administer VIJOICE oral granules directly onto the tongue with water, or prepare as a suspension or a mixture for oral use. To administer via feeding tubes prepare the suspension with water only.

- Each packet is for single use only.
- No packet should be used if the packet seal is broken.
- Do not attempt to use partial quantities of oral granules from 50 mg granules packets to prepare a dose. Do not combine VIJOICE tablets and VIJOICE oral granules to achieve the prescribed dose of 125 mg or 250 mg.
- For patients for whom a daily dose of 50 mg is prescribed, administer VIJOICE oral

granules [see *Clinical Pharmacology (12.3)*] in one of the following ways:

VIJOICE Oral Granules for Direct Oral Administration

- Pour the contents of one VIJOICE oral granules packet directly onto the tongue and swallow with approximately 2 to 4 ounces of water. If needed, rinse the mouth with additional water and swallow to ensure no particles remain in the mouth.

VIJOICE Oral Granules as a Suspension or a Mixture for Oral administration

- Pour the contents of one VIJOICE oral granules packet into a cup.
- Add 1 to 3 teaspoons (about 0.5 ounces) of a beverage (water, milk, or apple juice) or soft food (applesauce or yogurt) and stir with a spoon, then administer the suspension or the mixture immediately.
- Rinse the cup with up to 2 ounces of a beverage (water, milk or apple juice) and administer the rinse immediately to ensure the entire dose is administered. If particles remain, repeat until the full dose is administered.
- If not administered immediately after preparation, stir the suspension or the mixture with the same spoon to re-suspend any particles before administration.
- Discard the oral granules mixed with water, milk, apple juice, applesauce, or yogurt if they are not administered within 60 minutes after preparation.

VIJOICE Oral Granules Suspension for Feeding Tubes Administration

For patients who are not able to swallow VIJOICE orally, administer VIJOICE via feeding tubes.

Administer VIJOICE granules via French size 8 to 12 diameter silicone or polyurethane nasogastric tubes or via French size 12 to 24 diameter silicone gastric tubes.

- Pour the contents of one VIJOICE granules packet into a cup.
- Add 4 teaspoons (about 0.7 ounces or 20 mL) of water and stir gently with a spoon until a suspension is obtained. Make the suspension with water only.
- Immediately after preparation, withdraw the suspension from the cup into an enteral syringe and administer it via the nasogastric or gastric tube.
- After administration, add 4 teaspoons (about 0.7 ounces or 20 mL) of water to the same cup. Stir with the same spoon to re-suspend any remaining particles. Withdraw the contents of the cup into the same enteral syringe and administer it via the nasogastric or gastric tube. Repeat if particles remain.
- If not administered immediately after preparation, stir the suspension with the same spoon to re-suspend any particles before withdrawing into an enteral syringe for administration.
- Discard the suspension if it is not administered within 60 minutes after preparation.

2.5 Dosage Modifications for Adverse Reactions

The recommended VIJOICE dose reductions for adverse reactions in adult and pediatric patients are listed in Table 2 and Table 3, respectively.

Table 2: VIJOICE Dosage Reduction Recommendations for Adverse Reactions in Adult Patients

VIJOICE dose level	Dose and schedule
First-dose reduction	125 mg once daily
Second-dose reduction	50 mg once daily ^a

^aDose can be administered as VIJOICE tablets or VIJOICE oral granules.

Table 3: VIJOICE Dosage Reduction Recommendations for Adverse Reactions in Pediatric Patients

Action	VIJOICE dose prior to dose reduction	
	125 mg once daily	50 mg once daily
Dose reduction	50 mg once daily ^a	Not applicable

^aDose can be administered as VIJOICE tablets or VIJOICE oral granules.

Discontinue VIJOICE in adults or pediatric patients who cannot tolerate 50 mg daily.

Tables 4, 5, 6, 7, 8, and 9 summarize recommendations for dose interruption, reduction, or discontinuation of VIJOICE in the management of specific adverse reactions.

Cutaneous Adverse Reactions

If a severe cutaneous adverse reaction (SCAR) is confirmed, permanently discontinue VIJOICE. Do not reintroduce VIJOICE in patients who have experienced previous SCAR during VIJOICE treatment [see *Warnings and Precautions (5.2)*].

Table 4: Dosage Modification and Management for Rash and Severe Cutaneous Adverse Reactions (SCARs)

[see *Warnings and Precautions (5.1, 5.2)*]

Grade^{a,b}	Recommendation for adult and pediatric patients^c
Grade 1 (< 10% body surface area [BSA] with active skin toxicity)	No VIJOICE dosage modification is required unless the etiology is determined to be SCAR. Initiate topical corticosteroid treatment. Consider adding oral antihistamine to manage symptoms. If active rash is not improved within 28 days of appropriate treatment, add a low dose systemic corticosteroid. If the etiology is determined to be SCAR,

Grade 2 (10% to 30% BSA with active skin toxicity)	<p>permanently discontinue VIJOICE.</p> <p>No VIJOICE dosage modification is required unless the etiology is determined to be SCAR.</p> <p>Initiate or intensify topical corticosteroid and oral antihistamine treatment.</p> <p>Consider low dose systemic corticosteroid treatment.</p> <p>If rash improves to Grade ≤ 1 within 10 days, systemic corticosteroid may be discontinued.</p> <p>If the etiology is determined to be SCAR, permanently discontinue VIJOICE.</p>
Grade 3 (e.g., severe rash not responsive to medical management) (> 30% BSA with active skin toxicity)	<p>Interrupt VIJOICE and initiate or intensify topical/systemic corticosteroid and oral antihistamine treatment.</p> <p>If the etiology is determined to be SCAR, permanently discontinue VIJOICE.</p> <p>For rashes other than SCAR</p> <p><u>Adult Patients:</u></p> <ul style="list-style-type: none"> • Upon improvement to Grade ≤ 1, resume VIJOICE at the next lower dose level. <p><u>Pediatric Patients:</u></p> <ul style="list-style-type: none"> • Upon improvement to Grade ≤ 1, either resume VIJOICE at 50 mg while continuing oral antihistamine treatment or permanently discontinue VIJOICE. • Permanently discontinue VIJOICE if: <ul style="list-style-type: none"> ◦ Patient was receiving antihistamines at the time of rash onset and antihistamine dose cannot be increased ◦ Grade ≥ 3 rash recurs

Grade 4 (e.g., severe bullous, blistering or exfoliating skin conditions) (any % BSA associated with extensive superinfection, with IV antibiotics indicated; life-threatening consequences)	Permanently discontinue VIJOICE.
---	----------------------------------

^aGrading according to Common Terminology Criteria for Adverse Events (CTCAE) Version 5.0.

^bFor all grades of rash, consider consultation with a dermatologist.

^cAntihistamines administered prior to rash onset may decrease incidence and severity of rash.

Hyperglycemia

Before initiating treatment with VIJOICE, test fasting plasma glucose (FPG), HbA1c, and optimize blood glucose. After initiating treatment with VIJOICE, monitor fasting glucose (FPG or fasting blood glucose) at least once every week for the first 2 weeks, then at

least once every 4 weeks, and as clinically indicated. Monitor HbA1c every 3 months and as clinically indicated. In patients with risk factors for hyperglycemia, monitor fasting glucose more closely and as clinically indicated [see *Warnings and Precautions (5.3)*].

Table 5: Dosage Modification and Management for Hyperglycemia

[see *Warnings and Precautions (5.3)*]

Fasting plasma glucose (FPG)/Fasting blood glucose values^a	Recommendation for adult and pediatric patients
Dose modifications and management should only be based on fasting glucose values (FPG or fasting blood glucose).	
Grade 1 Fasting glucose > ULN -160 mg/dL or > ULN -8.9 mmol/L	No VIJOICE dosage modification is required. Initiate or intensify oral anti-hyperglycemic treatment ^b .
Grade 2 Fasting glucose > 160 - 250 mg/dL or > 8.9 - 13.9 mmol/L	No VIJOICE dosage modification is required. Initiate or intensify oral anti-hyperglycemic treatment ^b .
	<p><u>Adult Patients:</u></p> <ul style="list-style-type: none"> If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 21 days under appropriate anti-hyperglycemic treatment^b, reduce VIJOICE dose by 1 dose level and follow fasting glucose value-specific recommendations.
	<p><u>Pediatric Patients:</u></p> <ul style="list-style-type: none"> If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 21 days under appropriate anti-hyperglycemic treatment^b, interrupt VIJOICE until improvement to Grade ≤ 1, then resume VIJOICE at 50 mg and follow fasting glucose value-specific recommendations.
Grade 3 Fasting glucose > 250 - 500 mg/dL or > 13.9 - 27.8 mmol/L	<p>Interrupt VIJOICE. Initiate or intensify oral anti-hyperglycemic treatment^b and consider additional anti-hyperglycemic medications for 1-2 days until hyperglycemia improves, as clinically indicated. Administer intravenous hydration and consider appropriate treatment (e.g., intervention for electrolyte/ketoacidosis/hyperosmolar disturbances).</p> <p><u>Adult Patients:</u></p> <ul style="list-style-type: none"> If fasting glucose decreases to ≤ 160 mg/dL or 8.9 mmol/L within 3 to 5 days

under appropriate anti-hyperglycemic treatment, resume VIJOICE at 1 lower dose level.

- If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 3 to 5 days under appropriate anti-hyperglycemic treatment, consultation with a physician with expertise in the treatment of hyperglycemia is recommended.
- If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 21 days following appropriate anti-hyperglycemic treatment^b, permanently discontinue VIJOICE.

Pediatric Patients:

- If fasting glucose decreases to ≤ 160 mg/dL or 8.9 mmol/L within 3 to 5 days under appropriate anti-hyperglycemic treatment, resume VIJOICE at 50 mg.
- If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 3 to 5 days under appropriate anti-hyperglycemic treatment, consultation with a physician with expertise in the treatment of hyperglycemia is recommended to determine if treatment with VIJOICE should be resumed or permanently discontinued.
- If fasting glucose does not decrease to ≤ 160 mg/dL or 8.9 mmol/L within 21 days following appropriate anti-hyperglycemic treatment^b, permanently discontinue VIJOICE.
- If hyperglycemia recurs at Grade ≥ 3 , consider permanent discontinuation of VIJOICE.

Grade 4
Fasting glucose > 500 mg/dL or > 27.8 mmol/L

Interrupt VIJOICE.
Initiate or intensify appropriate oral anti-hyperglycemic treatment^b.
Administer intravenous hydration and consider appropriate treatment (e.g., intervention for electrolyte/ketoacidosis/hyperosmolar disturbances).
Re-check fasting glucose within 24 hours and as clinically indicated.

- If fasting glucose decreases to ≤ 500

mg/dL or 27.8 mmol/L, follow fasting glucose value-specific recommendations for Grade 3.

- If fasting glucose is confirmed at > 500 mg/dL or 27.8 mmol/L, permanently discontinue VIJOICE.

Abbreviation: ULN, upper limit of normal.

^aFPG/Fasting Blood Glucose/Grade levels reflect hyperglycemia grading according to Common Terminology Criteria for Adverse Events (CTCAE) Version 4.03.

^bInitiate applicable anti-hyperglycemic medications, including metformin in adult and pediatric patients ≥ 10 years, SGLT2 inhibitors or insulin sensitizers (such as thiazolidinediones or dipeptidyl peptidase-4 inhibitors) in adult patients, and review respective prescribing information for dosing and dose titration recommendations, including local hyperglycemic treatment guidelines [see *Warnings and Precautions (5.3)*].

Pneumonitis

Table 6: Dosage Modification for Pneumonitis

[see *Warnings and Precautions (5.4)*]

Grade^a	Recommendation for adult and pediatric patients
--------------------------	--

Any Grade	<ul style="list-style-type: none">• Interrupt VIJOICE if pneumonitis is suspected.• Permanently discontinue VIJOICE if pneumonitis is confirmed.
-----------	---

^aGrading according to CTCAE Version 5.0.

Diarrhea or Colitis

In pediatric patients, consider consultation with a physician with experience in the treatment of gastrointestinal conditions.

Table 7: Dosage Modification and Management for Diarrhea or Colitis

[see *Warnings and Precautions (5.5)*]

Grade^a	Recommendation for adult and pediatric patients
--------------------------	--

Grade 1	No VIJOICE dosage modification is required. Initiate appropriate medical therapy and monitor as clinically indicated.
---------	---

Grade 2	Interrupt VIJOICE dose until improvement to Grade ≤ 1 , then resume VIJOICE at the same dose level. Initiate or intensify appropriate medical therapy
---------	--

and monitor as clinically indicated^b.

Adult Patients:

- For recurrent Grade ≥ 2 , interrupt VIJOICE dose until improvement to Grade ≤ 1 , then resume VIJOICE at the next lower dose level.

Pediatric Patients:

- For recurrent Grade ≥ 2 , interrupt VIJOICE dose until improvement to Grade ≤ 1 , then resume VIJOICE at 50 mg.

Grade 3	Interrupt VIJOICE dose until improvement to Grade ≤ 1 . Initiate or intensify appropriate medical therapy and monitor as clinically indicated ^b . <u>Adult Patients:</u> <ul style="list-style-type: none">• Once improved to Grade ≤ 1, then resume VIJOICE at the next lower dose level. <u>Pediatric Patients:</u> <ul style="list-style-type: none">• Once improved to Grade ≤ 1, either resume VIJOICE at 50 mg or permanently discontinue VIJOICE.• For recurrent Grade ≥ 3, consider permanent discontinuation of VIJOICE.
---------	---

Grade 4	Permanently discontinue VIJOICE.
---------	----------------------------------

^aGrading according to CTCAE Version 5.0.

^bFor Grade 2 and 3 colitis consider additional treatment, such as enteric-acting and/or systemic steroids.

Pancreatitis

Table 8: Dosage Modification for Pancreatitis

Grade^a	Recommendation for adult and pediatric patients
Grade 2	Interrupt VIJOICE dose until improvement to Grade < 2 . <u>Adult Patients:</u> <ul style="list-style-type: none">• Resume VIJOICE at the next lower dose level (only one dose reduction is permitted).• If pancreatitis recurs, permanently discontinue VIJOICE. <u>Pediatric Patients:</u> <ul style="list-style-type: none">• Resume VIJOICE at 50 mg.• If pancreatitis recurs, permanently discontinue VIJOICE.

Grade 3	<p><u>Adult Patients:</u></p> <ul style="list-style-type: none"> • Interrupt VIJOICE dose until improvement to Grade < 2. • Resume VIJOICE at the next lower dose level (only one dose reduction is permitted). • If pancreatitis recurs, permanently discontinue VIJOICE. <p><u>Pediatric Patients:</u></p> <ul style="list-style-type: none"> • Permanently discontinue VIJOICE.
---------	---

Grade 4 Permanently discontinue VIJOICE.

^aGrading according to CTCAE Version 5.0.

Other Adverse Reactions

Table 9: Dosage Modification and Management for Other Adverse Reactions (Excluding Rash and Severe Cutaneous Adverse Reactions, Hyperglycemia, Pneumonitis, Diarrhea or Colitis, and Pancreatitis)

Grade^a	Recommendation for adult and pediatric patients
Grade 1 or 2 ^{b,c,d}	No VIJOICE dosage modification is required. Initiate appropriate medical therapy and monitor as clinically indicated ^{b,c,d} .
Grade 3	<p>Interrupt VIJOICE dose until improvement to Grade ≤ 1. Initiate or intensify appropriate medical therapy and monitor as clinically indicated.</p> <p><u>Adult Patients:</u></p> <ul style="list-style-type: none"> • Once improved to Grade ≤ 1, then resume VIJOICE at the next lower dose level. <p><u>Pediatric Patients:</u></p> <ul style="list-style-type: none"> • Once improved to Grade ≤ 1, either resume VIJOICE at 50 mg or permanently discontinue VIJOICE. • If adverse reaction recurs at Grade ≥ 3, consider permanent discontinuation of VIJOICE. • Consider consultation with a qualified physician with specific expertise in the field of the concerned adverse reaction.
Grade 4	Permanently discontinue VIJOICE.

^aGrading according to CTCAE Version 5.0.

^bFor Grade 2 total bilirubin elevation in adult patients, interrupt VIJOICE dose until improvement to Grade ≤ 1. If improvement

occurs in ≤ 14 days, resume at the same dose level. If improvement occurs in > 14 days, resume VIJOICE at the next lower dose level.

^cFor Grade 2 total bilirubin elevation in pediatric patients, interrupt VIJOICE dose until improvement to Grade ≤ 1 . If improvement occurs in ≤ 14 days, resume at the same dose level. If improvement occurs in > 14 days, resume VIJOICE at 50 mg.

^dIf alopecia becomes a concern, consider consulting a dermatologist.

3 DOSAGE FORMS AND STRENGTHS

Tablets

50 mg: Light yellow, unscored, round and curved with beveled edges film-coated tablet, debossed with “C7” on one side and “NVR” on the other side.

125 mg: Dark yellow, unscored, ovaloid and curved with beveled edges film-coated tablet, debossed with “Y7” on one side and “NVR” on the other side.

200 mg: Pale yellow, unscored, ovaloid and curved with beveled edges film-coated tablet, debossed with “CL7” on one side and “NVR” on the other side.

Oral Granules

50 mg: White to almost white free flowing mixture of powder and granules in packets.

4 CONTRAINDICATIONS

VIJOICE is contraindicated in patients with severe hypersensitivity to alpelisib or any of its ingredients [see *Warnings and Precautions (5.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Severe Hypersensitivity

Severe hypersensitivity reactions, including anaphylaxis, anaphylactic shock, and angioedema, have occurred in adult patients treated with alpelisib in the oncology setting and in patients treated with VIJOICE [see *Adverse Reactions (6.1)*]. VIJOICE is not approved for use in the oncology setting.

Permanently discontinue VIJOICE in the event of severe hypersensitivity [see *Contraindications (4)*].

5.2 Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), erythema multiforme (EM), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS), have occurred in adult patients treated with alpelisib in the oncology setting and may occur in patients treated with VIJOICE. VIJOICE is not approved for use in the oncology setting.

If signs or symptoms of SCARs occur, interrupt VIJOICE until the etiology of the reaction

has been determined. Consultation with a dermatologist is recommended.

If a SCAR is confirmed, permanently discontinue VIJOICE.

If a SCAR is not confirmed, VIJOICE may require dose modifications, topical or systemic corticosteroids, or oral antihistamine treatment as described in Table 4 [see *Dosage and Administration (2.5)*].

5.3 Hyperglycemia

Severe hyperglycemia, in some cases associated with hyperglycemic hyperosmolar non-ketotic syndrome (HHNKS) or fatal cases of ketoacidosis, has occurred in adult patients treated with alpelisib in the oncology setting and may occur in patients treated with VIJOICE. VIJOICE is not approved for use in the oncology setting.

In the EPIK-P1 study, Grade 1 or 2 hyperglycemia was reported in 12% of patients treated with VIJOICE [see *Adverse Reactions (6.1)*].

Before initiating treatment with VIJOICE, test fasting plasma glucose (FPG), HbA1c, and optimize blood glucose. After initiating treatment with VIJOICE, monitor fasting glucose (FPG or fasting blood glucose) at least once every week for the first 2 weeks, then at least once every 4 weeks, and as clinically indicated. Monitor HbA1c every 3 months and as clinically indicated. Monitor fasting glucose more frequently for the first few weeks during treatment with VIJOICE in patients with risk factors for hyperglycemia, such as obesity (BMI \geq 30), elevated FPG, HbA1c at the upper limit of normal or above, use of concomitant systemic corticosteroids, or age \geq 75 [see *Use in Specific Populations (8.5)*].

If a patient experiences hyperglycemia after initiating treatment with VIJOICE, monitor fasting glucose as clinically indicated, and at least twice weekly until fasting glucose decreases to normal levels. During treatment with anti-hyperglycemic medication, continue monitoring fasting glucose at least once a week for 8 weeks, followed by once every 2 weeks and as clinically indicated. Consider consultation with a healthcare practitioner with expertise in the treatment of hyperglycemia and counsel patients on lifestyle changes.

The safety of VIJOICE in patients with Type 1 and uncontrolled Type 2 diabetes has not been established. Patients with a history of diabetes mellitus may require intensified hyperglycemic treatment. Closely monitor patients with diabetes.

Interrupt, reduce the dose, or permanently discontinue VIJOICE based on the severity as described in Table 5 [see *Dosage and Administration (2.5)*].

5.4 Pneumonitis

Severe pneumonitis, including acute interstitial pneumonitis and interstitial lung disease, has occurred in adult patients treated with alpelisib in the oncology setting and may occur in patients treated with VIJOICE. VIJOICE is not approved for use in the oncology setting.

In patients who have new or worsening respiratory symptoms or are suspected to have developed pneumonitis, interrupt VIJOICE immediately and evaluate the patient for pneumonitis. Consider a diagnosis of non-infectious pneumonitis in patients presenting with non-specific respiratory signs and symptoms, such as hypoxia, cough, dyspnea, or interstitial infiltrates on radiologic exams and in whom infectious, neoplastic, and other

causes have been excluded by means of appropriate investigations.

Permanently discontinue VIJOICE in all patients with confirmed pneumonitis [see *Dosage and Administration (2.5)*].

5.5 Diarrhea or Colitis

Severe diarrhea, resulting in dehydration and in some cases in acute kidney injury, as well as colitis, have occurred in adult patients treated with alpelisib in the oncology setting and may occur in patients treated with VIJOICE. VIJOICE is not approved for use in the oncology setting.

In the EPIK-P1 study, 16% of patients experienced Grade 1 diarrhea during treatment with VIJOICE [see *Adverse Reactions (6.1)*].

Monitor patients for diarrhea and additional symptoms of colitis, such as abdominal pain and mucus or blood in stool. Interrupt, reduce the dose or permanently discontinue VIJOICE based on the severity of diarrhea or colitis [see *Dosage and Administration (2.5)*].

Patients with colitis may require additional treatment, such as enteric-acting and/or systemic steroids [see *Dosage and Administration (2.5)*].

5.6 Embryo-Fetal Toxicity

Based on findings in animals and its mechanism of action, VIJOICE can cause fetal harm when administered to a pregnant woman. In animal reproduction studies, oral administration of alpelisib to pregnant animals during organogenesis caused adverse developmental outcomes, including embryo-fetal mortality (post-implantation loss), reduced fetal weights, and increased incidences of fetal malformations at doses that were approximately equivalent to the recommended pediatric and adult doses. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with VIJOICE and for 1 week after the last dose. Advise male patients with female partners of reproductive potential to use condoms and effective contraception during treatment with VIJOICE and for 1 week after the last dose [see *Use in Specific Populations (8.1, 8.3)* and *Clinical Pharmacology (12.1)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are discussed elsewhere in the labeling:

- Severe Hypersensitivity [see *Warnings and Precautions (5.1)*]
- Severe Cutaneous Adverse Reactions [see *Warnings and Precautions (5.2)*]
- Hyperglycemia [see *Warnings and Precautions (5.3)*]
- Pneumonitis [see *Warnings and Precautions (5.4)*]
- Diarrhea or Colitis [see *Warnings and Precautions (5.5)*]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of VIJOICE was evaluated in EPIK-P1 (NCT04285723), a single-arm clinical study in patients who were treated as part of an expanded access program for compassionate use. Fifty-seven patients 2 years of age and older with severe or life-threatening PIK3CA-Related Overgrowth Spectrum (PROS) received VIJOICE based on age at dosages ranging from 50 mg to 250 mg orally once daily [see *Clinical Studies (14)*]. Among patients who received VIJOICE, 95% were exposed for 6 months or longer and 79% were exposed for greater than one year.

The median age of patients who received VIJOICE was 14 years (range, 2 to 50); 58% were female; 12% were White and race was not reported for 88%.

Serious adverse reactions occurred in 12% of patients who received VIJOICE. Serious adverse reactions occurring in two or more patients included dehydration (n = 2) and cellulitis (n = 2).

Dosage interruption of VIJOICE due to an adverse reaction occurred in 11% of patients. Adverse reactions which required dosage interruption in two or more patients included dizziness (n = 2) and vomiting (n = 2). Dose reductions of VIJOICE due to an adverse reaction occurred in 5% of patients. Adverse reactions which required dose reduction included alopecia, memory impairment, and soft tissue infection.

The most common adverse reactions ($\geq 10\%$) were diarrhea, stomatitis, and hyperglycemia. The most common Grade 3 or 4 laboratory abnormalities ($\geq 2\%$) were increased glucose, decreased hemoglobin, decreased phosphate, increased bilirubin, decreased sodium, and decreased platelets.

Adverse reactions and laboratory abnormalities are listed in Table 10 and Table 11, respectively.

Table 10: Adverse Reactions ($\geq 5\%$) in Patients with PROS Who Received VIJOICE in EPIK-P1

Adverse reactions	VIJOICE N = 57	
	All Grades (%)	Grade 3 or 4 (%)
Gastrointestinal disorders		
Diarrhea	16	0
Stomatitis ^a	16	0
Metabolism and nutrition disorders		
Hyperglycemia	12	0
Skin and subcutaneous tissue disorders		
Eczema	7	0
Dry skin	7	0
Alopecia	5	0
Nervous system disorders		
Headache	5	0

Infections and infestations

Cellulitis	5	3.5
------------	---	-----

Grading according to CTCAE Version 4.03.

^aStomatitis: including stomatitis and aphthous ulcer.

Clinically relevant adverse reactions in < 5% of patients who received VIJOICE included nausea, vomiting, dehydration, and mucosal dryness.

Table 11: Laboratory Abnormalities Worsening from Baseline in $\geq 10\%$ of Patients with PROS Who Received VIJOICE in EPIK-P1

Laboratory abnormality	VIJOICE ^a N = 57	
	All Grades %	Grade 3 or 4 %
Chemistry		
Decreased calcium (corrected)	60	0
Decreased phosphate	59	5 ^b
Increased glucose ^c	56	11 ^b
Increased glycosylated hemoglobin (HbA1c) ^d	38 ^d	N/A ^d
Increased creatinine	31	0
Increased bilirubin	29	2 ^b
Increased potassium	24	0
Increased triglycerides	19	0
Decreased magnesium	18	0
Increased aspartate aminotransferase (AST)	17	0
Increased cholesterol	13	0
Decreased albumin	13	0
Decreased sodium	12	2 ^b
Decreased potassium	12	0
Increased gamma glutamyl transferase (GGT)	11	0
Increased alanine aminotransferase (ALT)	10	0

Hematology

Decreased leukocyte	22	0
Decreased hemoglobin	20	6 ^b
Decreased lymphocyte	20	0
Decreased neutrophil	19	0
Increased lymphocyte	17	0
Decreased platelets	14	2 ^b

Grading according to CTCAE Version 4.03.

Abbreviation: N/A, not available.

^aThe denominator used to calculate the rate varied from 9 to 50 based on the number of patients with a baseline value and at least one post-treatment value.

^bNo Grade 4 laboratory abnormalities were reported.

^cGlucose increase is an expected laboratory abnormality of PI3K inhibition.

^dNo CTCAE grade available. For HbA1c, baseline values increasing post-treatment to a value above the upper limit of the normal range ($\geq 5.7\%$) are considered increased.

Other Clinical Trials Experience

The following additional adverse reactions and laboratory abnormality have been identified following administration of VIJOICE: hypersensitivity, lipase increased, dermatitis, and abdominal pain.

6.2 Postmarketing Experience and Other Spontaneous Adverse Reaction Reports

The following adverse reactions have been identified with VIJOICE use in patients with PROS in an expanded access program for compassionate use. Because these reactions are reported from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Metabolism and nutrition disorders: Decreased appetite.

Skin and subcutaneous tissue disorders: Pruritus, rash (including rash maculo-papular, rash erythematous, rash papular, and rash pruritic), acne (including dermatitis acneiform).

7 DRUG INTERACTIONS

7.1 Effect of Other Drugs on VIJOICE

CYP3A4 Inducers

Avoid coadministration of VIJOICE with strong CYP3A4 inducers and consider an alternative concomitant drug with no or minimal potential to induce CYP3A4.

Alpelisib is metabolized by CYP3A4. Concomitant use of VIJOICE with a strong CYP3A4 inducer may decrease alpelisib concentration [see *Clinical Pharmacology (12.3)*], which may decrease alpelisib activity.

Breast Cancer Resistance Protein Inhibitors (BCRP)

Avoid the use of BCRP inhibitors in patients treated with VIJOICE. If unable to use alternative drugs, when VIJOICE is used in combination with BCRP inhibitors, closely monitor for increased adverse reactions.

Alpelisib is transported by BCRP. Concomitant use of VIJOICE with a BCRP inhibitor may increase alpelisib exposure [see *Clinical Pharmacology (12.3)*], which may increase the risk of adverse reactions.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal data and mechanism of action, VIJOICE can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology (12.1)*]. There are no available data in pregnant women to inform the drug-associated risk. In animal reproduction studies, oral administration of alpelisib to pregnant rats and rabbits during organogenesis caused adverse developmental outcomes, including embryo-fetal mortality (post-implantation loss), reduced fetal weights, and increased incidences of fetal malformations at doses described below (see *Data*). Advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. However, the estimated background risk of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies in the U.S. general population.

Data

Animal Data

In embryo-fetal development studies in rats and rabbits, pregnant animals received oral doses of alpelisib during the period of organogenesis. In the rat study, animals were dosed at 3, 10, or 30 mg/kg/day from gestation day 6 to 17; and in the rabbit study, animals were dosed at 3, 15, 25, and 30 mg/kg/day from gestation day 7 to 20.

In rats, oral administration of alpelisib resulted in maternal toxicity (body weight loss, low food consumption) and no viable fetuses (post-implantation loss) at 30 mg/kg/day (approximately 3.6 to 1.2 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA). At a dose of 10 mg/kg/day (approximately 1.2 to 0.4 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA), toxicities included reduced fetal weight and increased incidences of skeletal malformations (bent scapula and thickened

or bent long bones) and fetal variations (enlarged brain ventricle, decreased bone ossification).

In a pilot embryo-fetal development study in rabbits, a dose of 30 mg/kg/day (approximately 7 to 2.2 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients based on BSA) resulted in no viable fetuses (post-implantation loss). Doses \geq 15 mg/kg/day (approximately 3.5 to 1.1 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA) resulted in increased embryo-fetal deaths, reduced fetal weights, and malformations, mostly related to the tail and head.

8.2 Lactation

Risk Summary

There are no data on the presence of alpelisib in human milk, its effects on milk production, or the breastfed child. Because of the potential for serious adverse reactions in the breastfed child, advise lactating women to not breastfeed during treatment with VIJOICE and for 1 week after the last dose.

8.3 Females and Males of Reproductive Potential

Pregnancy Testing

Verify the pregnancy status in females of reproductive potential prior to initiating VIJOICE.

Contraception

Females

VIJOICE can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with VIJOICE and for 1 week after the last dose.

Males

Advise male patients with female partners of reproductive potential to use condoms and effective contraception during treatment with VIJOICE and for 1 week after the last dose.

Infertility

Based on findings from animal studies, VIJOICE may impair fertility in males and females of reproductive potential [see *Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of VIJOICE have been established in pediatric patients 2 to less than 18 years of age with PROS based on results from a single-arm clinical study of VIJOICE (EPIK-P1) that enrolled 39 pediatric patients: 11 patients aged 2 to 5 years, 12 patients aged 6 to 11 years, and 16 patients aged 12 to less than 18 years of age [see *Adverse Reactions (6.1)* and *Clinical Studies (14)*].

The safety and effectiveness of VIJOICE in pediatric patients below the age of 2 years have not been established.

Although there were no new safety signals observed in pediatric patients, there is insufficient data to determine whether VIJOICE has an adverse impact on growth and development in pediatric patients with PROS. Based on the animal toxicity data (described below), regular monitoring of growth and development in pediatric patients treated with VIJOICE is recommended.

Animal Toxicity Data

In a 4-week general toxicology study, rats administered alpelisib had growth plate thickening and decreased trabeculae of the knee joint, dentin thinning, and degenerative odontoblasts at the dose of 30 mg/kg/day (approximately 2.8 to 1.2 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA). Dentin thinning/irregular dentin was also observed in the 13-week toxicology study in rats at the high dose of 20 mg/kg/day (approximately 2 to 0.8 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA).

8.5 Geriatric Use

There were no adult patients aged 65 years of age or older who received VIJOICE in EPIK-P1.

10 OVERDOSAGE


There is limited experience of overdose with alpelisib in clinical trials.

In cases where accidental overdose of alpelisib was reported in the clinical studies, the adverse reactions associated with the overdose were consistent with the known safety profile of alpelisib and included hyperglycemia, nausea, asthenia, and rash.

Initiate general symptomatic and supportive measures in all cases of overdose where necessary. There is no known antidote for VIJOICE.

11 DESCRIPTION

VIJOICE (alpelisib) is a kinase inhibitor. The chemical name of alpelisib is (2S)-N¹-[4-Methyl-5-[2-(2,2,2-trifluoro-1,1-dimethylethyl)-4-pyridinyl]-2-thiazolyl]-1,2-pyrrolidinedicarboxamide. Alpelisib is a white to almost white powder. The molecular formula for alpelisib is C₁₉H₂₂F₃N₅O₂S and the relative molecular mass is 441.47 g/mol. The pH of a 1.0% (m/V) solution of alpelisib in water/ethanol (50:50 V/V) is approximately 6.2. The chemical structure of alpelisib is shown below:



VIJOICE film-coated tablets are supplied for oral administration with three strengths that contain 50 mg, 125 mg, and 200 mg of apelisib. The tablets also contain hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, and sodium starch glycolate. The film-coating contains hypromellose, iron oxide red (applicable only to 50 mg and 200 mg strengths), iron oxide yellow, macrogol/polyethylene glycol (PEG) 4000, talc, and titanium dioxide.

VIJOICE oral granules are supplied for oral administration with one strength that contains 50 mg of apelisib. The granules also contain hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, and sodium starch glycolate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Apelisib is an inhibitor of phosphatidylinositol-3-kinase (PI3K) with inhibitory activity predominantly against PI3K α . Gain-of-function mutations in the gene encoding the catalytic α -subunit of PI3K (PIK3CA) lead to activation of PI3K α and Akt-signaling, cellular transformation and the generation of tumors in in vitro and in vivo models.

Activating mutations in PIK3CA have been found to induce a spectrum of overgrowths and malformations comprising a wide group of clinically recognizable disorders commonly known as PROS.

In an inducible mouse model of Congenital Lipomatous Overgrowth, Vascular Malformations, Epidermal Nevi, Scoliosis/Skeletal and Spinal syndrome (CLOVES), a phenotype of PROS, apelisib inhibition of the PI3K pathway resulted in the prevention or improvement of organ abnormalities associated with the disease, depending on when apelisib treatment was started. These findings were reversed after withdrawal of apelisib.

12.2 Pharmacodynamics

The exposure-response relationship and time course of pharmacodynamic response for the safety and effectiveness of VIJOICE have not been characterized.

Cardiac Electrophysiology

At a dose of 300 mg, apelisib does not prolong the QT interval to any clinically relevant extent in the oncology setting. VIJOICE is not approved for use in the oncology setting.

12.3 Pharmacokinetics

The pharmacokinetics of apelisib has been studied in healthy subjects and adult patients with solid tumors and are presented as mean (% CV) under fed conditions unless otherwise specified. Apelisib maximum plasma concentration (C_{max}) was 277 ng/mL (24%) and area under the curve (AUC) was 2,090 hr*ng/mL (24%) following administration of a single 50 mg dose of VIJOICE. Steady state apelisib C_{max} and AUC increased proportionally over the dose range of 30 mg (0.6 times the lowest approved recommended dosage) to 450 mg (1.8 times the highest approved recommended dosage) under fed conditions. The mean accumulation of apelisib was 1.3 to 1.5 and steady state plasma concentrations were reached within 3 days following daily dosage.

Absorption

The median time to reach peak plasma concentration (T_{max}) ranged between 2.0 to 4.0 hours.

Effect of Food

A high-fat high-calorie (HFHC) meal (985 calories with 58.1 g of fat) increased apelisib AUC by 73% and C_{max} by 84%, and a low-fat low-calorie (LFLC) meal (334 calories with 8.7 g of fat) increased apelisib AUC by 77% and C_{max} by 145% following administration of a single 300 mg dose of apelisib as tablets. No clinically relevant differences in apelisib AUC were observed between LFLC and HFHC meals.

No clinically relevant food effect on apelisib pharmacokinetics was observed after a single 50 mg oral dose of VIJOICE as granules taken with a LFLC meal as compared to that taken under fasted conditions.

No clinically relevant differences in apelisib C_{max} and AUC were observed following administration of a single 50 mg dose of VIJOICE as granules or tablets taken with a LFLC meal.

Distribution

The apparent volume of distribution of apelisib at steady state is 114 L (46%). Protein binding of apelisib is 89% and is independent of concentration.

Elimination

The half-life of apelisib is predicted to be 8 to 9 hours. The clearance of apelisib is 9.2 L/hr (21%) under fed conditions.

Metabolism

Apelisib is primarily metabolized by chemical and enzymatic hydrolysis to form its metabolite BZG791 and followed by CYP3A4 mediated hydroxylation.

Excretion

Following a single oral dose of 400 mg (1.6 times the highest approved recommended dosage) radiolabeled apelsisib under fasted condition, 81% of the administered dose was recovered in feces (36% unchanged) and 14% (2% unchanged) in urine. CYP3A4-mediated metabolites (12%) and glucuronides amounted to approximately 15% of the dose.

Specific Populations

No clinically significant differences in the pharmacokinetics of apelsisib were predicted based on age (21 to 87 years), sex, race/ethnicity (Japanese or Caucasian), body weight (37 to 181 kg), mild to moderate renal impairment (CL_{cr} 30 to < 90 mL/min based on the Cockcroft-Gault formula), or mild to severe hepatic impairment (Child-Pugh Class A, B, and C). The effect of severe renal impairment (CL_{cr} < 30 mL/min) on the pharmacokinetics of apelsisib is unknown.

Drug Interaction Studies

Clinical Studies and Model-Informed Approaches

Acid Reducing Agents: No clinically significant differences in the pharmacokinetics of apelsisib were observed when used concomitantly with ranitidine (H₂ receptor antagonist) and administered with food as directed.

Concomitant use of ranitidine decreased apelsisib AUC approximately 30% and C_{max} by 51% with a single 300 mg oral dose (1.2 times the highest approved recommended dosage) of apelsisib under the fasted state. In the presence of a low-fat low-calorie meal, AUC was decreased by 21% and C_{max} by 36% with ranitidine.

CYP3A4, CYP2C8, CYP2C9, CYP2C19 and CYP2B6 Substrates: Coadministration of repeated doses of apelsisib 300 mg with a single-dose of sensitive substrates of CYP3A4 (midazolam), CYP2C8 (repaglinide), CYP2C9 (warfarin), CYP2C19 (omeprazole) and CYP2B6 (bupropion), administered as a cocktail did not show clinically significant pharmacokinetic interactions.

No clinically significant differences in pharmacokinetics of everolimus (a substrate of CYP3A4 and P-gp) were observed when used concurrently with apelsisib.

Effect of CYP3A4 Inducers on Apelsisib: Coadministration of repeat doses of rifampin (a strong CYP3A4 inducer) with a single 300 mg dose of apelsisib decreased apelsisib C_{max} by 38% and AUC by 57%, respectively. Coadministration of rifampin with repeat doses of 300 mg apelsisib decreased apelsisib C_{max} by 59% and AUC by 74%, respectively.

Model-Informed Approaches

Coadministration of repeat doses of ketoconazole (a strong CYP3A4 inhibitor) with a single 300 mg dose of apelsisib is expected to increase apelsisib AUC by 37% or less.

Coadministration of repeat doses of efavirenz (a moderate CYP3A4 inducer) with a single 300 mg dose of apelsisib is expected to decrease apelsisib AUC by 30% or less.

In Vitro Studies

Effect of Transporter on Apelsisib: Apelsisib is a substrate of BCRP.

Effect of Apelsisib on Transporters: Apelsisib is an inhibitor of P-gp. Apelsisib has a low potential to inhibit BCRP, MRP2, BSEP, OATP1B1, OATP1B3, OCT1, OAT1, OAT3, OCT2, MATE1, and MATE2K at clinically relevant concentrations.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Alpelisib was not carcinogenic in a 2-year carcinogenicity study conducted in rats when administered by daily oral gavage at doses up to 4 mg/kg (approximately 0.5 to 0.2 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA).

Alpelisib was not mutagenic in an in vitro bacterial reverse mutation (Ames) assay, or aneugenic or clastogenic in human cell micronucleus and chromosome aberration tests. Alpelisib was not genotoxic in an in vivo rat micronucleus test.

In a fertility and early embryonic development study in rats, female animals were administered alpelisib doses of 3, 10, and 20 mg/kg/day orally. Animals were dosed for 4-weeks prior to pairing, during the mating period, and up to Gestation Day 6. At the dose of 20 mg/kg/day, alpelisib increased pre- and post-implantation losses, leading to reduced numbers of implantation sites and live embryos. These findings were observed at doses approximately 2.4 to 0.8 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on body surface areas (BSA). In a repeated-dose toxicity study in rats, adverse effects in female reproductive organs included vaginal atrophy and estrous cycle variations in rats at doses \geq 6 mg/kg/day (approximately 0.7 to 0.2 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA).

In a male fertility study, alpelisib administered orally at doses of 3, 10 and 20 mg/kg/day for up to 99 days (10-weeks prior to pairing, during mating period and continuing during post-pairing) to male rats, resulted in reduced weights of seminal vesicles and prostate, which correlated with atrophy and/or reduced secretion in prostate and seminal vesicles at \geq 10 mg/kg/day (approximately 1.2 to 0.4 times the initial recommended doses of 50 mg and 250 mg in pediatric and adult patients, respectively, based on BSA). No adverse effects on male fertility parameters were observed at doses up to 20 mg/kg/day.

14 CLINICAL STUDIES

The efficacy of VIJOICE was assessed in EPIK-P1 (NCT04285723), a single-arm clinical study in patients who were treated as part of an expanded access program for compassionate use which enrolled patients across seven sites in five countries (France, Spain, US, Ireland, and Australia). Eligible patients 2 years of age and older with PIK3CA-Related Overgrowth Spectrum (PROS) who received VIJOICE had clinical manifestations of PROS that were assessed by the treating physician as severe or life-threatening and necessitating systemic treatment and had documented evidence of mutation in the PIK3CA gene. Patients received VIJOICE at dosages based on age ranging from 50 mg to 250 mg orally once daily.

The efficacy of VIJOICE was evaluated in a total of 37 patients with at least one target lesion identified on imaging performed within 24 weeks prior to receipt of the first dose of VIJOICE. The median age of patients was 14 years (range: 2 to 38); 22% of patients were 2 to 5 years, 22% were 6 to 11 years, 27% were 12 to less than 18 years of age, and 30% were \geq 18 years; 57% were female, 11% were White and race was not

reported for 89%. Ninety-two percent of patients had congenital overgrowth and 8% had early childhood-onset. Patients had heterogeneous manifestations of PROS, including CLOVES (81%), Megalencephaly-Capillary Malformation Polymicrogyria (MCAP; 8%), Klippel-Trenaunay Syndrome (KTS; 2.7%), Facial Infiltrating Lipomatosis (FIL; 8%), and Other (5%). Five percent (5%) of patients had concurrent manifestations of CLOVES and MCAP.

The major efficacy outcome measure for the study was the proportion of patients with radiological response at Week 24 as determined by blinded independent central review (BICR), defined as a $\geq 20\%$ reduction from baseline in the sum of measurable target lesion volume (1 to 3 lesions) confirmed by at least one subsequent imaging assessment, in the absence of a $\geq 20\%$ increase from baseline in any target lesion, progression of non-target lesions, or appearance of a new lesion. An additional efficacy outcome measure was duration of response, defined as the time from the first documented response to the date of the first documented disease progression or death due to any cause.

Efficacy results are presented in Table 12.

Table 12: Efficacy Results at Week 24 in EPIK-P1

Efficacy parameters	All patients N = 37
Response rate^{a,b}	
Responders, n (%)	10 (27)
95% CI	(14, 44)
Duration of response (DOR)	
Median in months (range)	NR (0.9+, 42.9+)
% ≥ 6 months	70
% ≥ 12 months	60

Abbreviation: +, censored observation.

^aConfirmed response as determined by blinded independent central review (BICR).

^bPatients without any response assessment at Week 24 were considered non-responders.

16 HOW SUPPLIED/STORAGE AND HANDLING

VIJOICE (alpelisib) 50 mg, 125 mg, and 200 mg film-coated tablets are available in blister packs based on daily dose as described in Table 13.

Table 13: VIJOICE Tablets Daily Dose Blister Packs

Daily dose	Each carton contains	Each child-resistant blister pack contains	Appearance of tablet	NDC

50 mg daily dose	One 28-day supply blister pack	28 tablets: 50 mg apelisib per tablet	Light yellow, unscored, round and curved with beveled edges, debossed with "C7" on one side and "NVR" on the other side	NDC 0078-1021-84
125 mg daily dose	One 28-day supply blister pack	28 tablets: 125 mg apelisib per tablet	Dark yellow, unscored, ovaloid and curved with beveled edges, debossed with "Y7" on one side and "NVR" on the other side	NDC 0078-1028-84
250 mg daily dose	Two 14-day supply blister packs (56 tablets total)	14 tablets: 200 mg apelisib per tablet, and 14 tablets: 50 mg apelisib per tablet	Pale yellow, unscored, ovaloid and curved with beveled edges, debossed with "CL7" on one side and "NVR" on the other side Light yellow, unscored, round and curved with beveled edges, debossed with "C7" on one side and "NVR" on the other side	NDC 0078-1035-02

Table 14: VIJOICE Oral Granules

Daily dose	Each carton contains	Each child-resistant packet contains	Appearance	NDC
50 mg daily dose	28 packets (28-day supply of single-use packets)	50 mg apelisib	White to almost white free-flowing mixture of powder and granules	NDC 0078-1175-51

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient and their caregivers to read the FDA-approved patient labeling (Patient Information).

Severe Hypersensitivity

Inform patients and their caregivers of the signs and symptoms of hypersensitivity. Advise patients and their caregivers to contact their healthcare provider immediately for signs and symptoms of hypersensitivity [see *Warnings and Precautions (5.1)*].

Severe Cutaneous Adverse Reactions

Inform patients and their caregivers of the signs and symptoms of severe cutaneous adverse reactions (SCARs). Advise patients and their caregivers to contact their

healthcare provider immediately for signs and symptoms of SCARs (e.g., a prodrome of fever, flu-like symptoms, mucosal lesions, progressive skin rash, or lymphadenopathy) [*see Warnings and Precautions (5.2)*].

Hyperglycemia

Advise patients and their caregivers that VIJOICE may cause hyperglycemia and the need to monitor fasting glucose periodically during therapy. Advise patients and their caregivers of the signs and symptoms of hyperglycemia (e.g., excessive thirst, urinating more often than usual or higher amount of urine than usual, or increased appetite with weight loss) [*see Warnings and Precautions (5.3)*].

Pneumonitis

Inform patients and their caregivers that VIJOICE may cause pneumonitis and to immediately report new or worsening respiratory symptoms [*see Warnings and Precautions (5.4)*].

Diarrhea or Colitis

Advise patients and their caregivers that VIJOICE may cause diarrhea, which may be severe, and to start anti-diarrheal treatment, increase oral fluids, and notify their healthcare provider if diarrhea occurs while taking VIJOICE [*see Warnings and Precautions (5.5)*].

Advise patients and their caregivers that VIJOICE may cause colitis and to notify their healthcare provider immediately of any symptoms of colitis, such as abdominal pain and mucus or blood in stool, while taking VIJOICE [*see Warnings and Precautions (5.5)*].

Alopecia

Advise patients and caregivers that VIJOICE may cause alopecia [*see Adverse Reactions (6.1)*]. Advise patients and caregivers to contact a dermatologist if alopecia becomes a concern [*see Dosage and Administration (2.5)*].

Embryo-Fetal Toxicity

- Inform pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females to inform their healthcare provider of a known or suspected pregnancy [*see Warnings and Precautions (5.6)* and *Use in Specific Populations (8.1)*].
- Advise females of reproductive potential to use effective contraception during treatment with VIJOICE and for 1 week after the last dose [*see Use in Specific Populations (8.3)*].
- Advise male patients with female partners of reproductive potential to use condoms and effective contraception during treatment with VIJOICE and for 1 week after the last dose [*see Use in Specific Populations (8.3)*].

Lactation

Advise women not to breastfeed during treatment with VIJOICE and for 1 week after the last dose [*see Use in Specific Populations (8.2)*].

Infertility

Advise males and females of reproductive potential that VIJOICE may impair fertility [*see Use in Specific Populations (8.3)*].

Drug Interactions

Advise patients and their caregivers to inform their healthcare providers of all concomitant medications, herbal and dietary supplements [see *Drug Interactions (7.1)*].

Dosing

Instruct patients and their caregivers of the following:

- Take VIJOICE with food at approximately the same time each day [see *Dosage and Administration (2.3)*].
- Swallow the tablets whole (tablets should not be crushed, chewed, or split prior to swallowing) [see *Dosage and Administration (2.4)*].
- For patients unable to swallow tablets, advise how to prepare an oral suspension using the tablets or how to take the oral granules (if applicable) [see *Dosage and Administration (2.4)*].
- If a dose of VIJOICE is missed, it can be taken with food within 9 hours after the time it is usually taken. After more than 9 hours, skip the dose for that day. The next day, take VIJOICE at the usual time. Instruct patients not to take 2 doses to make up for a missed dose [see *Dosage and Administration (2.3)*].
- If vomiting occurs after taking the dose of VIJOICE, they should not take an additional dose on that day and should resume the usual dosing schedule the next day at the usual time [see *Dosage and Administration (2.3)*].

Distributed by
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

© Novartis

T2025-38

PATIENT INFORMATION	
VIJOICE® (vi' joiz) (alpelisib) tablets	VIJOICE® (vi' joiz) (alpelisib) oral granules
What is VIJOICE? VIJOICE is a prescription medicine used to treat adults and children 2 years of age and older with severe phosphatidylinositol-3-kinase catalytic subunit alpha (PIK3CA)-Related Overgrowth Spectrum (PROS). It is not known if VIJOICE is safe and effective in children below 2 years of age.	
Do not take VIJOICE if you have had a severe allergic reaction to alpelisib or are allergic to any of the ingredients in VIJOICE. <ul style="list-style-type: none">• See the end of this Patient Information leaflet for a complete list of the ingredients in VIJOICE.• See “What are the possible side effects of VIJOICE?” for signs and symptoms of severe allergic reactions.	
Before you take VIJOICE, tell your healthcare provider about all of your medical conditions, including if you: <ul style="list-style-type: none">• have a history of diabetes.• have a history of skin rash, redness of skin, blistering of the lips, eyes or mouth, or skin peeling.	

- are pregnant or plan to become pregnant. VIJOICE can harm your unborn baby.
Females who are able to become pregnant:
 - Your healthcare provider will check to see if you are pregnant before you start treatment with VIJOICE.
 - You should use effective birth control (contraception) during treatment with VIJOICE and for 1 week after the last dose. Talk to your healthcare provider about birth control methods that may be right for you during this time.
 - If you become pregnant or think you are pregnant, tell your healthcare provider right away.
- **Males** with female partners who are able to become pregnant should use condoms and effective birth control during treatment with VIJOICE and for 1 week after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away.
- are breastfeeding or plan to breastfeed. It is not known if VIJOICE passes into your breast milk. Do not breastfeed during treatment with VIJOICE and for 1 week after the last dose.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. VIJOICE and other medicines may affect each other causing side effects. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take VIJOICE?

- Read the detailed **“Instructions for Use”** that comes with VIJOICE for information about the right way to prepare and take or give a dose of VIJOICE tablets or oral granules.
- Take VIJOICE exactly as your healthcare provider tells you.
- Do not change your dose or stop taking VIJOICE unless your healthcare provider tells you.
- Take VIJOICE 1 time each day, at about the same time each day.
- Take VIJOICE with food.
- VIJOICE comes as tablets and oral granules. Your healthcare provider will prescribe the dosage form that is right for you:
 - If your prescribed daily dose is 50 mg, your healthcare provider may prescribe VIJOICE tablets or VIJOICE oral granules.
 - If your prescribed daily dose is 125 mg or 250 mg, you can only take VIJOICE tablets.
- **Do not** use multiple packets of 50 mg oral granules or part of a packet to prepare a dose. **Do not** combine VIJOICE tablets with VIJOICE oral granules to prepare your prescribed dose.
- If you miss a dose of VIJOICE, you may take it with food up to 9 hours after the time you usually take it. If it has been more than 9 hours after you usually take your dose, skip the dose for that day. The next day, take the dose at your usual time. Do not take 2 doses to make up for a missed dose.
- If you vomit after taking a dose of VIJOICE, do not take another dose on that day. Take your next dose at your usual time.
- If you take too much VIJOICE, call your healthcare provider or go to the nearest hospital emergency room right away.

VIJOICE tablets:

- VIJOICE tablets can be swallowed whole, or prepared as a suspension to take or give by mouth or through a feeding tube.
- If you are taking or giving VIJOICE tablets whole:
 - Swallow VIJOICE tablets whole and take with food. Do not chew, divide, or crush the tablets.
 - Do not take VIJOICE tablets that are broken, cracked, or that look damaged.
- If you cannot swallow tablets whole or need to use a feeding tube, you will need to mix VIJOICE tablets with water before taking or giving the dose.

VIJOICE oral granules:

- VIJOICE oral granules can be placed directly onto the tongue and swallowed with water, or mixed with a beverage or soft food to take or give by mouth, or mixed with water to give through a feeding tube.

What are the possible side effects of VIJOICE?

VIJOICE may cause serious side effects, including:

- **Severe allergic reactions.** Tell your healthcare provider or get medical help right away if you have trouble breathing, swelling of the face or throat, flushing, rash, fever, or fast heart rate during treatment with VIJOICE.
- **Severe skin reactions.** Tell your healthcare provider or get medical help right away if you get severe rash or rash that keeps getting worse, reddened skin, flu-like symptoms, blistering of the lips, eyes or mouth, blisters on the skin or skin peeling, with or without fever.
- **High blood sugar levels (hyperglycemia).** Hyperglycemia is common with VIJOICE and may be severe. Your healthcare provider will monitor your sugar levels before you start and during treatment with VIJOICE. Your healthcare provider may monitor your sugar levels more often if you have a history of diabetes. Tell your healthcare provider right away if you develop symptoms of hyperglycemia, including:
 - excessive thirst
 - dry mouth
 - more frequent urination than usual or a higher amount of urine than normal
 - increased appetite with weight loss
 - confusion
 - nausea
 - vomiting
 - fruity odor on breath
 - difficulty breathing
 - dry or flushed skin
- **Lung problems (pneumonitis).** Tell your healthcare provider right away if you develop new or worsening symptoms of lung problems, including:
 - shortness of breath or trouble breathing
 - cough
 - chest pain
- **Diarrhea or colitis (inflammation of your intestines).** Diarrhea is common with VIJOICE and may be severe. Severe diarrhea can lead to the loss of too much body water (dehydration) and kidney injury. Tell your healthcare provider right away if you develop diarrhea, stomach-area (abdominal) pain, or see mucus or blood in your stool during treatment with VIJOICE. Your healthcare provider may tell you to drink more fluids or take medicines to treat diarrhea or colitis.

Your healthcare provider may tell you to decrease your dose, temporarily stop your

treatment, or completely stop your treatment with VIJOICE if you get certain serious side effects.

The most common side effects of VIJOICE include:

- diarrhea
- mouth sores (stomatitis)
- high blood sugar

VIJOICE may cause hair loss (alopecia). Talk to a skin specialist (dermatologist) if this is a concern for you.

VIJOICE may affect fertility in males and in females who are able to become pregnant.

Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of VIJOICE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store VIJOICE?

- Store VIJOICE at room temperature between 68°F to 77°F (20°C to 25°C).

Keep VIJOICE and all medicines out of the reach of children.

General information about the safe and effective use of VIJOICE.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use VIJOICE for a condition for which it was not prescribed. Do not give VIJOICE to other people, even if they have the same symptoms you have. It may harm them. You can ask your healthcare provider or pharmacist for more information about VIJOICE that is written for health professionals.

What are the ingredients in VIJOICE?

Active ingredient: apelisib

Inactive ingredients for VIJOICE tablets: hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, and sodium starch glycolate. The film-coating contains hypromellose, iron oxide red (applicable only to 50 mg and 200 mg strengths), iron oxide yellow, macrogol/polyethylene glycol (PEG) 4000, talc, and titanium dioxide.

Inactive ingredients for VIJOICE oral granules: hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, and sodium starch glycolate.

Distributed by: Novartis Pharmaceuticals Corporation, East Hanover, New Jersey 07936
© Novartis

For more information, go to www.VIJOICE.com or call 1-888-VIJOICE (1-888-845-6423).

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 7/2025

T2025-39

INSTRUCTIONS FOR USE

VIJOICE® [vi' joiz]
(apelisib)
tablets

This Instructions for Use contains information on how to prepare and take or give a dose of VIJOICE tablets.

Read this Instructions for Use carefully before taking or giving VIJOICE tablets.

Talk to your healthcare provider or pharmacist if you have any questions. You can also call Novartis Pharmaceuticals Corporation at 1-888-845-6423 for more information.

Important Information You Need to Know Before Taking or Giving VIJOICE

- Take or give VIJOICE exactly as your healthcare provider tells you.
- **Do not** change your dose or stop taking VIJOICE unless your healthcare provider tells you.
- Take VIJOICE 1 time each day, at about the same time each day.
- Take VIJOICE with food.
- **Do not** combine VIJOICE tablets with VIJOICE oral granules to prepare your prescribed dose.
- If you miss a dose of VIJOICE, you may take it with food up to 9 hours after the time you usually take it. If it has been more than 9 hours after you usually take your dose, skip the dose for that day. The next day, take the dose at your usual time. **Do not** take 2 doses to make up for a missed dose.
- If you vomit after taking a dose of VIJOICE, **do not** take another dose on that day. Take your next dose at your usual time.
- If you take too much VIJOICE, call your healthcare provider or go to the nearest hospital emergency room right away.

Storing VIJOICE tablets

- Store VIJOICE tablets at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep VIJOICE and all medicines out of the reach of children.

Preparing and taking or giving VIJOICE tablets

You can take or give VIJOICE tablets whole by mouth, or mixed with water (suspension) by mouth or through a feeding tube as described in the sections below:

Taking or giving VIJOICE tablets whole by mouth

Swallow VIJOICE tablets whole and take with food.

- **Do not** chew, divide, or crush the tablets.
- **Do not** take VIJOICE tablets that are broken, cracked, or that look damaged.

Taking or giving VIJOICE tablets mixed with water (suspension) by mouth

If you cannot swallow tablets whole, VIJOICE tablets can be prepared as a suspension and taken or given by mouth.

Gather the following supplies (see Figure A):

- the number of VIJOICE tablet(s) needed for your prescribed dose
- water
- a clean empty cup
- a spoon

Step 1. Wash and dry your hands.

Step 2. Place VIJOICE tablets into a cup that contains 2 to 4 ounces of water. Use water only.

Step 3. Let the tablets stand in water **for about 5 minutes.**

Step 4. Crush the tablets with a spoon and

Figure A



Figure B

stir to disperse the tablets in water (suspension). The suspension will be cloudy and you may see tablet particles (see **Figure B**).

Step 5. Drink all of the suspension right away. If you cannot take or give it right away, take or give the suspension within 60 minutes of preparing the dose. Stir the suspension again with the same spoon before drinking it.

Step 6. Add about 1 to 2 ounces of water to the same cup and stir with the same spoon. Then drink all of the suspension to make sure the entire dose is taken.

Step 7. Repeat Step 6 until there are no tablet particles remaining in the cup.

Step 8. Wash your hands and all supplies used to take or give VIJOICE tablets. Throw away any VIJOICE suspension that is not taken within 60 minutes after it is prepared.

Giving VIJOICE tablets mixed with water (suspension) through a feeding tube

VIJOICE tablets may be given through a feeding tube, according to the manufacturer's instructions and as directed by your healthcare provider. **Only use silicone or polyurethane nasogastric tubes with a French size 5 to 12, or silicone gastric tubes with a French size 12 to 24.**

Gather the following supplies (see Figure C):

- the number of VIJOICE tablet(s) needed for your prescribed dose
- water
- a clean empty cup
- a spoon
- an enteral syringe

Step 1. Wash and dry your hands.

Step 2. Place VIJOICE tablets into a cup that contains 1 to 2 ounces (about 30 to 60 mL) of water. Use water only.

Step 3. Let the tablets stand in water for about 5 minutes.

Step 4. Crush the tablets with a spoon and stir to disperse the tablets in water (suspension). The suspension will be cloudy



Figure C



and you may see tablet particles (see **Figure B**).

Step 5. Draw up all of the suspension from the cup into an enteral syringe (see **Figure D**) and give it through the feeding tube right away. If you cannot give it right away, give the suspension within 60 minutes of preparing the dose. Stir the suspension again with the same spoon before drawing it up.


Step 6. Add about 1 to 2 ounces (about 30 to 60 mL) of water to the same cup and stir with the same spoon.

Step 7. Draw up all of the suspension into the same enteral syringe and give it through the feeding tube.

Step 8. Repeat Steps 6 and 7 if there are tablet particles remaining in the enteral syringe or in the cup.

Step 9. Wash your hands and all supplies used to give VIJOICE tablets. Follow the manufacturer's instructions to clean your enteral syringe.

Throw away any VIJOICE suspension that is not taken within 60 minutes after it is prepared.



Distributed by:

Novartis Pharmaceuticals Corporation East Hanover, New Jersey 07936

© Novartis

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Issued: 7/2025

T2025-40

INSTRUCTIONS FOR USE

VIJOICE® [vi' joiz]
(alpelisib)
oral granules

This Instructions for Use contains information on how to prepare and take or give a dose of VIJOICE oral granules.

Read this Instructions for Use carefully before taking or giving VIJOICE oral granules.

Talk to your healthcare provider or pharmacist if you have any questions. You can also call Novartis Pharmaceuticals Corporation at 1-888-845-6423 for more information.

Important Information You Need to Know Before Taking or Giving VIJOICE

- Take or give VIJOICE exactly as your healthcare provider tells you.
- **Do not** change your dose or stop taking VIJOICE unless your healthcare provider tells you.

- Take VIJOICE 1 time each day, at about the same time each day.
- Take VIJOICE with food.
- Each VIJOICE oral granules packet is for single use only.
- **Do not** use the packet if the packet seal is broken.
- **Do not** use multiple packets of 50 mg oral granules or part of a packet to prepare a dose.
- **Do not** combine VIJOICE tablets with VIJOICE oral granules to prepare your prescribed dose.
- If you miss a dose of VIJOICE, you may take it with food up to 9 hours after the time you usually take it. If it has been more than 9 hours after you usually take your dose, skip the dose for that day. The next day, take the dose at your usual time. **Do not** take 2 doses to make up for a missed dose.
- If you vomit after taking a dose of VIJOICE, **do not** take another dose on that day. Take your next dose at your usual time.
- If you take too much VIJOICE, call your healthcare provider or go to the nearest hospital emergency room right away.

Storing VIJOICE oral granules

- Store VIJOICE oral granules at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep VIJOICE and all medicines out of the reach of children.

Preparing and taking or giving VIJOICE oral granules

You can take or give VIJOICE oral granules directly onto the tongue swallowed with water, or mixed with a beverage or soft food by mouth, or mixed with water through a feeding tube as described in the sections below:

Taking or giving VIJOICE oral granules directly onto the tongue

Gather the following supplies (see Figure A):

- 1 VIJOICE oral granules packet
- scissors
- water

Step 1. Wash and dry your hands.

Step 2. Hold the packet of oral granules with the cut line on top.

Step 3. Shake the packet gently to ensure the VIJOICE oral granules are in the lower part of the packet.

Step 4. Using scissors, open the packet along the cut line.


Step 5. Pour all the oral granules from 1 VIJOICE oral granules packet directly onto the tongue by tapping the side and top of the packet, and swallow it with about 2 to 4 ounces of water (see **Figure B**).

Step 6. If there is any medicine remaining in the mouth, rinse the mouth with

Figure A



Figure B



additional water and swallow to make sure the entire dose is taken.

Step 7. Wash your hands and all supplies used to take or give VIJOICE oral granules.

Taking or giving VIJOICE oral granules mixed with a beverage or soft food by mouth

Gather the following supplies (see Figure C):

- 1 VIJOICE oral granules packet
- scissors
- the beverage **or** soft food of your choice among water, milk, apple juice, applesauce or yogurt. See Step 6.
- a clean empty cup
- a spoon

Step 1. Wash and dry your hands.

Step 2. Hold the packet of oral granules with the cut line on top.

Step 3. Shake the packet gently to ensure the VIJOICE oral granules are in the lower part of the packet.

Step 4. Using scissors, open the packet along the cut line.

Step 5. Pour the oral granules from 1 VIJOICE oral granules packet into a cup by tapping the side and top of the packet (see **Figure D**).

Step 6. Add 1 to 3 teaspoons of a beverage (water, milk, or apple juice) or soft food (applesauce or yogurt), stir with a spoon and take or give the mixture right away. If you do not take or give it right away, take or give the mixture within 60 minutes of preparing the dose. Stir the mixture with the same spoon before taking or giving it.

Step 7. Rinse the cup with up to 2 ounces of water, milk, or apple juice. Drink the rinse right away to make sure the entire dose is taken. Repeat this step until there are no oral granules remaining in the cup.

Step 8. Wash your hands and all supplies used to take or give VIJOICE oral granules.

Throw away any VIJOICE oral granules mixture that is not taken within 60 minutes after it is prepared.

Giving VIJOICE oral granules mixed with water (suspension) through a

Figure C



Figure D



feeding tube

VIJOICE oral granules may be given through a feeding tube, according to the manufacturer's instructions and as directed by your healthcare provider. **Only use silicone or polyurethane nasogastric tubes with a French size 8 to 12, or silicone gastric tubes with a French size 12 to 24.**

Gather the following supplies (see Figure E):

- 1 VIJOICE oral granules packet
- water
- a clean empty cup
- a spoon
- an enteral syringe

Step 1. Wash and dry your hands.

Step 2. Pour the oral granules from 1 VIJOICE oral granules packet into a cup by tapping the side and top of the packet (see Figure D).

Step 3. Add 4 teaspoons (about 20 mL) of water to the same cup and stir gently with a spoon to disperse the oral granules in the water (suspension). Use water only.

Step 4. Draw up all of the suspension from the cup into an enteral syringe (see **Figure F**) and give it through the feeding tube right away. If you cannot give it right away, give the suspension within 60 minutes of preparing the dose. Stir the suspension again with the same spoon before drawing it up.

Step 5. Add 4 teaspoons (about 20 mL) of water to the same cup and stir with the same spoon.

Step 6. Draw up all of the suspension from the cup into the same enteral syringe and give it through the feeding tube.

Step 7. Repeat Steps 5 and 6 until there are no oral granules remaining in the cup or in the syringe.

Step 8. Wash your hands and all supplies used give VIJOICE oral granules. Follow the manufacturer's instructions to clean your enteral syringe.

Throw away any VIJOICE suspension that is not taken within 60 minutes after it is prepared.

Figure E



Figure F



Distributed by: Novartis Pharmaceuticals Corporation East Hanover, New Jersey 07936
© Novartis

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Issued: 7/2025

T2025-41

PRINCIPAL DISPLAY PANEL

Vijoice®
(alpelisib) tablets

NDC 0078-1021-84

50 mg daily dose

Take one 50 mg tablet once daily

Rx only


Recommended Dosage: Take one 50 mg tablet once daily with food. Swallow tablets whole. DO NOT chew, crush, or split tablets.

Directions: See enclosed Instructions for Use for administration instructions, including options for patients who are unable to swallow whole tablets.

28-Day Supply

Contains: One blister pack containing 28 tablets

NOVARTIS



PRINCIPAL DISPLAY PANEL

Vioice®
(alpelisib) tablets

NDC 0078-1028-84

125 mg daily dose

Take one 125 mg tablet once daily

Rx only

Recommended Dosage: Take one 125 mg tablet once daily with food.
Swallow tablets whole. DO NOT chew, crush, or split tablets.

Directions: See enclosed Instructions for Use for administration instructions, including options for patients who are unable to swallow whole tablets.

28-Day Supply

Contains: One blister pack containing 28 tablets



PRINCIPAL DISPLAY PANEL

Vioice[®]
 (alpelisib) tablets

NDC 0078-1035-02

250 mg daily dose

Take one 200 mg tablet and one 50 mg tablet once daily

Rx only

Recommended Dosage: Take one 200 mg tablet and one 50 mg tablet once daily with food. Swallow tablets whole. DO NOT chew, crush, or split tablets.

Directions: See enclosed Instructions for Use for administration instructions, including options for patients who are unable to swallow whole tablets.

28-Day Supply (56 Tablets)

Contains: Two 14-day blister packs each containing 28 tablets:

200 mg
14 tablets per pack

50 mg
14 tablets per pack

NOVARTIS



PRINCIPAL DISPLAY PANEL

Rx only

NDC 0078-1175-51

Vioice[®]

(apelsib) oral granules

50 mg daily dose

Take one 50 mg packet once daily

Directions: See enclosed Instructions For Use for administration instructions.

28 Packets

NOVARTIS



VIOICE

alpelisib tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0078-1021
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPELISIB (UNII: 08W5N2C97Q) (ALPELISIB - UNII:08W5N2C97Q)	ALPELISIB	50 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	YELLOW (light)	Score	no score
Shape	ROUND (round and curved with beveled edges)	Size	7mm
Flavor		Imprint Code	C7;NVR
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0078-1021-84	1 in 1 CARTON	04/05/2022	
1	NDC:0078-1021-51	28 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0078-1021-91	1 in 1 CARTON	04/05/2022	
2	NDC:0078-1021-90	28 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215039	04/05/2022	

VIJOICE

alpelisib tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0078-1028
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPELISIB (UNII: 08W5N2C97Q) (ALPELISIB - UNII:08W5N2C97Q)	ALPELISIB	125 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	YELLOW (dark)	Score	no score
Shape	OVAL (ovaloid and curved with beveled edges)	Size	13mm
Flavor		Imprint Code	Y7;NVR
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0078-1028-84	1 in 1 CARTON	04/05/2022	
1	NDC:0078-1028-51	28 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0078-1028-91	1 in 1 CARTON	04/05/2022	
2	NDC:0078-1028-90	28 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215039	04/05/2022	

VIJOICE

alpelisib kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0078-1035
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0078-1035-02	2 in 1 CARTON	04/05/2022	
1	NDC:0078-1035-61	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0078-1035-92	2 in 1 CARTON	04/05/2022	
2	NDC:0078-1035-94	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1		14
Part 2		14

Part 1 of 2

VIJOICE

alpelisib tablet

Product Information

Item Code (Source)	NDC:0078-1133
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPELISIB (UNII: 08W5N2C97Q) (ALPELISIB - UNII:08W5N2C97Q)	ALPELISIB	200 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

MAGNESIUM STEARATE (UNII: 70097M6I30)

TALC (UNII: 7SEV7J4R1U)

POLYETHYLENE GLYCOL 4000 (UNII: 4R4HF16D95)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

FERRIC OXIDE RED (UNII: 1K09F3G675)

Product Characteristics

Color	YELLOW (pale)	Score	no score
Shape	OVAL (ovaloid and curved with beveled edges)	Size	16mm
Flavor		Imprint Code	CL7;NVR
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215039	04/05/2022	

Part 2 of 2

VIJOICE

alpelisib tablet

Product Information

Item Code (Source)	NDC:0078-1140
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPELISIB (UNII: 08W5N2C97Q) (ALPELISIB - UNII:08W5N2C97Q)	ALPELISIB	50 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HF16D95)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

FERRIC OXIDE RED (UNII: 1K09F3G675)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

Product Characteristics

Color	YELLOW (light)	Score	no score
Shape	ROUND (round and curved with beveled edges)	Size	7mm
Flavor		Imprint Code	C7;NVR
Contains			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215039	04/05/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA215039	04/05/2022	

VIJOICE

alpelisib granule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0078-1175
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALPELISIB (UNII: 08W5N2C97Q) (ALPELISIB - UNII:08W5N2C97Q)	ALPELISIB	50 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0078-1175-51	28 in 1 CARTON	04/24/2024	
1	NDC:0078-1175-19	1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA218466	04/24/2024	

Labeler - Novartis Pharmaceuticals Corporation (002147023)

Revised: 11/2025

Novartis Pharmaceuticals Corporation