## IODENT ORAL ANALGESIC- benzocaine gel United Exchange Corp.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### Iodent Oral Analgesic Gel 0.42 oz 490 (ZDP 2020)

Active ingredient Purpose

Benzocaine 20%......Oral Pain Reliever

Uses

for the temporary relief of pain due to:

- toothache
- canker sores
- gum pain
- denture pain
- cheek bites

### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert**: do not use this product if your child has a history of allergy or local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- more than directed
- for more than 7 days unless directed by a doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness does not go away
- swelling, rash or fever develops

**If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children:

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Adults and children Remove cap and cut tip of tube. Apply a small amount of product to the cavity and 2 years of age and around gum surrounding the teeth. Use up to 4 times daily or as directed by a

older. dentist or doctor.

Children under 12 years of age. Should be supervised in the use of this product.

Children under 2 years of age. Do not use.

#### Other information

• store at room temperature 20° to 25°C (68° to 77°)

- this preparation is intended for use in cases of toothaches, only as a temporary expedient until a dentist or doctor can be consulted
- do not use continuously

#### Inactive ingredients

glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

Distributed by:

United Exchange Corp.

5836 Corporate Ave.

Cypress, CA 90630 USA

1-800-814-8028

Made in China



#### **MAXIMUM STRENGTH**

# Oral Analgesic Gel

Benzocaine 20% (Oral Pain Reliever)

#### Fast Relief From:

• Toothache • Canker Sores • Gum Pain • Denture Pain • Cheek Bites

#### Fast Relief From:

- Toothache
   Canker Sores
- Gum Pain
   Denture Pain
   Cheek Bites

**MAXIMUM STRENGTH** 

SAFETY-SEALED TUBE TIP

# Oral Analgesic Gel

Benzocaine 20% (Oral Pain Reliever)

NET WT 0.42 OZ (11.9 G)



Distributed By: UNITED EXCHANGE CORP. Cypress, CA 90630 USA 1-888-645-8204

Made In China 490-ZDP-BX-R3 | OC-MD-ZDP105-R3

### Call 1-888-645-8204 Monday-Friday 9AM-5PM (PST)

Inactive ingredients glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

°C8-07 and more managed as the strice and more more more more of 1-68-07 and 1-68-07 and

years of age.	
Children under 2	Do not use.
years of age.	product.
Children under 12	Should be supervised in the use of this
Adults and children 2 years of age and blder.	Remove cap and cut tip of tube. Apply a small amount of product to the cavity and around gum aurrounding the feeth. Use the product of the capital of the capital of dentist or doctor.
DIRECTIONS	

ontact a Poison Control Center (1-800-222-1222) right away yeep out of reach of children. If swallowed, get medical help or

Dung Facts (continued)

Stop use and sek a doctor if ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain or redness does not go away ■ swey use swey.

unless directed by a doctor 

for teething 

in children under

Figure 2 and deserver and the control of the contro

methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen camed in blood. This can occur even if you have used this product before opinemia warning: Use of this product may cause

**Ness ■** tor the temborary relief of pein de to: ■ cheek bites

.Oral pain reliever Renzocaine 20%. Purpose Active ingredient Drug Facts



#### **IODENT ORAL ANALGESIC**

benzocaine gel

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65923-094	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BENZO CAINE (UNII: U3RS Y48 JW5) (BENZO CAINE - UNII: U3RS Y48 JW5)	BENZOCAINE	200 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
GLYCERIN (UNII: PDC6A3C0OX)		
PEPPERMINT OIL (UNII: AV092KU4JH)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6 D95)		
SORBITOL (UNII: 506T60A25R)		
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)		

Packaging			
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1 NDC:65923-094-11	1 in 1 CARTON	0 1/0 2/20 18	
1	11.9 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	0 1/0 2/20 18	

### Labeler - United Exchange Corp. (840130579)

Revised: 3/2020 United Exchange Corp.