

**QC STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid  
QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION)**

-----  
**Quality Choice Max Strength Bismuth Subsalsicylate 525mg**

**Active ingredient (in each 15 mL)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever/Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - upset stomach
  - nausea
  - gas
  - belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## **Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- shake well before use
- For accurate dosing, use dose cup
- **adults and children 12 years and over:**
  - 1 dose (2 tbsp. or 30 ml) every 1 hour as needed
  - **do not exceed 4 doses (8tbsp or 120 mL) in 24 hours**
  - use until diarrhea stops but not more than 2 days
- **children under 12 years:**ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

## **Other information**

- **each tbsp. contains:**sodium 10 mg
- low sodium
- sugar free
- salicylate 236 mg
- protect from freezing
- avoid excessive heat (more than 104°F or 40°C)
- **TAMPER EVIDENT: Do not use if imprinted shrink band is missing or broken**

## **Inactive ingredients**

Carboxymethylcellulose sodium, flavor, microcrystalline cellulose, potassium hydroxide,

potassium sorbate, purified water, red 22, red 28, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum.

## **Principal Display Panel**

Compare to the active ingredient in **Pepto-Bismol® Maximum Strength**

Pink Bismuth

Maximum Strength

Stomach Relief

**Bismuth Subsalicylate 525 mg**

**Soothing Relief from:**

Upset Stomach

Indigestion Nausea

Heartburn

Diarrhea

Original Flavor

FL OZ (mL)

QC SATISFACTION GURANTEED

**\*This product is not manufactured or distributed by The Procter & Gamble, distributors of Pepto-Bismol®.**

**Distributed by C.D.M.A., Inc.**

**43157 W. Nine Mile**

**Novi, MI 48376-0995**

**[www.qualitychoice.com](http://www.qualitychoice.com)**

**Questions: 248-449-9300**

**Drug facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- shake well before use
- for accurate dosing, use dose cup
- adults and children 12 years and over:** 1 dose (2 tsp or 30 ml) every 1/2 to 1 hour as needed
- do not exceed 4 doses (8 Tbsp or 120 ml) in 24 hours**
- use until diarrhea stops, but not more than 2 days
- children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other Information**

- each tsp contains: sodium 10 mg
- low sodium ■ sugar free ■ salicylate 236 mg
- protect from freezing
- avoid excessive heat (more than 104°F or 40°C)
- TAMPER EVIDENT: Do not use if imprinted shrink band is missing or broken.**

**Inactive ingredients**  
 carboxymethylcellulose sodium, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, red 22, red 28, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

70708QCLR

LOT: UNVARNISHED AREA

EXP:

Distributed by C.D.M.A., Inc.  
 43157 W. Nine Mile  
 Novi, MI 48376-0995  
 www.qualitychoice.com  
 Questions: 248-449-9300

6 35515 98132 4

QC QUALITY CHOICE

NDC 63868-304-08

\*Compare to the active ingredient in PEPTO-BISMOL® Maximum Strength

# Pink Bismuth

## Maximum Strength

### Stomach Relief

Bismuth Subsalicylate 525 mg

Soothing Relief from:  
 Upset Stomach  
 Indigestion Nausea  
 Heartburn  
 Diarrhea

Original Flavor

8 FL OZ 236 mL



**Drug Facts**

**Active ingredient (In each 15 ml Tablespoon)** Bismuth subsalicylate 525 mg

**Purpose** Upset stomach reliever and antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ upset stomach ■ nausea ■ gas ■ belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

\* This product is not manufactured or distributed by Procter & Gamble, distributors of Pepto-Bismol®.

## Package Label

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- shake well before use
- for accurate dosing, use dose cup
- adults and children 12 years and over:** 1 dose (2 tsp or 30 ml) every 1 hour as needed
- do not exceed 4 doses (8 tsp or 120 ml) in 24 hours**
- use until diarrhea stops, but not more than 2 days
- children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- each tsp contains: sodium 10 mg
- low sodium ■ sugar free ■ salicylate 236 mg
- protect from freezing
- avoid excessive heat (more than 104°F or 40°C)
- TAMPER EVIDENT: Do not use if imprinted shrink band is missing or broken.**

**Inactive ingredients**  
 carboxymethylcellulose sodium, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, red 22, red 28, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

70712QCLR

LOT: UNVARNISHED AREA

EXP:

Distributed by C.D.M.A., Inc.  
 43157 W. Nine Mile  
 Novi, MI 48376-0995  
 www.qualitychoice.com  
 Questions: 248-449-9300

6 35515 97000 7

QC QUALITY CHOICE

NDC 63868-304-12

\*Compare to the active ingredient in PEPTO-BISMOL® Maximum Strength

# Pink Bismuth

## Maximum Strength

### Stomach Relief

Bismuth Subsalicylate 525 mg

Soothing Relief from:  
 Upset Stomach  
 Indigestion Nausea  
 Heartburn  
 Diarrhea

Original Flavor

12 FL OZ 354 mL



**Drug Facts**

**Active ingredient (In each 15 ml Tablespoon)** Bismuth subsalicylate 525 mg

**Purposes** Upset stomach reliever and antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ upset stomach ■ nausea ■ gas ■ belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

\* This product is not manufactured or distributed by Procter & Gamble, distributors of Pepto-Bismol®.

## QC STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:63868-304

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>DIMETHICONE, UNSPECIFIED</b> (UNII: 92RU3N3Y10)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	pink (viscous)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-304-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/16/2019	
2	NDC:63868-304-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/16/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	04/16/2019	

**Labeler** - QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION) (011920774)

Revised: 12/2025

QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION)