

PHARBEST REGULAR STRENGTH ASPIRIN- aspirin tablet
Pharbest Pharmaceuticals, Inc

Drug Facts

Active ingredient (in each tablet)

Aspirin 325mg (NSAID)[†]

[†]nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscle pain
- toothache
- menstrual pain
- colds
- minor pains of arthritis
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding •feel faint •vomit blood •have bloody or black stools •have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

If pregnant or breast-feeding

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose

adults and children 12 years and over	take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)
- avoid excess heat and moisture

Inactive ingredient

corn starch

Questions or comments?

(866)562-2756 (Mon-Fri 8 AM to 4 PM EST)

PHARBEST

NDC 16103-365-11

Manufactured in the USA

***Compare to the active ingredient in Genuine Bayer® Aspirin**


Aspirin

Regular Strength

Pain Reliever/Fever Reducer (NSAID)

1000 TABLETS

Aspirin, 325 mg each

PHARBEST  **Manufactured in the USA**
NDC 16103-365-11

Compare to the active ingredient in Genuine Bayer® Aspirin

Aspirin

Regular Strength

Pain Reliever/Fever Reducer (NSAID)

1000 TABLETS
Aspirin, 325 mg each

THIS PACKAGE FOR HOUSEHOLD WITHOUT YOUNG CHILDREN

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Aspirin, 325 mg (NSAID) Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to: ■ headache ■ muscle pain ■ toothache ■ menstrual pain ■ colds ■ minor pain of arthritis ■ temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from influenza or chickenpox or other viral infections should not take this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) drug ■ take aspirin or other drugs such as ibuprofen or other NSAIDs ■ take aspirin, naproxen, or other pain relievers 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy. Needs change rapidly during pregnancy. Do not stop use if you are breastfeeding. The use of this product during breastfeeding may cause problems for the child or infant. For more information, call the Poison Control Center (1-800-222-1222).

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222).

Directions

- drink a full glass of water with each dose

adults and children 12 years and over	take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F) ■ avoid excess heat and moisture


Inactive ingredient corn starch

Questions or comments? (866) 562-2756 (Mon-Fri 8 AM to 4 PM EST)

This product is not manufactured or distributed by Bayer. It is a generic equivalent of Bayer's Aspirin.

Manufactured by:
Pharbest Pharmaceuticals Inc.,
Farmingdale, NY 11735

LRev 06/20



0 16103-365-11 3

PHARBEST REGULAR STRENGTH ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:16103-365
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
Inactive Ingredients				
Ingredient Name		Strength		
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND (ROUND TABLET)	Size	10mm	
Flavor		Imprint Code	PH033	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16103-365-08	1 in 1 CARTON	01/12/2007	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:16103-365-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/12/2007	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	01/12/2007		

Labeler - Pharbest Pharmaceuticals, Inc (557054835)

Registrant - Pharbest Pharmaceuticals, Inc (557054835)

Establishment

Name	Address	ID/FEI	Business Operations
Pharbest Pharmaceuticals, Inc		557054835	analysis(16103-365) , manufacture(16103-365) , label(16103-365) , pack(16103-365)

Revised: 12/2024

Pharbest Pharmaceuticals, Inc