

**EFFERVESCENT ORANGE COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent**  
**H. E. Butt Grocery Company**

-----  
**HEB Effervescent Cold Relief**

**Active Ingredients**

Aspirin 325 mg (NSAID)\*

Chlorpheniramine maleate 2 mg

Phenylephrine Bitartrate 7.8 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Aspirin 325 mg (NSAID)\*. . . . .Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg . . . . .Antihistamine

Phenylephrine bitartrate 7.8 mg . . . . .Nasal decongestant

\*Nonsteroidal anti-inflammatory drug

**Uses**

temporarily relieves these symptoms of the common cold:

- fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose
- sinus congestion and pressure

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than

directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor before us if you have** ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- presently taking a prescription drug
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

### **When using this product**

- do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by a doctor.
- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does

not get better

- an allergic reaction occurs. Seek medical help right away.
- pain or nasal congestion gets worse or lasts more than 7 days.
- fever gets worse or lasts more than 3 days.
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

### **If pregnant or breastfeeding,**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

### **Keep out of reach of children**

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 ounces of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- *children under 12 years: ask a doctor*

### **Other information**

#### **each tablet contains:**

sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

### **Inactive ingredients**

acesulfame potassium, aspartame, citric acid, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?**

**Call (1-888-228-6937)**

**Package Label.Principle Display Panel**

FPO

**Drug Facts**

**Active ingredients**  
(in each effervescent tablet)  
Aspirin 325 mg (NSAID)  
Phenylephrine 2 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine 2 mg  
Aspirin 325 mg (NSAID)  
Phenylephrine 2 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine 2 mg  
Aspirin 325 mg (NSAID)  
Phenylephrine 2 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine 2 mg

**Purpose**  
Relief of minor aches and pains, runny nose, sneezing, nasal congestion, sore throat, headache, fever, and pain.

**Uses**  
Generally relieves these symptoms of the common cold:  
● sneezing ● nasal congestion ● sore throat ● headache ● minor aches and pains ● runny nose

**Warnings**  
● Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior, such as irritability, loss of appetite, unusual drowsiness, or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
● Allergy alert: Aspirin may cause a severe allergic reaction (anaphylaxis) in some people. ● Do not swallow. ● Do not use if you are allergic to aspirin, salicylates, or other drugs containing salicylates or non-steroidal anti-inflammatory drugs (NSAIDs), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older. ● Have had stomach ulcers or bleeding problems. ● Take other drugs containing aspirin, naproxen, or ibuprofen (NSAIDs) during naproxen, or ibuprofen (NSAIDs) use. ● Take other drugs containing aspirin, naproxen, or ibuprofen (NSAIDs) during naproxen, or ibuprofen (NSAIDs) use. ● Take other drugs containing aspirin, naproxen, or ibuprofen (NSAIDs) during naproxen, or ibuprofen (NSAIDs) use.

**Drug Facts (continued)**

Some throat warbling. If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ● if you have ever had an allergic reaction to any other pain reliever/fever reducer ● if you are allergic to aspirin ● if you are now taking a prescription medicine called thiazide (MIH) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MIH drug, if you do not know if your prescription drug contains an MIH. ● Ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if** you have ● glaucoma ● diabetes ● thyroid disease ● trouble urinating due to an enlarged prostate gland ● a breathing problem such as emphysema or chronic bronchitis ● been placed on a sodium-restricted diet.

**Ask a doctor or pharmacist before use if you are** ● taking sedatives or tranquilizers ● presently taking a prescription drug ● taking a prescription drug for anticoagulation (thinning the blood, diabetes, gout, or arthritis) ● using the product ● do not take more than 6 tablets daily and

**When using this product** ● do not take more than 6 tablets daily and

**do not use more than directed** ● you may get drowsy ● avoid alcohol drinks ● exactly may occur, especially in children ● alcohol, sedatives and tranquilizers may increase drowsiness ● be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding ● heartburn ● vomit blood ● have a body or black stools ● take more or for a longer time than directed.



## Effervescent Cold Plus

Chlorpheniramine Maleate / Antihistamine  
Phenylephrine Bitartrate / Nasal Decongestant  
Aspirin (NSAID) / Pain Reliever/Fever Reducer

Cold

20 EFFERESCENT TABLETS

Compare to Alka-Seltzer Plus® Cold Orange Zest active ingredients\*



## Effervescent Cold Plus

Chlorpheniramine Maleate / Antihistamine  
Phenylephrine Bitartrate / Nasal Decongestant  
Aspirin (NSAID) / Pain Reliever/Fever Reducer

Cold

- Relief of:
- Minor Aches and Pains
  - Headache
  - Runny Nose
  - Fever
  - Nasal Congestion
  - Sneezing
  - Sore Throat



Orange Flavor

20 EFFERESCENT TABLETS

### Drug Facts (continued)

Seek medical help right away

- pain or nasal congestion gets worse or lasts more than 7 days.
- fever gets worse or lasts more than 3 days.
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor



FPO

### Drug Facts (continued)

**Other information**  
each tablet contains:  
sodium 46.4 mg

- phenylephrine HCl: contains phenylephrine 9 mg per tablet
- store at room temperature (59°-86°F)

**Inactive ingredients**  
ascorbic acid, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments?**  
Call (1-888-228-6837)

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED



\*This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold Orange Zest.



PN: 87560 HE C2  
REV: 09/17

# EFFERVESCENT ORANGE COLD RELIEF

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-768
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE BITARTRATE</b> (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-768-20	10 in 1 CARTON	01/30/2018	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/30/2018	

**Labeler** - H. E. Butt Grocery Company (007924756)

Revised: 11/2024

H. E. Butt Grocery Company