

EFFERVESCENT ANTACID AND PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent
H. E. Butt Grocery Company

HEB Antacid & Pain Relief

Active ingredients

(in each effervescent tablet)

Aspirin 325 mg **(NSAID)***

Citric acid 1000 mg

Sodium bicarbonate (heat-treated) 1916 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Aspirin 325 mg **(NSAID)***.Pain reliever

Citric acid 1000 mg Antacid

Sodium bicarbonate (heat-treated) 1916 mg Antacid

*Nonsteroidal anti-inflammatory drug

Uses

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- pain alone (headaches or body aches and pains)

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or

nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if overly full from food or drink. Ask a doctor if severe stomach pain occurs after taking this product.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 10 days

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away.

■ pain lasts for more than 10 days or gets worse ■ new symptoms occur

■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ■ ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over (up to 60 years of age): 2 tablets every 4 hours - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours - do not exceed 4 tablets in 24 hours

Take this product any time - morning, noon or night - when you need relief from heartburn, acid indigestion, sour stomach with headache or body aches and pains. Place two (2) tablets in 4 oz. of water. Dissolve tablets completely. You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Other information

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59-86°F)

Inactive ingredients

■ none

Questions or comments?

1-888-228-6937

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL



EFFERVESCENT ANTACID AND PAIN RELIEF

aspirin, citric acid, sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-421
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	L421
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-421-68	18 in 1 CARTON	08/10/2011	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/10/2011	12/31/2026

Labeler - H. E. Butt Grocery Company (007924756)

Revised: 11/2024

H. E. Butt Grocery Company