


## **DILTIAZEM HYDROCHLORIDE- diltiazem hydrochloride capsule, extended release**

**Ingenus Pharmaceuticals, LLC**

-----

### **DESCRIPTION**

Diltiazem hydrochloride is a calcium ion cellular influx inhibitor (slow channel blocker or calcium antagonist). Chemically, diltiazem hydrochloride is (+)-5-[2-(Dimethylamino)ethyl]-*cis*-2,3-dihydro-3-hydroxy-2-(*p*-methoxyphenyl) -1,5-benzothiazepin-4(5*H*)-one acetate (ester) monohydrochloride. The chemical structure is:



Diltiazem hydrochloride, USP is a white to off-white crystalline powder with a bitter taste. It is soluble in water, methanol, and chloroform. It has a molecular weight of 450.98. Each diltiazem hydrochloride extended-release capsule, USP contains either 60 mg diltiazem hydrochloride, USP (equivalent to 55.1 mg diltiazem), 90 mg diltiazem hydrochloride, USP (equivalent to 82.7 mg diltiazem), or 120 mg diltiazem hydrochloride, USP (equivalent to 110.3 mg diltiazem).

Inactive ingredients: diethyl phthalate, gelatin, hypromellose, methacrylic acid and methyl methacrylate copolymer (1:2), sodium lauryl sulfate, polyethylene glycol, sugar spheres, talc, yellow iron oxide, titanium dioxide, polydextrose, triacetin, and FD&C Red No.40.

In addition, the black imprinting ink contains black iron oxide, potassium hydroxide, propylene glycol and shellac.

For oral administration.

FDA approved dissolution test specifications differ from USP.

FDA approved organic impurities and assay procedures differ from USP.

### **CLINICAL PHARMACOLOGY**

The therapeutic effects of diltiazem hydrochloride are believed to be related to its ability to inhibit the influx of calcium ions during membrane depolarization of cardiac and vascular smooth muscle.

### **Mechanism of Action**

Diltiazem hydrochloride extended-release capsules produce their antihypertensive effect primarily by relaxation of vascular smooth muscle and the resultant decrease in peripheral vascular resistance. The magnitude of blood pressure reduction is related to the degree of hypertension; thus, hypertensive individuals experience an antihypertensive effect, whereas there is only a modest fall in blood pressure in normotensives.

### **Hemodynamic and Electrophysiological Effects**

Like other calcium channel antagonists, diltiazem decreases sinoatrial and atrioventricular conduction in isolated tissues and has a negative inotropic effect in isolated preparations. In the intact animal, prolongation of the AH interval can be seen at higher doses.

In man, diltiazem prevents spontaneous and ergonovine-provoked coronary artery spasm. It causes a decrease in peripheral vascular resistance and a modest fall in blood pressure in normotensive individuals and, in exercise tolerance studies in patients with ischemic heart disease, reduces the heart rate-blood pressure product for any given workload. Studies to date, primarily in patients with good ventricular function, have not revealed evidence of a negative inotropic effect; cardiac output, ejection fraction, and left ventricular end diastolic pressure have not been affected. Increased heart failure has, however, been reported in occasional patients with preexisting impairment of ventricular function. There are as yet few data on the interaction of diltiazem and beta-blockers in patients with poor ventricular function. Resting heart rate is usually slightly reduced by diltiazem.

Diltiazem hydrochloride extended-release capsules produce antihypertensive effects both in the supine and standing positions. Postural hypotension is infrequently noted upon suddenly assuming an upright position. No reflex tachycardia is associated with the chronic antihypertensive effects. Diltiazem hydrochloride extended-release capsules decrease vascular resistance, increase cardiac output (by increasing stroke volume), and produce a slight decrease or no change in heart rate. During dynamic exercise, increases in diastolic pressure are inhibited, while maximum achievable systolic pressure is usually reduced. Heart rate at maximum exercise does not change or is slightly reduced. Chronic therapy with diltiazem hydrochloride produces no change or an increase in plasma catecholamines. No increased activity of the renin-angiotensin-aldosterone axis has been observed. Diltiazem hydrochloride extended-release capsules antagonize the renal and peripheral effects of angiotensin II. Hypertensive animal models respond to diltiazem with reductions in blood pressure and increased urinary output and natriuresis without a change in urinary sodium/potassium ratio.

Intravenous diltiazem hydrochloride in doses of 20 mg prolongs AH conduction time and AV node functional and effective refractory periods by approximately 20%. In a study involving single oral doses of 300 mg of diltiazem hydrochloride in six normal volunteers, the average maximum PR prolongation was 14% with no instances of greater than first-degree AV block. Diltiazem-associated prolongation of the AH interval is not more pronounced in patients with first-degree heart block. In patients with sick sinus syndrome, diltiazem significantly prolongs sinus cycle length (up to 50% in some cases).

Chronic oral administration of diltiazem hydrochloride in doses of up to 360 mg/day has resulted in small increases in PR interval, and on occasion produces abnormal prolongation (see WARNINGS).

## **Pharmacokinetics and Metabolism**

Diltiazem is well absorbed from the gastrointestinal tract and is subject to an extensive first-pass effect, giving an absolute bioavailability (compared to intravenous administration) of about 40%. Diltiazem hydrochloride undergoes extensive metabolism in which 2% to 4% of the unchanged drug appears in the urine. *In vitro* binding studies show diltiazem hydrochloride is 70% to 80% bound to plasma proteins. Competitive *in vitro* ligand binding studies have also shown diltiazem hydrochloride binding is not altered by therapeutic concentrations of digoxin, hydrochlorothiazide, phenylbutazone, propranolol, salicylic acid, or warfarin. The plasma elimination half-life following single or multiple drug administration is approximately 3.0 to 4.5 hours. Desacetyl diltiazem is also present in the plasma at levels of 10% to 20% of the parent drug and is 25% to 50% as potent a coronary vasodilator as diltiazem. Minimum therapeutic plasma levels of diltiazem hydrochloride appear to be in the range of 50 - 200 ng/mL. There is a departure from linearity when dose strengths are increased; the half-life is slightly increased with dose. A study that compared patients with normal hepatic function to patients with cirrhosis found an increase in half-life and a 69% increase in bioavailability in the hepatically impaired patients. A single study in nine patients with severely impaired renal function showed no difference in the pharmacokinetic profile of diltiazem compared to patients with normal renal function.

## **Diltiazem Hydrochloride Extended-Release Capsules (Twice-a-Day Dosage)**

A single 120 mg dose of the capsule results in detectable plasma levels within 2 to 3 hours and peak plasma levels at 6 to 11 hours. The apparent elimination half-life after single or multiple dosing is 5 to 7 hours. A departure from linearity similar to that observed with the diltiazem hydrochloride tablet is observed. As the dose of diltiazem hydrochloride extended-release capsules is increased from a daily dose of 120 mg (60 mg b.i.d.) to 240 mg (120 mg b.i.d.) daily, there is an increase in area-under-the-curve of 2.6 times. When the dose is increased from 240 mg to 360 mg daily, there is an increase in area-under-the-curve of 1.8 times. The average plasma levels of the capsule dosed twice daily at steady-state are equivalent to the tablet dosed four times daily when the same total daily dose is administered.

## **INDICATIONS AND USAGE**

Diltiazem hydrochloride extended-release capsules (Twice-a-Day Dosage) are indicated for the treatment of hypertension. They may be used alone or in combination with other antihypertensive medications, such as diuretics.

## **CONTRAINDICATIONS**

Diltiazem hydrochloride is contraindicated in (1) patients with sick sinus syndrome except in the presence of a functioning ventricular pacemaker, (2) patients with second- or third-degree AV block except in the presence of a functioning ventricular pacemaker, (3) patients with hypotension (less than 90 mm Hg systolic), (4) patients who have demonstrated hypersensitivity to the drug, and (5) patients with acute myocardial infarction and pulmonary congestion documented by X-ray on admission.

## **WARNINGS**

### **Cardiac Conduction**

Diltiazem hydrochloride prolongs AV node refractory periods without significantly prolonging sinus node recovery time, except in patients with sick sinus syndrome. This effect may rarely result in abnormally slow heart rates (particularly in patients with sick sinus syndrome) or second- or third-degree AV block (9 of 2,111 patients or 0.43%). Concomitant use of diltiazem with beta-blockers or digitalis may result in additive effects on cardiac conduction. A patient with Prinzmetal's angina developed periods of asystole (2 to 5 seconds) after a single dose of 60 mg of diltiazem. (See ADVERSE REACTIONS.)

### **Congestive Heart Failure**

Although diltiazem has a negative inotropic effect in isolated animal tissue preparations, hemodynamic studies in humans with normal ventricular function have not shown a reduction in cardiac index nor consistent negative effects on contractility (dp/dt). An acute study of oral diltiazem in patients with impaired ventricular function (ejection fraction  $24\% \pm 6\%$ ) showed improvement in indices of ventricular function without significant decrease in contractile function (dp/dt). Experience with the use of diltiazem hydrochloride in combination with beta-blockers in patients with impaired ventricular function is limited. Caution should be exercised when using this combination.

### **Hypotension**

Decreases in blood pressure associated with diltiazem hydrochloride therapy may occasionally result in symptomatic hypotension.

### **Acute Hepatic Injury**

Mild elevations of transaminases with and without concomitant elevation in alkaline phosphatase and bilirubin have been observed in clinical studies. Such elevations were usually transient and frequently resolved even with continued diltiazem treatment. In rare instances, significant elevations in enzymes such as alkaline phosphatase, LDH, SGOT, SGPT, and other phenomena consistent with acute hepatic injury have been noted. These reactions tended to occur early after therapy initiation (1 to 8 weeks) and have been reversible upon discontinuation of drug therapy. The relationship to diltiazem hydrochloride is uncertain in some cases, but probable in some. (See PRECAUTIONS.)

## **PRECAUTIONS**

### **General**

Diltiazem hydrochloride is extensively metabolized by the liver and excreted by the kidneys and in bile. As with any drug given over prolonged periods, laboratory parameters of renal and hepatic function should be monitored at regular intervals. The drug should be used with caution in patients with impaired renal or hepatic function. In subacute and chronic dog and rat studies designed to produce toxicity, high doses of diltiazem were associated with hepatic damage. In special subacute hepatic studies, oral doses of 125 mg/kg and higher in rats were associated with histological changes in the liver which were reversible when the drug was discontinued. In dogs, doses of 20 mg/kg were also associated with hepatic changes; however, these changes were reversible

with continued dosing.

Dermatological events (see ADVERSE REACTIONS) may be transient and may disappear despite continued use of diltiazem hydrochloride. However, skin eruptions progressing to erythema multiforme and/or exfoliative dermatitis have also been infrequently reported. Should a dermatologic reaction persist, the drug should be discontinued.

## **Drug Interactions**

Due to the potential for additive effects, caution and careful titration are warranted in patients receiving diltiazem hydrochloride concomitantly with any agents known to affect cardiac contractility and/or conduction (see WARNINGS). Pharmacologic studies indicate that there may be additive effects in prolonging AV conduction when using beta-blockers or digitalis concomitantly with diltiazem hydrochloride (see WARNINGS).

As with all drugs, care should be exercised when treating patients with multiple medications. Diltiazem is both a substrate and an inhibitor of the cytochrome P-450 3A4 enzyme system. Other drugs that are specific substrates, inhibitors, or inducers of this enzyme system may have a significant impact on the efficacy and side effect profile of diltiazem. Patients taking other drugs that are substrates of CYP450 3A4, especially patients with renal and/or hepatic impairment, may require dosage adjustment when starting or stopping concomitantly administered diltiazem in order to maintain optimum therapeutic blood levels.

## **Anesthetics**

The depression of cardiac contractility, conductivity, and automaticity as well as the vascular dilation associated with anesthetics may be potentiated by calcium channel blockers. When used concomitantly, anesthetics and calcium blockers should be titrated carefully.

## **Benzodiazepines**

Studies showed that diltiazem increased the AUC of midazolam and triazolam by 3- to 4-fold and the  $C_{max}$  by 2-fold, compared to placebo. The elimination half-life of midazolam and triazolam also increased (1.5- to 2.5-fold) during coadministration with diltiazem. These pharmacokinetic effects seen during diltiazem coadministration can result in increased clinical effects (e.g., prolonged sedation) of both midazolam and triazolam.

## **Beta-blockers**

Controlled and uncontrolled domestic studies suggest that concomitant use of diltiazem hydrochloride and beta-blockers is usually well tolerated, but available data are not sufficient to predict the effects of concomitant treatment in patients with left ventricular dysfunction or cardiac conduction abnormalities.

Administration of diltiazem hydrochloride concomitantly with propranolol in five normal volunteers resulted in increased propranolol levels in all subjects and bioavailability of propranolol was increased approximately 50%. *In vitro*, propranolol appears to be displaced from its binding sites by diltiazem. If combination therapy is initiated or withdrawn in conjunction with propranolol, an adjustment in the propranolol dose may be warranted (see WARNINGS).

### ***Buspirone***

In nine healthy subjects, diltiazem significantly increased the mean buspirone AUC 5.5-fold and  $C_{max}$  4.1-fold compared to placebo. The  $T_{1/2}$  and  $T_{max}$  of buspirone were not significantly affected by diltiazem. Enhanced effects and increased toxicity of buspirone may be possible during concomitant administration with diltiazem. Subsequent dose adjustments may be necessary during coadministration and should be based on clinical assessment.

### ***Carbamazepine***

Concomitant administration of diltiazem with carbamazepine has been reported to result in elevated serum levels of carbamazepine (40% to 72% increase), resulting in toxicity in some cases. Patients receiving these drugs concurrently should be monitored for a potential drug interaction.

### ***Cimetidine***

A study in six healthy volunteers has shown a significant increase in peak diltiazem plasma levels (58%) and area-under-the-curve (53%) after a 1-week course of cimetidine at 1,200 mg per day and a single dose of diltiazem 60 mg. Ranitidine produced smaller, nonsignificant increases. The effect may be mediated by cimetidine's known inhibition of hepatic cytochrome P-450, the enzyme system responsible for the first-pass metabolism of diltiazem. Patients currently receiving diltiazem therapy should be carefully monitored for a change in pharmacological effect when initiating and discontinuing therapy with cimetidine. An adjustment in the diltiazem dose may be warranted.

### ***Clonidine***

Sinus bradycardia resulting in hospitalization and pacemaker insertion has been reported in association with the use of clonidine concurrently with diltiazem. Monitor heart rate in patients receiving concomitant diltiazem and clonidine.

### ***Cyclosporine***

A pharmacokinetic interaction between diltiazem and cyclosporine has been observed during studies involving renal and cardiac transplant patients. In renal and cardiac transplant recipients, a reduction of cyclosporine dose ranging from 15% to 48% was necessary to maintain cyclosporine trough concentrations similar to those seen prior to the addition of diltiazem. If these agents are to be administered concurrently, cyclosporine concentrations should be monitored, especially when diltiazem therapy is initiated, adjusted or discontinued. The effect of cyclosporine on diltiazem plasma concentrations has not been evaluated.

### ***Digitalis***

Administration of diltiazem hydrochloride with digoxin in 24 healthy male subjects increased plasma digoxin concentrations approximately 20%. Another investigator found no increase in digoxin levels in 12 patients with coronary artery disease. Since there have been conflicting results regarding the effect of digoxin levels, it is recommended that digoxin levels be monitored when initiating, adjusting, and discontinuing diltiazem hydrochloride therapy to avoid possible over- or under-digitalization (see WARNINGS).

## ***Ivabradine***

Concurrent use of diltiazem increases exposure to ivabradine and may exacerbate bradycardia and conduction disturbances. Avoid concomitant use of ivabradine and diltiazem.

## ***Quinidine***

Diltiazem significantly increases the AUC ( $0 \rightarrow \infty$ ) of quinidine by 51%,  $T_{1/2}$  by 36%, and decreases its  $CL_{oral}$  by 33%. Monitoring for quinidine adverse effects may be warranted and the dose adjusted accordingly.

## ***Rifampin***

Coadministration of rifampin with diltiazem lowered the diltiazem plasma concentrations to undetectable levels. Coadministration of diltiazem with rifampin or any known CYP3A4 inducer should be avoided when possible, and alternative therapy considered.

## ***Statins***

Diltiazem is an inhibitor of CYP3A4 and has been shown to increase significantly the AUC of some statins. The risk of myopathy and rhabdomyolysis with statins metabolized by CYP3A4 may be increased with concomitant use of diltiazem. When possible, use a non-CYP3A4-metabolized statin together with diltiazem; otherwise, dose adjustments for both diltiazem and the statin should be considered along with close monitoring for signs and symptoms of any statin related adverse events.

In a healthy volunteer cross-over study (N = 10), coadministration of a single 20 mg dose of simvastatin at the end of a 14 day regimen with 120 mg BID diltiazem extended-release resulted in a 5-fold increase in mean simvastatin AUC versus simvastatin alone. Subjects with increased average steady-state exposures of diltiazem showed a greater fold increase in simvastatin exposure. Computer-based simulations showed that at a daily dose of 480 mg of diltiazem, an 8- to 9-fold mean increase in simvastatin AUC can be expected. If coadministration of simvastatin with diltiazem is required, limit the daily doses of simvastatin to 10 mg and diltiazem to 240 mg.

In a ten-subject randomized, open label, 4-way cross over study, co-administration of diltiazem (120 mg bid, diltiazem SR for 2 weeks) with single 20 mg dose of lovastatin resulted in 3- to 4- fold increase in mean lovastatin AUC and  $C_{max}$  versus lovastatin alone. In the same study, there was no significant change in 20 mg single dose pravastatin AUC and  $C_{max}$  during diltiazem co-administration. Diltiazem plasma levels were not significantly affected by lovastatin or pravastatin.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

A 24-month study in rats and a 21-month study in mice showed no evidence of carcinogenicity. There was also no mutagenic response in *in vitro* bacterial tests. No intrinsic effect on fertility was observed in rats.

## **Pregnancy**

Reproduction studies have been conducted in mice, rats, and rabbits. Administration of doses ranging from five to ten times greater (on a mg/kg basis) than the daily

recommended therapeutic dose has resulted in embryo and fetal lethality. These doses, in some studies, have been reported to cause skeletal abnormalities. In the perinatal/postnatal studies, there was some reduction in early individual pup weights and survival rates. There was an increased incidence of stillbirths at doses of 20 times the human dose or greater.

There are no well controlled studies in pregnant women; therefore, use diltiazem hydrochloride in pregnant women only if the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

Diltiazem is excreted in human milk. One report suggests that concentrations in breast milk may approximate serum levels. If use of diltiazem hydrochloride is deemed essential, an alternative method of infant feeding should be instituted.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **Geriatric Use**

Clinical studies of diltiazem did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## **ADVERSE REACTIONS**

Serious adverse reactions have been rare in studies carried out to date, but it should be recognized that patients with impaired ventricular function and cardiac conduction abnormalities have usually been excluded from these studies.

The adverse events described below represent events observed in clinical studies of hypertensive patients receiving either diltiazem hydrochloride tablets or diltiazem hydrochloride extended-release capsules, as well as experiences observed in studies of angina and during marketing. The most common events in hypertension studies are shown in a table with rates in placebo patients shown for comparison. Less common events are listed by body system; these include any adverse reactions seen in angina studies that were not observed in hypertension studies. In all hypertensive patients studied (over 900), the most common adverse events were edema (9%), headache (8%), dizziness (6%), asthenia (5%), sinus bradycardia (3%), flushing (3%), and first-degree AV block (3%). Only edema and perhaps bradycardia and dizziness were dose related. The most common events observed in clinical studies (over 2,100 patients) of angina patients and hypertensive patients receiving diltiazem hydrochloride tablets or diltiazem hydrochloride extended-release capsules were (i.e., greater than 1%) edema (5.4%), headache (4.5%), dizziness (3.4%), asthenia (2.8%), first-degree AV block (1.8%), flushing (1.7%), nausea (1.6%), bradycardia (1.5%), and rash (1.5%).



## Double Blind Placebo Controlled Hypertension Trials

Adverse	Diltiazem N = 315 No. pts (%)	Placebo N = 211 No. pts (%)
Headache	38 (12%)	17 (8%)
AV block first degree	24 (7.6%)	4 (1.9%)
Dizziness	22 (7%)	6 (2.8%)
Edema	19 (6%)	2 (0.9%)
Bradycardia	19 (6%)	3 (1.4%)
ECG abnormality	13 (4.1%)	3 (1.4%)
Asthenia	10 (3.2%)	1 (0.5%)
Constipation	5 (1.6%)	2 (0.9%)
Dyspepsia	4 (1.3%)	1 (0.5%)
Nausea	4 (1.3%)	2 (0.9%)
Palpitations	4 (1.3%)	2 (0.9%)
Polyuria	4 (1.3%)	2 (0.9%)
Somnolence	4 (1.3%)	—
Alk phos increase	3 (1%)	1 (0.5%)
Hypotension	3 (1%)	1 (0.5%)
Insomnia	3 (1%)	1 (0.5%)
Rash	3 (1%)	1 (0.5%)
AV block second degree	2 (0.6%)	—

In addition, the following events were reported infrequently (less than 1%) with diltiazem hydrochloride extended-release capsules or diltiazem hydrochloride tablets or have been observed in angina or hypertension trials:

**Cardiovascular:** Angina, arrhythmia, second- or third-degree AV block (see Conduction Warning), bundle branch block, congestive heart failure, syncope, tachycardia, ventricular extrasystoles.

**Nervous System:** Abnormal dreams, amnesia, depression, gait abnormality, hallucinations, nervousness, paresthesia, personality change, tremor.

**Gastrointestinal:** Anorexia, diarrhea, dry mouth, dysgeusia, mild elevations of SGOT, SGPT, and LDH (see Hepatic Warnings), thirst, vomiting, weight increase.

**Dermatological:** Petechiae, photosensitivity, pruritus, urticaria.

**Other:** Amblyopia, CPK increase, dyspnea, epistaxis, eye irritation, hyperglycemia, hyperuricemia, impotence, muscle cramps, nasal congestion, nocturia, osteoarticular pain, sexual difficulties, tinnitus.

The following post-marketing events have been reported infrequently in patients receiving diltiazem hydrochloride: acute generalized exanthematous pustulosis, allergic reactions, alopecia, angioedema (including facial or periorbital edema), asystole, erythema multiforme (including Stevens-Johnson Syndrome, toxic epidermal necrolysis), extrapyramidal symptoms, gingival hyperplasia, hemolytic anemia, increased bleeding time, leukopenia, photosensitivity (including lichenoid keratosis and hyperpigmentation at sun-exposed skin areas), purpura, retinopathy, myopathy, and thrombocytopenia.

There have been observed cases of a generalized rash, some characterized as leukocytoclastic vasculitis. In addition, events such as myocardial infarction have been observed which are not readily distinguishable from the natural history of the disease in these patients. A definitive cause and effect relationship between these events and diltiazem hydrochloride therapy cannot yet be established. Exfoliative dermatitis (proven by rechallenge) has also been reported.

## **OVERDOSAGE**

The oral LD<sub>50</sub>'s in mice and rats range from 415 to 740 mg/kg and from 560 to 810 mg/kg, respectively. The intravenous LD<sub>50</sub>'s in these species were 60 and 38 mg/kg, respectively. The oral LD<sub>50</sub> in dogs is considered to be in excess of 50 mg/kg, while lethality was seen in monkeys at 360 mg/kg.

The toxic dose in man is not known. Due to extensive metabolism, blood levels after a standard dose of diltiazem can vary over tenfold, limiting the usefulness of blood levels in overdose cases.

There have been reports of diltiazem overdose in doses ranging from 1 g to 18 g. Of cases with known outcome, most patients recovered and in cases with a fatal outcome, the majority involved multiple drug ingestion.

Events observed following diltiazem overdose included bradycardia, hypotension, heart block, and cardiac failure. Most reports of overdose described some supportive medical measure and/or drug treatment. Bradycardia frequently responded favorably to atropine, as did heart block, although cardiac pacing was also frequently utilized to treat heart block. Fluids and vasopressors were used to maintain blood pressure and in cases of cardiac failure inotropic agents were administered. In addition, some patients received treatment with ventilatory support, gastric lavage, activated charcoal, and/or intravenous calcium.

The effectiveness of intravenous calcium administration to reverse the pharmacological effects of diltiazem overdose has been inconsistent. In a few reported cases, overdose with calcium channel blockers associated with hypotension and bradycardia that was initially refractory to atropine became more responsive to atropine after the patients received intravenous calcium. In some cases, intravenous calcium has been administered (1 g calcium chloride or 3 g calcium gluconate) over 5 minutes and repeated every 10-20 minutes as necessary. Calcium gluconate has also been administered as a continuous infusion at a rate of 2 g per hour for 10 hours. Infusions of calcium for 24 hours or more may be required. Patients should be monitored for signs of hypercalcemia.

In the event of overdosage or exaggerated response, appropriate supportive measures should be employed in addition to gastrointestinal decontamination. Diltiazem does not appear to be removed by peritoneal or hemodialysis. Limited data suggest that plasmapheresis or charcoal hemoperfusion may hasten diltiazem elimination following overdose. Based on the known pharmacological effects of diltiazem and/or reported clinical experiences the following measures may be considered:

### **Bradycardia**

Administer atropine (0.6 to 1 mg). If there is no response to vagal blockade, administer

isoproterenol cautiously.

### **High-Degree AV Block**

Treat as for bradycardia above. Fixed high degree AV block should be treated with cardiac pacing.

### **Cardiac Failure**

Administer inotropic agents (isoproterenol, dopamine, or dobutamine) and diuretics.

### **Hypotension**

Vasopressors (e.g., dopamine or norepinephrine bitartrate).

Actual treatment and dosage should depend on the severity of the clinical situation and the judgment and experience of the treating physician.

**To report SUSPECTED ADVERSE REACTIONS, contact Ingenus Pharmaceuticals, LLC Toll-Free at 1-877-748-1970 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **DOSAGE AND ADMINISTRATION**

Dosages must be adjusted to each patient's needs, starting with 60 to 120 mg twice daily. Maximum antihypertensive effect is usually observed by 14 days of chronic therapy; therefore, dosage adjustments should be scheduled accordingly. Although individual patients may respond to lower doses, the usual optimum dosage range in clinical trials was 240 to 360 mg/day.

Diltiazem hydrochloride extended-release capsules have an additive antihypertensive effect when used with other antihypertensive agents. Therefore, the dosage of diltiazem hydrochloride extended-release capsules or the concomitant antihypertensives may need to be adjusted when adding one to the other. See WARNINGS and PRECAUTIONS regarding use with beta-blockers.

## **HOW SUPPLIED**

Diltiazem Hydrochloride Extended-Release Capsules, USP (Twice-a-Day Dosage) are available containing either 60 mg diltiazem hydrochloride, USP (equivalent to 55.1 mg diltiazem), 90 mg diltiazem hydrochloride, USP (equivalent to 82.7 mg diltiazem), or 120 mg diltiazem hydrochloride, USP (equivalent to 110.3 mg diltiazem).

The 60 mg capsules are hard-shell gelatin capsules with a medium orange opaque cap and white opaque body filed with white to off-white beads. The capsules are imprinted with "60 mg" in black ink on the cap and "NL2202" in black ink on the body. They are available as follows:

NDC 50742-564-01    bottles of 100 capsules

NDC 50742-564-05    bottles of 500 capsules

The 90 mg capsules are hard-shell gelatin capsules with a medium orange opaque cap and rich yellow opaque body filed with white to off-white beads. The capsules are

imprinted with “90 mg” in black ink on the cap and “NL2203” in black ink on the body. They are available as follows:

NDC 50742-565-01    bottles of 100 capsules  
NDC 50742-565-05    bottles of 500 capsules

The 120 mg capsules are hard-shell gelatin capsules with a medium orange opaque cap and body filed with white to off-white beads. The capsules are imprinted with “120 mg” in black ink on the cap and “NL2204” in black ink on the body. They are available as follows:

NDC 50742-566-01    bottles of 100 capsules  
NDC 50742-566-05    bottles of 500 capsules

**Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]  
Avoid excessive humidity.**

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Manufactured for:  
Ingenus Pharmaceuticals LLC  
Orlando, FL 32811-7193

Made in China  
I0186  
Iss. 04/2025  
Rev.A



## **PRINCIPAL DISPLAY PANEL - 60 mg**

**NDC 50742-564-01**

**Diltiazem HCl  
Extended-Release  
Capsules, USP  
60 mg\***

**Twice-a-Day Dosage**

**Rx only 100 Capsules**

\*Each capsule contains  
diltiazem hydrochloride, USP 60 mg  
(equivalent to 55.1 mg as diltiazem).

**Usual Dosage:** See accompanying  
prescribing information.

**Keep this and all medication out of the  
reach of children.**

**Store at 20° to 25°C (68° to 77°F). [See**

## USP Controlled Room Temperature.]

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Avoid excess humidity.

Rev. A Iss. 04/2025 P0406

### Manufactured for:

Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in China.

FDA approved organic impurities and assay procedures differ from USP.

564-01 v1/04-2025

Manufactured for:  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in China.


ingenus

NDC 50742-564-01

# Diltiazem HCl Extended-Release Capsules, USP

**60 mg\***

Twice-a-Day Dosage



Rx only  
100 Capsules

\*Each capsule contains:  
diltiazem hydrochloride, USP 60 mg  
(equivalent to 55.1 mg as diltiazem)


**Usual Dosage:** See accompanying  
prescribing information.

Keep this and all medication out of the  
reach of children.

Store at 20° to 25°C (68° to 77°F); [see  
USP Controlled Room Temperature].

Dispense in a tight, light-resistant container  
as defined in the USP using a child-resistant  
closure. Keep container tightly closed.  
Avoid excess humidity.

P04060425



3 50742 56401 9

GTIN: 00350742564019

LOT : 000000000  
EXP : 0000/00

SN : 000000000000

564-01 v1/04-2025

Rev.A Iss. 04/2025 P0406

## PRINCIPAL DISPLAY PANEL - 90 mg

NDC 50742-565-01

**Diltiazem HCl  
Extended-Release  
Capsules, USP  
90 mg\***

**Twice-a-Day Dosage**

**Rx only 100 Capsules**

\*Each capsule contains:  
diltiazem hydrochloride, USP 90 mg  
(equivalent to 82.7 mg as diltiazem).

**Usual Dosage:** See accompanying  
prescribing information.

**Keep this and all medication out of the  
reach of children.**

**Store at 20° to 25°C (68° to 77°F). [See**

## **USP Controlled Room Temperature.]**

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Avoid excess humidity.

Rev. A Iss. 04/2025 P0408

### **Manufactured for:**

Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in China.

FDA approved organic impurities and assay procedures differ from USP.

565-01 v1/04-2025

Manufactured for:  
Ingenus Pharmaceuticals, LLC  
Orlando, FL 32811-7193  
Made in China.

ingenus

NDC 50742-565-01

**Diltiazem HCl  
Extended-Release  
Capsules, USP**

**90 mg\***

Twice-a-Day Dosage

90mg NL2203

Rx only

100 Capsules

\*Each capsule contains:  
diltiazem hydrochloride, USP 90 mg  
(equivalent to 82.7 mg as diltiazem)

Usual Dosage: See accompanying prescribing information.

Keep this and all medication out of the reach of children.

Store at 20° to 25°C (68° to 77°F); [see USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure. Keep container tightly closed. Avoid excess humidity.

P04080425

3 50742 56501 6

GTIN: 00350742565016

LOT: 000000000

EXP: 0000/00

SN: 000000000000

565-01 v1/04-2025

Rev. A Iss. 04/2025 P0408

## **PRINCIPAL DISPLAY PANEL - 120 mg**

**NDC 50742-566-01**

**Diltiazem HCl  
Extended-Release  
Capsules, USP  
120 mg\***

**Twice-a-Day Dosage**

**Rx only 100 Capsules**

\*Each capsule contains:  
diltiazem hydrochloride, USP 120 mg  
(equivalent to 110.3 mg as diltiazem).

Usual Dosage: See accompanying  
prescribing information.

**Keep this and all medication out of the  
reach of children.**

DILTIAZEM HYDROCHLORIDE

diltiazem hydrochloride capsule, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50742-564
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
DILTIAZEM HYDROCHLORIDE (UNII: OLH94387TE) (DILTIAZEM - UNII:EE92BBP03H)		DILTIAZEM HYDROCHLORIDE	60 mg
Inactive Ingredients			
Ingredient Name			Strength
DIETHYL PHTHALATE (UNII: UF064M00AF)			

<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2)</b> (UNII: 5KY68S2577)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SHELLAC</b> (UNII: 46N107B71O)	

### Product Characteristics

<b>Color</b>	orange (medium orange opaque) , WHITE (white opaque)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	60mg;NL2202
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-564-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	
2	NDC:50742-564-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA218032	07/03/2025	

## DILTIAZEM HYDROCHLORIDE

diltiazem hydrochloride capsule, extended release

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50742-565
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety



Ingredient Name	Basis of Strength	Strength
<b>DILTIAZEM HYDROCHLORIDE</b> (UNII: OLVH94387TE) (DILTIAZEM - UNII:EE92BBP03H)	DILTIAZEM HYDROCHLORIDE	90 mg

Inactive Ingredients

Ingredient Name	Strength
<b>DIETHYL PHTHALATE</b> (UNII: UF064M00AF)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2)</b> (UNII: 5KY68S2577)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDWL1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SHELLAC</b> (UNII: 46N107B71O)	

Product Characteristics

<b>Color</b>	orange (medium orange opaque) , yellow (rich yellow opaque)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	90mg;NL2203
<b>Contains</b>			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-565-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	
2	NDC:50742-565-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA218032	07/03/2025	

DILTIAZEM HYDROCHLORIDE

diltiazem hydrochloride capsule, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:50742-566
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>DILTIAZEM HYDROCHLORIDE</b> (UNII: OLH94387TE) (DILTIAZEM - UNII:EE92BBP03H)	DILTIAZEM HYDROCHLORIDE	120 mg

Inactive Ingredients	
Ingredient Name	Strength
<b>DIETHYL PHTHALATE</b> (UNII: UF064M00AF)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:2)</b> (UNII: 5KY68S2577)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERROSOFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SHELLAC</b> (UNII: 46N107B71O)	

Product Characteristics			
Color	orange (medium orange opaque)	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	120mg;NL2204
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50742-566-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	
2	NDC:50742-566-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/03/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA218032	07/03/2025	

**Labeler -** Ingenus Pharmaceuticals, LLC (833250017)

**Registrant -** Novast Laboratories, Ltd. (527695995)

Establishment			
Name	Address	ID/FEI	Business Operations
Novast Laboratories, Ltd.		527695995	analysis(50742-564, 50742-565, 50742-566) , label(50742-564, 50742-565, 50742-566) , pack(50742-564, 50742-565, 50742-566) , manufacture(50742-564, 50742-565, 50742-566)