

**CLEAR LAX- polyethylene glycol 3350 powder, for solution**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. ClearLax<sup>®</sup> Drug Facts**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g

**Purpose**

Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product**

you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
- use once a day
- stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil pouch is open or broken

### **Inactive ingredients**

none

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

meijer®

ClearLax®

COMPARE TO MIRALAX® ACTIVE INGREDIENT

Polyethylene Glycol 3350 Powder for Solution Osmotic Laxative

RELIEVES OCCASIONAL CONSTIPATION/IRREGULARITY SOFTENS STOOL

Unflavored Powder Dissolves in Any Beverage!

Travel Size

20 PACKETS

EACH PACKET NET WT 0.5 OZ (17 g)

20 ONCE-DAILY DOSES

meijer.

# ClearLax®

Polyethylene Glycol 3350  
Powder for Solution  
Osmotic Laxative

RELIEVES OCCASIONAL  
CONSTIPATION/IRREGULARITY  
SOFTENS STOOL

Travel Size  
Packets

Unflavored  
Grit Free Powder

meijer.

# ClearLax®

Polyethylene Glycol 3350  
Powder for Solution  
Osmotic Laxative

RELIEVES OCCASIONAL  
CONSTIPATION/IRREGULARITY  
SOFTENS STOOL

NDC 41250-306-60

COMPARE TO  
MIRALAX®  
ACTIVE INGREDIENT\*

Unflavored Powder  
Dissolves in Any  
Beverage!

Travel Size  
20 PACKETS  
EACH PACKET NET WT 0.5 OZ (17 g)

20  
ONCE-DAILY  
DOSES

meijer.

# ClearLax®

Polyethylene Glycol 3350  
Powder for Solution  
Osmotic Laxative

RELIEVES OCCASIONAL  
CONSTIPATION/IRREGULARITY  
SOFTENS STOOL

NDC 41250-304-60

COMPARE TO  
MIRALAX®  
ACTIVE INGREDIENT\*

Unflavored Powder  
Dissolves in Any  
Beverage!

Travel Size  
20 PACKETS  
EACH PACKET  
NET WT 0.5 OZ (17 g)

20  
ONCE-DAILY  
DOSES

PD 5330680



30000 0E C1



**meijer.**

NDC 41250-306-60

# ClearLax<sup>®</sup>

COMPARE TO  
MIRALAX<sup>®</sup>  
ACTIVE INGREDIENT\*

**Polyethylene Glycol 3350  
Powder for Solution  
Osmotic Laxative**

**RELIEVES OCCASIONAL  
CONSTIPATION/IRREGULARITY  
SOFTENS STOOL**

Unflavored | Sugar-Free  
Dissolves in Any Beverage

20 Pre-Measured  
Unflavored, Single Doses

### Drug Facts

**Active ingredient (in each dose) Purpose**  
Polyethylene Glycol 3350, 17 g.....Osmotic Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

### Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older:
  - use once a day
  - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil pouch is open or broken

### Inactive ingredients

none

### Questions or comments?

1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark MIRALAX<sup>®</sup>.

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com



how2recycle.info



www.meijer.com/  
satisfaction



# CLEAR LAX

polyethylene glycol 3350 powder, for solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-306
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-306-02	238 g in 1 BOTTLE; Type 0: Not a Combination Product	10/06/2009	
2	NDC:41250-306-03	510 g in 1 BOTTLE; Type 0: Not a Combination Product	10/06/2009	
3	NDC:41250-306-04	850 g in 1 BOTTLE; Type 0: Not a Combination Product	05/27/2011	
4	NDC:41250-306-73	16 in 1 CARTON	10/07/2009	02/03/2013
4		17 g in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:41250-306-01	119 g in 1 BOTTLE; Type 0: Not a Combination Product	04/22/2015	
6	NDC:41250-306-60	20 in 1 CARTON	02/08/2024	
6		17 g in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/06/2009	

---

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 2/2024

Meijer Distribution Inc