DG HEALTH COUGH DM- dextromethorphan polistirex suspension Dolgencorp, LLC

Dolgencorp, LLC Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-888-309-9030

Package/Label Principal Display Panel

Compare to the active ingredient of Delsym®

Day or Night

Cough DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

Cough Suppressant

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

12 HOUR COUGH RELIEF

Alcohol free

Gluten free

Dosing cup included

Orange-Flavored Liquid

3 FL OZ (89 mL)



DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE GOODLETTSVILLE, TN 37072

Satisfaction Guaranteed! (888) 309-9030

Drug Facts

Active ingredient (in each 5 mL)
Destromethorphan polistrex equivalent to
30 mg dextromethorphan hydrobromide.......

.Cough suppr

Uses temporarily relieves

moough due to minor thread and bronchiel irritation as may occur with the
common cold or inhaled inflants

the impulse to cough to help you get to sleep

When implies to copy as new having a prescription monoamine existes inhibitor (MAO) (postular drags for depression, population or endotred conditions, or Parlamon's disease), or for 2 weeks after dopping the MAOI drag. If you do not know if your prescription drag contains an MAOI, ask a dockor or pharmacist when your prescription drag contains an MAOI, ask a dockor or pharmacist altergraphic for contains so dram melablesurflet, a suffer that may cause altergio-hype reactions.

Jake, a declore before use if you have a climate could be a declore before use if you have a contain an account of the cours with two much philagin (insular). Stop use and ask a doctor if may also a count of the cours with two much philagin (insular) are cough asts more than 7 days, cough comes back, or occurs with frever, read or headache that least. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health prefessional before use. Keep out of reach of children. In case of overdose, get mades help or confract a Poleon Control Center right away (1-800-222-1222).

- Directions

 shake bottle well before use
 measure only with dosing cup provided
 measure only with dosing cup provided
 med not use dosing cup with other products
 dose as follows or as directed by a doctor
- adults and children 12 years of age and over 10 mL every 12 hours, not to exceed 20 mL in 24 hours children 6 to under 5 mL every 12 hours, not to exceed 10 mL in 24 hours 2.5 mL every 12 hours, not to exceed 5 mL in 24 hours children 4 to under 6 years of age do not use

Other information

■ each 5 mL contains: sodium 5 mg ■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

Drug Facts (continued)

Inactive ingredients DBC Red a20 shimbur lege (Plator to BC Red 20 shimbur leke, DBC Velow #10 aluminum leke, DBC Velow #10 aluminum leke, DBC Velow #10 aluminum leke picerin, high fructose com syrup, methylpanben natural and artificial orange fewor, polycor beta 80 polyknyl acetake, polykore, propyparaben, purfiled velete, sodium metabkutifite, sodium polystyrene suffonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

"This product is not manufactured or distributed by

DOSING

Measure only with dosing oup provided. Do not use dosing oup with other products.		
Age (yr)	Dose	
12 years	10 mL	
to adult	EVERY 12 HOURS	
6 to	5 mL	
under 12	EVERY 12 HOURS	
4 to	2.5 mL	
under 6	EVERY 12 HOURS	
Lindor 4	Do not uso	

See back panel for full dosing directions.



Orange-Flavored Liquid Dosing cup included





Compare to the active ingredient of Delsym€*

Day or Night Cough DM

Dextromethorphan Polistirex **Extended-Release Oral Suspension**

Cough Suppressant

- Dosing cup included





Orange-Flavored Liquid

3 FL OZ (89 mL)



12 Hour Cough Relief Contains No Fever Reducer or I

ACTUAL SIZE

DG health

ough Suppressant

3 FL OZ (89 mL)





DG HEALTH COUGH DM

dextromethorphan polistirex suspension

Product Information

Route of Administration

HUMAN OTC DRUG NDC:55910-384 Product Type Item Code (Source)

ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength DEXTROMETHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH) **DEXTROMETHORPHAN** 30 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) HYDROBROMIDE in 5 mL

Inactive Ingredients

Ingredient Name Strength

POLISTIREX (UNII: 5H9W9GTW27)

D&C RED NO. 30 (UNII: 2S42T2808B)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A2I8 C7HI9 T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
PO VIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
NDC:55910-384-21	1 in 1 CARTON	07/02/2013	
1	89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	07/02/2013	

Labeler - Dolgencorp, LLC (068331990)

Revised: 6/2020 Dolgencorp, LLC