PRIMO TOPICAL ANESTHETIC- benzocaine gel Tri State Dental

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use. Use only as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Keep out of reach of children.

Other Information

Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

1-800-232-6874 or www.primodentalproducts.com

Manufactured for: Manufactured for: Principal Dental Product 845 Third Avenue 6th Floor New York, NY 100 800-232-6874 www.primodenta)22		Prime TOPICA TOPICA ANESTHETI (BENZOCAINE Gluten Free Net Content: 1 oz (IO L C G 20%)	B MADE IN THE USA	
directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly. • for teething • in children under 2 years of age When using this product	Avoid contact with eyes. If contact occurs, flush with water. Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast feeding, ask a physician before use. Keep out of reach of children.	Directions Apply only amount needed to Apply only amount needed to the oral mucosa to prevent or relieve pain. children under 2 vears of age: 	do not use Other information Store at 59° - 86°F (15 - 30°C). Protect from freezing Inactive ingredients flavoring, PEG 3350, PEG 400, sodium saccharin, water, Mav	contain blue #1, green #3, green #5, red #28, red #40,	yellow #5, (tartrazine), yellow #6, as a color additive. Questions or comments? 1-800-232-6874 or www.prinodentalproducts.com	
Drug Facts Active ingredients Benzocaine 200 mg (in each g) Benzocaine 200 mg (in each g) Purpose Oral Anesthetic Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures. Warnings	Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical	attention if you or a child in your care develops: • pale, gray or blue colored skin (cyanosis) • headache	 rapid heart rate shortness of breath dizziness or lightheadedness fatigue or lack of energy Allergy Alert: Do not use on patients with a history of allergies to local anesthetics 	such as procaine, butacaine, benzocaine or other "caine"	Anestnetics. Do not use • for more than 7 days unless	
PRIMO TOPICAL ANE benzocaine gel	STHETIC					
Product Information						
			(6	NDC		
Product Type	HUMAN OTC DRUG	ltem Code	(Source)	NDC:69509-006		
Route of Administration	DENTAL					
Active Ingredient/Active	Moiety					
Ingredient Name Basis of Stren				gth	n Strength	
BENZOCAINE (UNII: U3RSY48JW5)	BENZOCAINE	200 mg in 1 g				
Inactive Ingredients						
	Ingredient Name					
POLYETHYLENE GLYCOL 3350 (U		Strength				
POLYETHYLENE GLYCOL 400 (UI	NII: B697894SGQ)					
SACCHARIN SODIUM (UNII: SB8Z	JX40TY)					
FD&C YELLOW NO. 5 (UNII: 1753)	VB2F1M)					

FD&C YELLOW NO.	6 (UNII: H77VI	E193A8)					
FD&C RED NO. 3 (UN	III: PN2ZH5LO	QY)					
FD&C RED NO. 40 (U	JNII: WZB9127	XOA)					
FD&C BLUE NO. 1 (U	JNII: H3R47K3	TBD)					
FD&C GREEN NO. 3	(UNII: 3P3ONF	R601S)					
D&C RED NO. 28 (UN							
D&C GREEN NO. 5 (U	JNII: 8J6RDU8L	.9X)					
Product Charac	teristics						
Color purple		Score					
Shape			Size				
Flavor		GRAPE	Imprint	Imprint Code			
Contains							
Packaging							
# Item Code	Package Description			Ma	arketing Start Date	Marketin Dat	•
1 NDC:69509-006- 3030 g in 1 JAR; Type 0: Not a Combination Product				11/01/2018			
Marketing In	formati	on					
Marketing Category	Application Number or Monograp Citation		h	Marketing Start Date	Marketing End Date		
OTC monograph not final	part356			0	3/01/2014		
lillai							

Labeler - Tri State Dental (176048478)

Revised: 1/2022

Tri State Dental