

**CHILDRENS PAIN AND FEVER CHERRY FLAVOR- acetaminophen suspension**  
**Walgreen Company**

-----  
**Children's Pain & Fever**

**Drug Facts**

**Active ingredient**

(in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses** temporarily ■ reduces fever

■ relieves minor aches and pains due to:

■ the common cold ■ flu ■ headache ■ sore throat

■ toothache

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver

damage may occur if your child takes: ■ more than 5

doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin

reactions. Symptoms may include: ■ skin reddening

■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days,

is accompanied or followed by fever, headache, rash,

nausea or vomiting, consult a doctor promptly.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If your child is allergic to acetaminophen or any other of the inactive ingredients in this product.

**Ask a doctor before use if your child** has liver disease.

**Ask a doctor or pharmacist before use if your child**

**is** taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use.
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find the right dose on the chart below. If possible, use weight to dose, otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor.

---

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2 - 3 years	5 mL (1 tsp)
36-47	4 - 5 years	7.5 mL (1 1/2 tsp)
48-59	6 - 8 years	10 mL (2 tsp)
60-71	9 - 10 years	12.5. mL (2 1/2 tsp)
72-95	11 years	15 mL (3 tsp)

---

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**

- each 5 mL (1 tsp) contains; sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- Keep carton for full directions for use.

**Inactive ingredients** Anhydrous citric acid, butylparaben, FD&C red #40 flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and

carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

## **Product Label**

***Walgreens***

**Compare to Children's Tylenol®  
Oral Suspension active ingredient<sup>††</sup>**

**NDC 0363-0322-24**

**children's**

**Pain & Fever**

**ACETAMINOPHEN  
160 mg PER 5 mL**

**PAIN RELIEVER /  
FEVER REDUCER  
ORAL SUSPENSION**

**ALCOHOL FREE  
ASPIRIN FREE  
IBUPROFEN FREE**

**AGES  
2-11  
YEARS**

**CHERRY FLAVOR**

**4 FL OZ (118 mL)**

**This product is bot manufactured or distributed by Johnson & Johnson Consumer Inc., |McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.**

**ITEM 268958 W00000-0000-0**

**REV-1020**

**DO NOT USE IF PRINTED NECKBAND IS  
BROKEN OR MISSING**

**Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey**

**DISTRIBUTED BY WALGREEN CO.  
200 WILMONT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2020 Walgreen Co.**

**DOSAGE**

CUP PROVIDED  
BX-113

ORG1020-F

Carton



Bottle

*Walgreens*

NDC 0363-0322-24

**children's**  
**Pain & Fever**

ACETAMINOPHEN  
160 mg PER 5 mL  
PAIN RELIEVER /  
FEVER REDUCER  
ORAL SUSPENSION

Compare to  
Children's Tylenol®  
Oral Suspension  
active ingredient††

• Alcohol free

4 FL OZ (118 mL)

CHERRY  
FLAVOR

AGES  
2-11  
YEARS

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**Active ingredient (in each 5 mL = 1 teaspoonful)**      **Purpose**  
Acetaminophen 160 mg.....Pain reliever/fever reducer

**Uses:** Temporarily ■ reduces fever ■ relieves minor aches and pains due to:  
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings:** **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly. **Do not use:** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or a pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if your child has liver disease. Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product do not exceed recommended dose (see overdose warning) Stop use and ask a doctor if** ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. **Keep out of reach of children. Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-322-1223) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** ■ this product does not contain directions or complete warnings for adult use. ■ shake well before using ■ mL = milliliter; tsp = teaspoonful ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. **Other information** ■ each 5 mL (1 tsp) contains: sodium 3 mg ■ store between 20-25°C (68-77°F) ■ do not refrigerate

**Inactive ingredients** anhydrous citric acid, butylparaben, FD&C Red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

**Questions?** Call 1-877-798-5944      **LOT:**

† This product is not manufactured or distributed by Johnson & Johnson Consumer Inc. | McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY: WALGREEN CO.      ITEM 268958  
230 W. MONT RD., DEERFIELD, IL 60015      W00000-0000-0 LR-153 EXP:  
100% SATISFACTION GUARANTEED      ORG1020-F REV-1020  
walgreens.com ©2020 Walgreen Co.

res

## CHILDRENS PAIN AND FEVER CHERRY FLAVOR

acetaminophen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0322
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg in 5 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
BUTYLPARABEN (UNII: 3QPI1U3FV8)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor	CHERRY	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0322-24	1 in 1 CARTON	11/05/2020	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	11/05/2020		

**Labeler** - Walgreen Company (008965063)

**Registrant** - AptaPharma Inc. (790523323)

**Establishment**

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(0363-0322)

Revised: 12/2023

Walgreen Company