

HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet Family Dollar (FAMILY WELLNESS)

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives

- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have asthma
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis. or kidney disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask doctor if

- an allergic reaction occurs, Seek medical help right away
 - any new symptoms appear.
- you experience any of the following signs of stomach bleeding
 - feel faint

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen

these could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid, talc

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE THE ACTIVE INGREDIENTS IN EXCEDRIN® EXTRA STRENGTH†
EXTRA STRENGTH
HEADACHE RELIEF

Acetaminophen 250 mg - Pain reliever

Aspirin (NSAID) 250 mg - Pain reliever

Caffeine 65 mg - Pain reliever aid

CAPLETS

†This product is not manufactured or distributed by GSK Consumer Healthcare distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

Package Label

FAMILY
wellness™

COMPARE TO THE ACTIVE INGREDIENTS
IN EXCEDRIN® EXTRA STRENGTH*

EXTRA STRENGTH
HEADACHE RELIEF
Acetaminophen 250 mg - Pain reliever
Aspirin (NSAID) 250 mg - Pain reliever
Caffeine 65 mg - Pain reliever aid

NDC 55319-511-10



Actual Size



100 CAPLETS

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to
 - headache
 - a cold
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription

Drug Facts (continued)

NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have asthma
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- any new symptoms appear
- you experience any of the following signs or stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or loss of hearing occurs
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - painful area is red or swollen

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; do not take more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid, talc

Questions or comments?

Call 1-877-763-3635 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Excedrin® Extra Strength.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
500 VOLVO PARKWAY, CHESAPEAKE, VA 23320
NOT 100% SATISFIED? Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

SKU 090494



PLD-H881A F0007346
Product of India

Lot No.:

Exp. Date:

FAMILY WELLNESS Extra Strength Headache Relief Caplets

HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-511
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-511-10	1 in 1 BOX	11/27/2020	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	11/27/2020	

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)

Revised: 3/2024

Family Dollar (FAMILY WELLNESS)