

**CETIRIZINE HYDROCHLORIDE TABLETS, 5 MG - cetirizine hydrochloride tablet**  
**CETIRIZINE HYDROCHLORIDE TABLETS, 10 MG - cetirizine hydrochloride tablet**  
**CETIRIZINE HYDROCHLORIDE TABLETS, 10 MG (OVAL SHAPE) - cetirizine hydrochloride tablet**  
**MARKSANS PHARMA LIMITED**

-----  
**Allergy Relief**  
**Cetirizine Hydrochloride Tablets USP, 5 mg and 10 mg**

**Active ingredient (in each tablet)**

**For 5 mg:**

Cetirizine HCl USP 5 mg

**For 10 mg:**

Cetirizine HCl USP 10 mg

**For 10 mg (Oval Shape):**

Cetirizine HCl USP 10 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions****For 5 mg:**

<b>adults and children 6 years and over</b>	1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours
<b>adults 65 years and over</b>	1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours.
<b>children under 6 years of age</b>	ask a doctor
<b>consumers with liver or kidney disease</b>	ask a doctor

**For 10 mg:**

<b>adults and children 6 years and over</b>	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
<b>adults 65 years and over</b>	ask a doctor
<b>children under 6 years of age</b>	ask a doctor

<b>consumers with liver or kidney disease</b>	ask a doctor
---	--------------

### **Other information**

- store between 20° to 25°C (68° to 77°F)
- do not use if foil seal under cap is broken or missing

### **Inactive ingredients**

corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments? Call 1-877-376-4271**  
(weekdays 9 AM to 5 PM)

### **PRINCIPAL DISPLAY PANEL**

NDC 25000-219-03  
Cetirizine Hydrochloride Tablets USP, 5 mg  
30s count bottle label

NDC 25000-219-03

Original Prescription Strength  
Allergy

**Cetirizine Hydrochloride Tablets USP, 5 mg**

Antihistamine  
Indoor & Outdoor  
Allergies

**24** hour Relief of  
• Sneezing  
• Runny Nose  
• Itchy, Watery Eyes  
• Itchy Throat  
or Nose

Marksans

30 Tablets

**Active ingredient (in each tablet)**

Cetirizine HCl 5 mg.....Antihistamine

**Purpose**

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** liver or kidney disease. Your doctor

LOT :  
EXP :  
R27/23

should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask**

**a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions**

**adults and children 6 years and over:** 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours. **adults 65 years and over:** 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. **children under 6 years of age:** ask a doctor. **consumers with liver or kidney**

**disease:** ask a doctor. **Other information** ■ store between 20° to 25°C (68° to 77°F) ■ **Do not use if foil seal under cap is broken or missing.** **Inactive ingredients** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide

**Questions?** Call at 1-877-376-4271  
Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.  
Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-219-03  
Cetirizine Hydrochloride Tablets USP, 5 mg  
30s count carton label



LOT:

EXP:

**Drug Facts** (Continued)

■ be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

adults and children 6 years and over	1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours
adults 65 years and over	1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours.
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Drug Facts** (Continued)

- **Do not use if foil seal under cap is broken or missing.**

**Inactive ingredients:** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide

**Questions?**

Call at 1-877-376-4271

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.**

Compare to the active ingredient in Zyrtec® Tablets\*

NDC 25000-219-03

Original Prescription Strength  
Allergy

**Cetirizine Hydrochloride Tablets USP, 5 mg**

Antihistamine  
Indoor & Outdoor Allergies

24 Relief of Sneezing, Runny Nose, Itchy, Watery Eyes, Itchy Throat or Nose



30 Tablets

Take 1 to 2 tablets\* once daily depending on the severity of your symptoms. Adults 65 years and over: take only one tablet per day.



**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE. REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 5 mg	Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : G0/DRUGS/515

62/28

NDC 25000-219-08  
Cetirizine Hydrochloride Tablets USP, 5 mg  
100s count bottle label

NDC 25000-219-08

Original Prescription Strength  
Allergy

**Cetirizine Hydrochloride Tablets USP, 5 mg**

Antihistamine  
Indoor & Outdoor  
Allergies

**24** hour Relief of  
• Sneezing  
• Runny Nose  
• Itchy, Watery Eyes  
• Itchy Throat  
or Nose

Marksans

100 Tablets

**Active ingredient (in each tablet)**

Cetirizine HCl 5 mg.....Antihistamine

**Purpose**

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** liver or kidney disease. Your doctor

LOT :  
EXP :  
R27/23

should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask**

**a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions**

**adults and children 6 years and over:** 1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours. **adults 65 years and over:** 1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. **children under 6 years of age:** ask a doctor. **consumers with liver or kidney**

**disease:** ask a doctor. **Other information** ■ store between 20° to 25°C (68° to 77°F) ■ **Do not use if foil seal under cap is broken or missing.** **Inactive ingredients** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide

**Questions?** Call at 1-877-376-4271  
Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.  
Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-219-08  
Cetirizine Hydrochloride Tablets USP, 5 mg  
100s count carton label



**Drug Facts** (Continued)

- be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

<b>adults and children 6 years and over</b>	1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours
<b>adults 65 years and over</b>	1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours.
<b>children under 6 years of age</b>	ask a doctor
<b>consumers with liver or kidney disease</b>	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Drug Facts** (Continued)

- Do not use if foil seal under cap is broken or missing.**

**Inactive ingredients:** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide

**Questions?**  
Call at 1-877-376-4271

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.**

NDC 25000-219-08

Compare to the active ingredient in Zyrtec® Tablets\*

**Original Prescription Strength Allergy**

# Cetirizine Hydrochloride Tablets USP, 5 mg

**Antihistamine**

**Indoor & Outdoor Allergies**

**24 Relief of Sneezing, Runny Nose, Itchy, Watery Eyes, Itchy Throat or Nose**

Actual Size

**100 Tablets**

**Marksans**

Take 1 to 2 tablets\* once daily depending on the severity of your symptoms. Adults 65 years and over take only one tablet per day.

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : G0/DRUGS/515

62/28

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE. REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 5 mg	Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are taking** tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness

NDC 25000-219-14  
Cetirizine Hydrochloride Tablets USP, 5 mg  
1000s count bottle label

NDC 25000-219-14

Compare to the active ingredient in Zyrtec® Tablets†

# Original Prescription Strength Allergy

## Cetirizine Hydrochloride Tablets USP, 5 mg

### Antihistamine Indoor & Outdoor Allergies

- 24 hour** Relief of
- Sneezing
  - Runny Nose
  - Itchy, Watery Eyes
  - Itchy Throat or Nose

**THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN**

Take 1 to 2 tablets\* once daily depending on the severity of your symptoms  
\*Adults 65 and older take only one tablet per day



Actual Size

LOT :  
EXP :  
R27/23



# 1000 Tablets

### IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE.

#### Drug Facts

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 5 mg.....	Antihistamine

#### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

#### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. ➔

#### Drug Facts (Continued)

#### When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

#### If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

<b>adults and children 6 years and over</b>	1 tablet (5 mg) or 2 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 2 tablets (10 mg) in 24 hours
<b>adults 65 years and over</b>	1 tablet (5 mg) once daily; do not take more than 1 tablet (5 mg) in 24 hours. ➔

Drug Facts (Continued)

Drug Facts (Continued)

<b>children under 6 years of age</b>	ask a doctor	Manufactured for: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735, USA  Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa - 403722, India.  Mfg. Lic. No. : GO/DRUGS/515
<b>consumers with liver or kidney disease</b>	ask a doctor	
<b>Other information</b> ■ store between 20° to 25°C (68° to 77°F) ■ <b>Do not use if foil seal under cap is broken or missing.</b>		
<b>Inactive ingredients</b> lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide		
<b>Questions?</b> Call at 1-877-376-4271		
†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.		

NDC 25000-220-03  
 Cetirizine Hydrochloride Tablets USP, 10 mg  
 30s count bottle label

NDC 25000-220-03

Original Prescription Strength

Allergy

Cetirizine Hydrochloride Tablets USP, 10 mg

Antihistamine  
Indoor & Outdoor  
Allergies

24  
hour

Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

Marksans

30 Tablets

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg.....Antihistamine

**Purpose**

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat **Warnings** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** liver or kidney disease. Your doctor

LOT :  
EXP :  
R19/21

should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask**

**a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions**

**Adults and children 6 years and over:** one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between

20° to 25°C (68° to 77°F) ■ **Do not use if foil seal under cap is broken or missing.** **Inactive ingredients** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide **Questions?** Call at 1-877-376-4271

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA  
Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-220-03

Cetirizine Hydrochloride Tablets USP, 10 mg

30s count carton label



NDC 25000-220-08

Original Prescription Strength

Allergy

Cetirizine Hydrochloride Tablets USP, 10 mg

Antihistamine  
Indoor & Outdoor  
Allergies

24  
hour

Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

Marksans

100 Tablets

**Active ingredient (in each tablet)**

Cetirizine HCl 10 mg.....Antihistamine

**Purpose**

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat **Warnings** **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine. **Ask a doctor before use if you have** liver or kidney disease. Your doctor

LOT :  
EXP :  
R19/21

should determine if you need a different dose. **Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. **When using this product** ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. **Stop use and ask**

**a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding:** ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) **Directions**

**Adults and children 6 years and over:** one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. **Adults 65 years and over:** ask a doctor. **Children under 6 years of age:** ask a doctor. **Consumers with liver or kidney disease:** ask a doctor. **Other information** ■ store between

20° to 25°C (68° to 77°F) ■ **Do not use if foil seal under cap is broken or missing.** **Inactive ingredients** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide **Questions?** Call at 1-877-376-4271

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA  
Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-220-08  
Cetirizine Hydrochloride Tablets USP, 10 mg  
100s count carton label



:dX3  
:LOT:

**Drug Facts** (Continued)

■ be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding:**

■ if breast-feeding: not recommended  
■ if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

<b>Adults and children 6 years and over</b>	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
<b>Adults 65 years and over</b>	ask a doctor
<b>Children under 6 years of age</b>	ask a doctor
<b>Consumers with liver or kidney disease</b>	ask a doctor

**Other information**

■ store between 20° to 25°C (68° to 77°F)

**Drug Facts** (Continued)

■ **Do not use if foil seal under cap is broken or missing.**

**Inactive ingredients:** lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide

**Questions?**

Call at 1-877-376-4271

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.**

Compare to the active ingredient in Zyrtec® Tablets\*

NDC 25000-220-08

**Original Prescription Strength Allergy**

**Cetirizine Hydrochloride Tablets USP, 10 mg**

**Antihistamine Indoor & Outdoor Allergies**

**24 Relief of**  
• Sneezing  
• Runny Nose  
• Itchy, Watery Eyes  
• Itchy Throat or Nose



100 Tablets



**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE. REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 10 mg.....	Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : G0/DRUGS/515

1538723

NDC 25000-220-14  
Cetirizine Hydrochloride Tablets USP, 10 mg  
1000s count bottle label

NDC 25000-220-14

Compare to the active ingredient in Zyrtec® Tablets†

Original Prescription Strength

Allergy

# Cetirizine Hydrochloride Tablets USP, 10 mg

Antihistamine  
Indoor & Outdoor  
Allergies

24  
hour

Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

THIS PACKAGE FOR HOUSEHOLDS  
WITHOUT YOUNG CHILDREN



LOT :

EXP :

R38/23



M Marksans

1000 Tablets

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE.**

### Drug Facts

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 10 mg.....	Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives. ▶

### Drug Facts (Continued)

#### When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

#### If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

<b>Adults and children 6 years and over</b>	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
---	--

<b>Adults 65 years and over</b>	ask a doctor ▶
---------------------------------	----------------

Drug Facts (Continued)

Drug Facts (Continued)

<b>Children under 6 years of age</b>	ask a doctor	Manufactured for: Time-Cap Labs, Inc. 7 Michael Avenue, Farmingdale, NY 11735, USA  Manufactured by: Marksans Pharma Ltd. Plot No. L-82, L-83, Verna Indl. Estate, Verna, Goa - 403722, India.  Mfg. Lic. No. : GO/DRUGS/515
<b>Consumers with liver or kidney disease</b>	ask a doctor	
<b>Other information</b> ■ store between 20° to 25°C (68° to 77°F) ■ <b>Do not use if foil seal under cap is broken or missing.</b>		
<b>Inactive ingredients</b> lactose monohydrate, corn starch, povidone, magnesium stearate, hypromellose, polyethylene glycol, titanium dioxide		
<b>Questions?</b> Call at 1-877-376-4271		
†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.		

NDC 25000-220-78  
 Cetirizine Hydrochloride Tablets USP, 10 mg  
 7s count carton label

Allergy

# Cetirizine Hydrochloride Tablets USP, 10 mg

NDC 25000-220-78

Compare to the active ingredient in Zyrtec® Tablets†

Original Prescription Strength

Allergy

# Cetirizine Hydrochloride Tablets USP, 10 mg

Antihistamine  
Indoor & Outdoor  
Allergies

24  
hour

Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

PACKAGE NOT  
CHILD-RESISTANT



Actual Size

7 Tablets

M Marksans

Varnish/Print Omit area

Lot #: Exp. Date:

† This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division owner of Zyrtec® Tablets.

3 25000122078 5



Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.  
Mfg. Lic. No. : GO/DRUGS/515

R28/23

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE**  
**REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Cetirizine HCl 10 mg.....Antihistamine

**Uses**  
Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
■ runny nose  
■ sneezing  
■ itchy, watery eyes  
■ itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine.  
**Ask a doctor before use if you have** liver or kidney disease.  
Your doctor should determine if you need a different dose.  
**Ask a doctor or pharmacist before use if you are taking** tranquilizers or sedatives.  
**When using this product**  
■ drowsiness may occur ■ avoid alcoholic drinks  
■ alcohol, sedatives and tranquilizers may increase drowsiness  
■ be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.  
If pregnant or breast-feeding:


**Directions**  
**Adults and children 6 years and over**  
one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours.  
**Children 6 years and under**  
ask a doctor  
**Adults 65 years and over**  
ask a doctor  
**Children under 6 years of age**  
ask a doctor  
**Consumers with liver or kidney disease**  
ask a doctor

**Other information**  
■ Store between 20° to 25°C (68° to 77°F)  
■ Do not use if blister unit is torn or broken.

**Inactive ingredients** corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

**Drug Facts (Continued)**  
■ if breast-feeding: not recommended  
■ if pregnant: ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



Varnish/Print Omit area

C61 1 7R

NDC 25000-220-81

Cetirizine Hydrochloride Tablets USP, 10 mg

14s count carton label

# Cetirizine Hydrochloride Tablets USP, 10 mg

Allergy



**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE.**  
**REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**  
**Active ingredient (in each tablet)** Purpose  
 Cetirizine HCl 10 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ runny nose ■ sneezing  
 ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine.  
**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.  
**Ask a doctor or pharmacist before use if you are taking** tranquilizers or sedatives.  
**When using this product**  
 ■ drowsiness may occur ■ avoid alcoholic drinks ■ drowsiness and tranquilizers may increase  
 ■ alcohol, sedatives and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery.  
**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

**Directions**  
**Adults and children 6 years and over**  
 one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.  
**Adults 65 years and over**  
 ask a doctor  
**Children under 6 years of age**  
 ask a doctor  
**Consumers with liver or kidney disease**  
 ask a doctor

**Other information**  
 ■ Store between 20° to 25°C (68° to 77°F) ■ Do not use if blister unit is torn or broken.

**Inactive ingredients** corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

**Drug Facts (Continued)**  
 If pregnant or breast-feeding: not recommended  
 ■ if breast-feeding: not recommended  
 ■ if pregnant: ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Manufactured for:**  
 Time-Cap Labs, Inc.  
 7 Michael Avenue, Farmingdale,  
 NY 11735, USA

**Manufactured by:**  
 Marksans Pharma Ltd.  
 Plot No. L-82, L-83, Verna Indl. Estate,  
 Verna, Goa - 403722, India.  
 Mfg. Lic. No. : GO/DRUGS/515

**Lot No.:**  
**Exp Date:**

**Varnish Omit Area**

Manufactured for:  
 Time-Cap Labs, Inc.  
 7 Michael Avenue, Farmingdale,  
 NY 11735, USA

Manufactured by:  
 Marksans Pharma Ltd.  
 Plot No. L-82, L-83, Verna Indl. Estate,  
 Verna, Goa - 403722, India.  
 Mfg. Lic. No. : GO/DRUGS/515

R18/23

NDC 25000-220-81

Compare to the active ingredient in Zyrtec® Tablets†

Original Prescription Strength  
**Allergy**

# Cetirizine Hydrochloride Tablets USP, 10 mg

**Antihistamine**  
**Indoor & Outdoor**  
**Allergies**

**24**  
 hour

- Relief of**
- Sneezing
  - Runny Nose
  - Itchy, Watery Eyes
  - Itchy Throat or Nose

**PACKAGE NOT CHILD-RESISTANT**



**14 Tablets**




NDC 25000-160-81

Cetirizine Hydrochloride Tablets USP, 10 mg (Oval Shape)

14s count carton label

# Allegry Cetirizine Hydrochloride Tablets USP, 10 mg



**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE. REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**  
Cetirizine HCl 10 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ runny nose  
 ■ sneezing  
 ■ itchy, watery eyes  
 ■ itching of the nose or throat

**Warnings**  
**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine.  
**Ask a doctor before use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.  
**Ask a doctor or pharmacist before use if you are taking** tranquilizers or sedatives.  
**When using this product**  
 ■ drowsiness may occur ■ avoid alcoholic drinks  
 ■ alcohol, sedatives and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery.  
**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.  
 ■ if pregnant or breast-feeding: not recommended

**Drug Facts (Continued)**  
 ■ if pregnant: ask a health professional before use, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
**Adults and children 6 years and over**  
 one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.  
**Adults 65 years and over**  
 ask a doctor  
**Children under 6 years of age**  
 ask a doctor  
**Consumers with liver or kidney disease**  
 ask a doctor

**Other information**  
 ■ store between 20° to 25°C (68° to 77°F)  
 ■ Do not use if blister unit is torn or broken.

**Inactive ingredients**  
 corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division owner of Zyrtec® Tablets.

LOT:  
EXP:

Varnish Omit Area

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83, Verna Indl. Estate,  
Verna, Goa - 403722, India.  
Mfg. Lic. No. : GO/DRUGS/515

R05/25

NDC 25000-160-81

Compare to the active ingredient in Zyrtec® Tablets†

Original Prescription Strength  
Allergy

# Cetirizine Hydrochloride Tablets USP, 10 mg


Antihistamine  
Indoor & Outdoor  
Allergies

**24** hour Relief of  
 • Sneezing  
 • Runny Nose  
 • Itchy, Watery Eyes  
 • Itchy Throat or Nose




Varnish Omit Area

Varnish Omit Area



NDC 25000-160-03  
 Cetirizine Hydrochloride Tablets USP, 10 mg (Oval Shape)  
 30s count carton label



varnish  
omit area



**Drug Facts** (continued)

- be careful when driving a motor vehicle or operating machinery.

**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

<b>Adults and children 6 years and over</b>	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
<b>Adults 65 years and over</b>	ask a doctor
<b>Children under 6 years of age</b>	ask a doctor
<b>Consumers with liver or kidney disease</b>	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)

**Drug Facts** (continued)

- Do not use if foil seal under cap is broken or missing.**

**Inactive ingredients** corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec® Tablets.

NDC 25000-160-03

Compare to the active ingredient in Zyrtec® Tablets

**Original Prescription Strength Allergy**

**Cetirizine Hydrochloride Tablets USP, 10 mg**

**24** Relief of Sneezing, Runny Nose, Itchy, Watery Eyes, Itchy Throat or Nose

**Antihistamine Indoor & Outdoor Allergies**

**30** Tablets

**Marksans**

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE. REFER CARTON FOR COMPLETE INFORMATION**

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Cetirizine HCl 10 mg	Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine.

**Ask a doctor before use** if you have liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use** if you are taking tranquilizers or sedatives.

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness

Manufactured for:  
 Time-Cap Labs, Inc.  
 7 Michael Avenue, Farmingdale,  
 NY 11735, USA

Manufactured by:  
 Marksans Pharma Ltd.  
 Plot No. L-82, L-83,  
 Verma Indl. Estate,  
 Verma, Goa - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

RO/25

NDC 25000-160-03 (Oval Shape)  
 Cetirizine Hydrochloride Tablets USP, 10 mg

30s count bottle label

<p>NDC 25000-160-03</p> <p><b>Original Prescription Strength</b>  <b>Allergy</b></p> <p><b>Cetirizine Hydrochloride Tablets USP, 10 mg</b></p> <p><b>Antihistamine</b>  <b>Indoor &amp; Outdoor</b>  <b>Allergies</b></p> <p><b>24 hour</b> Relief of</p> <ul style="list-style-type: none"> <li>• Sneezing</li> <li>• Runny Nose</li> <li>• Itchy, Watery Eyes</li> <li>• Itchy Throat or Nose</li> </ul> <p><b>30 Tablets</b></p> <p>Marksans</p>	<p><b>Drug Facts</b></p> <table border="1"> <tr> <td><b>Active ingredient (in each tablet)</b></td> <td><b>Purpose</b></td> </tr> <tr> <td>Cetirizine HCl 10 mg.....</td> <td>Antihistamine</td> </tr> </table> <p><b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat</p> <p><b>Warnings</b>  <b>Do not use</b> if you have ever had an allergic reaction to this product ▶</p>	<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>	Cetirizine HCl 10 mg.....	Antihistamine	<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>Varnish Omit Area</b></p> <p>LOT :          EXP :          R05/25</p>
<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>					
Cetirizine HCl 10 mg.....	Antihistamine					

<p>or any of its ingredients, or to an antihistamine containing hydroxyzine. <b>Ask a doctor before use if you have</b> liver or kidney disease. Your doctor should determine if you need a different dose. <b>Ask a doctor or pharmacist before use if you are</b> taking tranquilizers or sedatives. <b>When using this product</b> ■ drowsiness may occur ■ avoid alcoholic drinks ▶</p>	<p>■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery. <b>Stop use and ask a doctor if</b> an allergic reaction to this product occurs. Seek medical help right away. <b>If pregnant or breast-feeding:</b> ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use. <b>Keep out of reach</b> ▶</p>	<p><b>of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)  <b>Directions Adults and children 6 years and over:</b> one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms. <b>Adults 65 years and over:</b> ask a doctor. <b>Children</b> ▶</p>
---	--	---

<p><b>under 6 years of age:</b> ask a doctor. <b>Consumers with liver or kidney disease:</b> ask a doctor.  <b>Other information</b> ■ store between 20° to 25°C (68° to 77°F) ■ <b>Do not use if foil seal under cap is broken or missing.</b>  <b>Inactive ingredients</b> corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, ▶</p>	<p>povidone, titanium dioxide  <b>Questions or comments?</b>          Call <b>1-877-376-4271</b>          (weekdays 9 AM to 5 PM)          Manufactured for:          Time-Cap Labs, Inc.          7 Michael Avenue, Farmingdale,          NY 11735, USA ▶</p>	<p>Manufactured by:          Marksans Pharma Ltd.          Plot No. L-82, L-83, Verna Indl. Estate,          Verna, Goa - 403722, India.          Mfg. Lic. No. : GO/DRUGS/515</p>
---	--	--

NDC 25000-160-14  
 Cetirizine Hydrochloride Tablets USP, 10 mg (Oval Shape)  
 1000s count bottle label

NDC 25000-160-14 Compare to the active ingredient in Zyrtec® Tablets†

**Original Prescription Strength**  
**Allergy**

**Cetirizine Hydrochloride Tablets USP, 10 mg**

**Antihistamine**  
**Indoor & Outdoor**  
**Allergies**

**24 hour** Relief of

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN



**1000 Tablets**

Varnish Omit Area

LOT: EXP: R05/25





IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USE.	
<b>Drug Facts</b>	
<b>Active ingredient (in each tablet)</b> Cetirizine HCl 10 mg.....	<b>Purpose</b> Antihistamine
<b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itchy, watery eyes</li> <li>■ itching of the nose or throat</li> </ul>	
<b>Warnings</b> <b>Do not use</b> if you have ever had an allergic reaction to this product or any of its ingredients, or to an antihistamine containing hydroxyzine. <b>Ask a doctor before use if you have</b> liver or kidney disease. Your doctor should determine if you need a different dose. <b>Ask a doctor or pharmacist before use if you are</b> taking tranquilizers or	
<b>Drug Facts</b> (continued)	
<b>When using this product</b> <ul style="list-style-type: none"> <li>■ drowsiness may occur</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery.</li> </ul>	
<b>Stop use and ask a doctor if</b> an allergic reaction to this product occurs. Seek medical help right away.	
<b>If pregnant or breast-feeding</b> <ul style="list-style-type: none"> <li>■ if breast-feeding: not recommended</li> <li>■ if pregnant: ask a health professional before use.</li> </ul> <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
<b>Directions</b>	
<b>Adults and children 6 years and over</b>	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
<b>Adults 65 years</b>	ask a doctor

sedatives.

and over

**Drug Facts** (continued)

<b>Children under 6 years of age</b>	ask a doctor
<b>Consumers with liver or kidney disease</b>	ask a doctor

**Other information**

- store between 20° to 25°C (68° to 77°F)
- **Do not use if foil seal under cap is broken or missing.**

**Inactive ingredients** corn starch, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, povidone, titanium dioxide

**Questions or comments?** Call 1-877-376-4271 (weekdays 9 AM to 5 PM)

<sup>†</sup>This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, distributor of Zyrtec<sup>®</sup> Tablets.

**Drug Facts** (continued)

Manufactured for:  
Time-Cap Labs, Inc.  
7 Michael Avenue, Farmingdale,  
NY 11735, USA

Manufactured by:  
Marksans Pharma Ltd.  
Plot No. L-82, L-83,  
Verna Indl. Estate,  
Verna, Goa - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

## CETIRIZINE HYDROCHLORIDE TABLETS, 5 MG

cetirizine hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:25000-219
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>STARCH, CORN</b> (UNII: O8232NY35J)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE (white to off white)	<b>Score</b>	no score
<b>Shape</b>	RECTANGLE (Rounded-off rectangular shaped)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	J219
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-219-03	1 in 1 CARTON	01/13/2022	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:25000-219-08	1 in 1 CARTON	01/13/2022	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-219-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/13/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078933	01/13/2022	

## CETIRIZINE HYDROCHLORIDE TABLETS, 10 MG

cetirizine hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:25000-220
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE (white to off white)	<b>Score</b>	no score
<b>Shape</b>	RECTANGLE (Rounded-off rectangular shaped)	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	J220
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-220-03	1 in 1 CARTON	01/13/2022	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:25000-220-08	1 in 1 CARTON	01/13/2022	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-220-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/13/2022	
4	NDC:25000-220-78	1 in 1 CARTON	04/17/2023	
4		7 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:25000-220-81	1 in 1 CARTON	07/21/2023	
5		14 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078933	01/13/2022	

## CETIRIZINE HYDROCHLORIDE TABLETS, 10 MG (OVAL SHAPE)

cetirizine hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:25000-160
---------------------	----------------	---------------------------	---------------

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETIRIZINE HYDROCHLORIDE (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	10mm
Flavor		Imprint Code	160
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-160-81	1 in 1 CARTON	04/10/2025	
1		14 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:25000-160-03	1 in 1 CARTON	04/10/2025	
2		30 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:25000-160-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/10/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078933	04/10/2025	

Labeler - MARKSANS PHARMA LIMITED (925822975)

## Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-160, 25000-219, 25000-220)

Revised: 7/2025

MARKSANS PHARMA LIMITED