

IBUPROFEN- ibuprofen tablet, coated
Velocity Pharma

Ibuprofen Tablets, USP 200mg

Active Ingredient

(in each tablet)

Ibuprofen 200 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- minor pains of arthritis
- muscular aches
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: This product may cause a severe allergic reaction, especially in people allergic to aspirin, symptoms may include

- hives
- facial swelling
- asthma(wheezing)
- shock
- skin reddning
- rash
- blisters

if an allergic reaction occurs, stop use and seek medical help right away

Stomach bleeding warning: this product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning(anticoagulant) or steroid drug.
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

do not use

- this product if you have ever had an allergic reaction to any pain reliever/fever reducer
- right before or after heart surgery

ask your doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma

ask your doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

when using this product

- take with food or milk
- The risk of heart attack or stroke may increase if you are more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. it is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persists.
- If pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other Information

- read all warnings and directions before use. keep carton
- store at 20-25 °C (68-77 °F)
- avoid excessive heat above 40°C (104°F)


- do not use if imprinted safety seal under the cap is broken or missing

Inactive Ingredients

Colloidal silicon Dioxide, corn starch, Croscarmellose Sodium, hypermellose, Iron Oxide red, lactose monohydrate, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Polyvinyl Alcohol, povidone, Pregelatinized Starch, stearic acid, Talc, Titanium Dioxide, triacetin

Questions or Comments

Call toll free 1-855-314-1850



7.2813 in

1.8750

3.5313

1.8750

Made in India

Distributed by
Vedochy Pharma LLC
Farmingdale, NY 11735

3 76168 01220 0

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- stop use and ask a doctor if
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- the smallest effective dose should be used

DOSE TABLE	
Age	Dose
Adults and children 12 years of age and over	Take 1 tablet every 4 to 6 hours while symptoms persist. Do not exceed 4 tablets in 24 hours. Always directed by a doctor.
Children under 12 years of age	Ask a doctor

Other information

- Each Tablet contains: Calcium 0.01mg, Magnesium 0.24mg, Sodium 0.71mg
- Read all warnings and directions before use.
- Store at 20 - 25°C (68 - 77°F)
- Avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, polyethylene glycol, polyvinylidene chloride, sodium starch glycolate, stearic acid, titanium dioxide, xanthan gum

Questions or comments?

Call 1-855-314-1850

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

*This product is an authorized or licensed by Pfizer Consumer Health Care, under the registered trademark ARIFF Tablet.

Ibuprofen

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

200 Tablets

Pain • Body Aches • Fever Reducer

Drug Facts

Active ingredient Purpose

Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, rash, facial swelling, skin redness or blisters, asthma (wheezing). If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

03/19-01

Ibuprofen

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

200 Tablets

Pain • Body Aches • Fever Reducer

Drug Facts

Active ingredient Purpose

Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, rash, facial swelling, skin redness or blisters, asthma (wheezing). If an allergic reaction occurs, stop use and seek medical help right away.


Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.



IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76168-095
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 1500 (UNII: 1212Z7S33A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	brown	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	Ibu200
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76168-095-24	24 in 1 BOTTLE	03/27/2019	
1		1 in 1 CARTON; Type 0: Not a Combination Product		
2	NDC:76168-095-30	30 in 1 BOTTLE	03/27/2019	
2		1 in 1 CARTON; Type 0: Not a Combination Product		
3	NDC:76168-095-20	200 in 1 BOTTLE	03/27/2019	
3		1 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	03/27/2019	

Labeler - Velocity Pharma (962198409)