

**STOOL SOFTENER- docusate sodium capsule, liquid filled**  
**Bi-Mart**

-----

**ACTIVE INGREDIENT (IN EACH SOFTGEL)**

Docusate sodium 100 mg

**PURPOSE**

Stool softener laxative

**USES**

- for the prevention of dry, hard stools
- for relief of occasional constipation
- this product generally produces a bowel movement within 12 to 72 hours

**WARNINGS**

**Do not use** if you are presently taking mineral oil, unless directed by a doctor.

**Ask a doctor before use if you have**

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over 2 weeks

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**DIRECTIONS**

- **adults and children 12 years of age and over:** take 1 softgel daily or as directed by a doctor
- **children under 12 years of age:** ask a doctor

**OTHER INFORMATION**

- **each softgel contains:** sodium 7 mg
- store at 15°- 25°C (59°- 77°F)

INACTIVE INGREDIENTS

Edible ink, FD&C red #40, FD&C yellow #6, gelatin (bovine), glycerin, polyethylene glycol, purified water, sorbitol special.

QUESTIONS?

call toll free 1-844-912-4012

Distributed by:

BI-MART

Eugene, OR 97402

BI-MART

NDC 37835-223-01

STOOL  
SOFTENER

DOCUSATE SODIUM 250 mg

100 SOFTGELS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active ingredient (in each softgel)**  
Docusate sodium 250 mg.....Stool softener laxative

**Uses**  
■ for the prevention of dry, hard stools  
■ for relief of occasional constipation  
■ this product generally produces a bowel movement within 12 to 72 hours

**Warnings**  
Do not use if you are presently taking mineral oil, unless directed by a doctor.  
**Ask a doctor before use if you have**  
■ stomach pain ■ nausea ■ vomiting  
■ noticed a sudden change in bowel habits that lasts over 2 weeks

**Stop use and ask a doctor if** ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.  
■ you need to use a laxative for more than 1 week

**if pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
■ adults and children 12 years of age and over: take 1 softgel daily or as directed by a doctor  
■ children under 12 years of age: ask a doctor

**Other information**  
■ each softgel contains: sodium 7 mg  
■ store at 15° - 25°C (59° - 77°F)

**Inactive ingredients**  
Edible ink, FD&C red #40, FD&C yellow #6, gelatin (bovine), glycerin, polyethylene glycol, purified water, sorbitol special.

**Questions?** call toll-free 1-844-912-4012

Distributed by: BI-MART, Eugene, OR 97402 223-01-BM 02/25

BI-MART UNUSUAL MONEY BACK GUARANTEE  
If you are not completely satisfied with your BI-MART purchase, return it to the store for a full refund. No questions asked. The unused portion and your entire purchase price will be covered. No restocking fee.

0 71357 00773 8

STOOL SOFTENER

docusate sodium capsule, liquid filled

| Product Information   |                |                    |               |
|---|----------------|--------------------|---------------|
| Product Type  | HUMAN OTC DRUG | Item Code (Source) | NDC:37835-223 |
| Route of Administration   | ORAL           |                    |               |
|   |                |                    |               |
| Active Ingredient/Active Moiety                                 |                |                    |               |
| Ingredient Name   |                | Basis of Strength  | Strength      |
| DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG) |                | DOCUSATE SODIUM    | 250 mg        |
|   |                |                    |               |
| Inactive Ingredients  |                |                    |               |
| Ingredient Name   |                |                    | Strength      |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8)                            |                |                    |               |
| FD&C RED NO. 40 (UNII: WZB9127XOA)                              |                |                    |               |

|   |  |
|---|--|
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                          |  |
| <b>GELATIN TYPE B BOVINE (160 BLOOM)</b> (UNII: 1T8387508X) |  |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)  |  |
| <b>WATER</b> (UNII: 059QF0KO0R)                             |  |
| <b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)                 |  |

### Product Characteristics

|                 |               |                     |          |
|-----------------|---------------|---------------------|----------|
| <b>Color</b>    | red           | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL (oblong) | <b>Size</b>         | 20mm     |
| <b>Flavor</b>   |               | <b>Imprint Code</b> | SCU1     |
| <b>Contains</b> |               |                     |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:37835-223-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 03/01/2025           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007                                     | 03/01/2025           |                    |

**Labeler -** Bi-Mart (027630078)