

IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE- ibuprofen and pseudoephedrine hydrochloride tablet, sugar coated
Ohm Laboratories Inc.

Ibuprofen and Pseudoephedrine hydrochloride

Drug Facts

Active ingredients (in each Purposes caplet)

Ibuprofen, USP 200 mg (NSAID) *	Pain reliever/fever reducer
Pseudoephedrine HCl, USP 30 mg	Nasal decongestant

* nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms associated with the common cold or flu:

- headache
- fever
- sinus pressure
- nasal congestion
- minor body aches and pains

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- symptoms continue or get worse
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years of age and over:
 - take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used.
 - do not use more than 6 caplets in any 24-hour period unless directed by a doctor
- children under 12 years of age: do not use

Other information

- store at 20 - 25° C (68 - 77° F). Avoid excessive heat above 40° C (104° F).
- read all warnings and directions before use. Keep carton.

Inactive ingredients

acacia, calcium carbonate, carnauba wax, confectioner's sugar, corn starch, croscarmellose sodium, crospovidone, FD&C Blue no. 2 Aluminum Lake, FD&C Red no. 40 Aluminum Lake, FD&C Yellow no. 6 Aluminum Lake, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, iron oxide black, kaolin, polyethylene glycol, powdered cellulose, povidone, pregelatinized starch, propylene glycol, shellac, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

Questions?

call **1-800-406-7984**

Distributed by: Ohm Laboratories Inc.
North Brunswick, NJ 08902

PRINCIPAL DISPLAY PANEL - 20 Caplet Blister Pack Carton

**†Compare To
the active ingredients of
Advil[®] Cold & Sinus**

NDC 51660-423-21

See New Warnings Information

ohm[®]

Non-Drowsy

Cold & Sinus

Ibuprofen and Pseudoephedrine

HCl Tablets, USP

Ibuprofen, USP 200 mg - Pain Reliever/Fever Reducer (NSAID)*

Pseudoephedrine HCl, USP 30 mg - Nasal Decongestant

*nonsteroidal anti-inflammatory drug

- ***Relieves Sinus Pressure,
Nasal Congestion and Fever***

20 Coated Caplets**

(oval-shaped tablets)**

Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives
facial swelling
asthma (wheezing)
shock
stomach bleeding
rash
diarrhea
if an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer
if you have had an allergic reaction to any other pain reliever/fever reducer
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if:
stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers or fever reducers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
under a doctor's care for any serious condition
taking any other product that contains pseudoephedrine or any other nasal decongestant

Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives
facial swelling
asthma (wheezing)
shock
stomach bleeding
rash
diarrhea
if an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer
if you have had an allergic reaction to any other pain reliever/fever reducer
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if:
stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers or fever reducers
you have a history of stomach problems, such as heartburn
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, have trouble urinating due to an enlarged prostate gland, or had a stroke
you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
under a doctor's care for any serious condition
taking any other product that contains pseudoephedrine or any other nasal decongestant

Drug Facts (continued)

Active Ingredients (in each caplet)
Ibuprofen, USP 200 mg (NSAID)
Pseudoephedrine HCl, USP 30 mg
Pain reliever/fever reducer
nasal decongestant

Purposes
Ibuprofen, USP 200 mg (NSAID)
Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Drug Facts (continued)

Uses temporarily relieves these symptoms associated with the common cold or flu: headache
fever
sinus pressure
nasal congestion
minor body aches and pains

Drug Facts (continued)

Directions
Do not take more than directed
the smallest effective dose should be used
adults and children 12 years of age and over:
take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. Do not use more than 6 caplets in any 24-hour period unless directed by a doctor
children under 12 years of age: do not use

Other Information
Store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).
Read all warnings and directions before use. Keep caplets in their original container.

ohm
LABORATORIES INC.
a SUN PHARMA company

Compare To
the active ingredients of
Advil® Cold & Sinus

NDC 51660-423-21

See New
Warnings Information

Expiration Date: _____

Batch No. _____

Non-Drowsy

Cold & Sinus

Ibuprofen and Pseudoephedrine HCl Tablets, USP

Ibuprofen, USP 200 mg – Pain Reliever/Fever Reducer (NSAID)*
Pseudoephedrine HCl, USP 30 mg – Nasal Decongestant

*nonsteroidal anti-inflammatory drug

- **Relieves Sinus Pressure, Nasal Congestion and Fever**

20 Coated Caplets**
(**oval-shaped tablets)

Drug Facts (continued)

Inactive Ingredients acacia, calcium carbonate, camphor wax, croscarmellose sodium, croscopolone, FD&C Blue no. 2, Aluminum Lake, FD&C Red no. 40, Aluminum Lake, FD&C Yellow no. 6, Aluminum Lake, gelatin, guar gum, hydrogenated vegetable oil, hydroxypropyl cellulose, iron oxide black, lactin, polyethylene glycol, powdered cellulose, powderson, pregelatinized starch, propylene glycol, shellac, silicon dioxide, sodium benzoate, sucrose, talc, titanium dioxide, white wax

Questions? call 1-800-406-7984

Drug Facts (continued)

Directions
Do not take more than directed
the smallest effective dose should be used
adults and children 12 years of age and over:
take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. Do not use more than 6 caplets in any 24-hour period unless directed by a doctor
children under 12 years of age: do not use

Other Information
Store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).
Read all warnings and directions before use. Keep caplets in their original container.

Drug Facts (continued)

Directions
Do not take more than directed
the smallest effective dose should be used
adults and children 12 years of age and over:
take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. Do not use more than 6 caplets in any 24-hour period unless directed by a doctor
children under 12 years of age: do not use

Other Information
Store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).
Read all warnings and directions before use. Keep caplets in their original container.

Drug Facts (continued)

Directions
Do not take more than directed
the smallest effective dose should be used
adults and children 12 years of age and over:
take 1 caplet every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 caplet, 2 caplets may be used. Do not use more than 6 caplets in any 24-hour period unless directed by a doctor
children under 12 years of age: do not use

Other Information
Store at 20° - 25° C (68° - 77° F). Avoid excessive heat above 40° C (104° F).
Read all warnings and directions before use. Keep caplets in their original container.

IBUPROFEN AND PSEUDOEPHEDRINE HYDROCHLORIDE

ibuprofen and pseudoephedrine hydrochloride tablet, sugar coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51660-423
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
KAOLIN (UNII: 24H4NWX5CO)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
GUAR GUM (UNII: E89I1637KE)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL (caplets)	Size	14mm
Flavor		Imprint Code	423

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51660-423-21	20 in 1 BLISTER PACK; Type 0: Not a Combination Product	10/13/2001	
2	NDC:51660-423-41	40 in 1 BLISTER PACK; Type 0: Not a Combination Product	10/13/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074567	10/13/2001	

Labeler - Ohm Laboratories Inc. (184769029)**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)**Establishment**

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(51660-423)

Revised: 2/2025

Ohm Laboratories Inc.