GADAVYT - magnesium citrate liquid Gadal Laboratories Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredient Purpose

Magnesium citrate 1.745g per fl. Oz..... Saline Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet
- a sodium restricted diet
- · stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 1 week.

Ask a doctor or pharmacist before use if you are

• presently taking a prescription drug

When using this product

• do not exceed the maximum recommended daily dosage in a 24 hour period

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after use. These could be signs of a serious condition
- you need to use a laxative for more than 1 week

Keep this and all drugs out of the reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center (1-800-222-1222) immediately

If pregnant or breast-feeding ask a health professional before use.

Directions

	1/2 to 1 bottle (10 fl. oz.) Drink a full 8 oz glass of liquid with each dose. These dose may be taken as a single daily dose or in divided doses
children 6 to under 12 years of age	1/3 to 1/2 bottle with a fill 8 oz glass of liquid
children under 6 years of age	consult a doctor

Other information

• each fluid once contains: sodium 36 mg, magnesium 199 mg

- store at room temperature
- protect from excessive heat and freezing
- discard unused product within 24 hours of opening

Inactive ingredients

citric acid, flavors, purified water, red 40, blue #1, saccharin sodium, sodium bicarbonate


Questions or comments?

(305) 382-3288

Gadal Laboratories Inc

Miami Fl 33186





Drug Facts Active ingredient Magnesium citrate 1.745g po	Purpose er fl. OzSaline laxativ
Uses - relieves occasional constip - generally produces bowel r	
Warnings Ask a doctor before use if - kidney disease - a magnesium restricted die - a sodium restricted diet - stomach pain, nausea or v - noticed a sudden change ir more than 1 week	t omiting
Ask a doctor or pharmacist • presently taking a prescript	
When using this product do not exceed the maximum dosage in a 24 hour period	n recommended daily
Stop use and ask a doctor you have rectal bleeding or use. These could be signs you need to use a laxative	no bowel movement afte of a serious condition
If pregnant or breast-feedi professional before use.	ng, ask a health
Keep this and all drugs ou In case of accidental overdo assistance or contact a Pois (1-800-222-1222) immediate	se, seek professional on Control Center

GADAVYT

magnesium citrate liquid

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53113-102	
Route of Administration	ORAL			

I	Active Ingredient/Active Moiety				
I	Ingredient Name	Basis of Strength	Strength		
	MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL		

Inactive Ingredients			
Ingredient Name	Strength		
CITRIC ACID MONO HYDRATE (UNII: 2968 PHW8 QP)			
FD&C RED NO. 40 (UNII: WZB9 127XOA)			
SACCHARIN SO DIUM DIHYDRATE (UNII: SB8 ZUX40 TY)			
WATER (UNII: 059QF0KO0R)			

SODIUM BICARBONATE (UNII: 8 MDF5 V39 QO)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	GRAPE (Grape)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53113-102-10	296 mL in 1 BOTTLE, GLASS		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part334	10/01/2012	

Labeler - Gadal Laboratories Inc (841305639)

Revised: 10/2012 Gadal Laboratories Inc