FIRST AID MAXIMUM STRENGTH BACITRAYCIN PLUS- bacitracin, and pramoxine hcl ointment

First Aid Research Corp

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

First Aid Maximum Strength Bacitraycin Plus Ointment 1oz 49555 (2018)

Active ingredients (each gram contains) Purpose

Bacitracin 500 units......First Aid Antibiotic

Uses

first aid to help prevent infection in

- minor cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- in the eyes
- if you are allergic to any of the ingredients
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use in case of deep or puncture wounds, animal bites, or serious burns

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Other information

- store at 15° to 25°C (59° to 77°F)
- lot no. & exp. date: see box or see crimp of tube

Inactive ingredients

aloe vera leaf, methylparaben, mineral oil, petrolatum, propylparaben, stearyl alcohol

Distributed by:

First Aid Research Corp.

Jupiter, FL 33478

Made in Korea











Distributed by: FIRST AID RÉSEARCH CORP. Jupiter, FL 33478

Made in Korea

Purpose



Inactive ingredients aloe vera leaf, methylparaben, mineral oil, petrolatum, propylparaben, stearyl alcohol

■ store at 15° to 25° C (59° to 77° F)
■ lot no. & exp. date: see box or see crimp of tube

Other information

Directions ■ clean the affected area
■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area
1 to 3 times daily
■ may be covered with a sterile bandage

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Stop use and ask a doctor if

the condition persists or gets worse

a stash or other allergic reaction develops Ask a doctor before use in case of deep or puncture wounds, animal bites, or serious burns.

■ over large areas of the body

longer than 1 week unless directed by a doctor

■ in the eyes ■ if you are allergic to any of the ingredients

esu ton od Warnings For external use only.

suunq ■ scusbes ■ minor cuts Uses first aid to help prevent infection in

First Aid Antibiotic

Bacitracin 500 units... Pramoxine HCl 10 mg..

Active ingredients (each gram contains)

Drug Facts

with Pain Reliever First Aid Ointment





LOT & EXP



FIRST AID MAXIMUM STRENGTH BACITRAYCIN PLUS

bacitracin, and pramoxine hclointment

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:75983-055	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
BACITRACIN (UNII: 58 H6 RWO52I) (BACITRACIN - UNII:58 H6 RWO52I)	BACITRACIN	500 [USP'U] in 1 g		
PRAMO XINE HYDRO CHLO RIDE (UNII: 88 AYB867L5) (PRAMO XINE - UNII: 068 X84E056)	PRAMO XINE HYDRO CHLO RIDE	10 mg in 1 g		

Inactive Ingredients	
Ingredient Name	Strength
METHYLPARABEN (UNII: A2I8C7HI9T)	
ALOE VERA LEAF (UNII: ZY8 1Z8 3H0 X)	
MINERAL OIL (UNII: T5L8T28FGP)	
PETROLATUM (UNII: 4T6H12BN9U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:75983-055-28	1 in 1 CARTON	0 3/0 1/20 18		
1		28 g in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part333B	0 3/0 1/20 18		

Labeler - First Aid Research Corp (089405927)

Revised: 3/2018 First Aid Research Corp