

**UPSET STOMACH RELIEF-BISMUTH SUBSALICYLATE 262 MG- bismuth subsalicylate 262 mg capsule, liquid filled**  
**NUVICARE LLC**

-----  
**Nuvicare-Upset Stomach Relief-Bismuth Subsalsicylate 262 mg**

**Active Ingredient (in each softgel)**

Bismuth subsalicylate 262 mg

**Purpose**

Upset stomach reliever and antidiarrheal

**Uses**

relieves:

■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**If pregnant or breast feeding,** ask a health professional before use.

**KEEP OUT OF REACH OF CHILDREN** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
  - 2 softgels every ½ hour or 4 softgels every hour as needed for diarrhea
  - 2 softgels every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 softgels) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- each softgel contains:
  - salicylate 99 mg
  - low sodium
- do not exceed 25°C

### **Inactive ingredients**

Polyethylene Glycol-400, Polysorbate 80, Soya Lecithin, Povidone (PVPK 30), Gelatin, Glycerin, Sorbitol Sorbitan Solution, D & C Red 33 , Titanium Dioxide, Purified Water

### **Questions or Comments?**

Call 1 (718) 337-8733 or email: [support@nuvicare.com](mailto:support@nuvicare.com)



<b>Gelatin</b> (UNII: 2G86QN327L)				
<b>Glycerin</b> (UNII: PDC6A3C0OX)				
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)				
<b>SORBITAN</b> (UNII: 6O92ICV9RU)				
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)				
<b>Titanium Dioxide</b> (UNII: 15FIX9V2JP)				
<b>WATER</b> (UNII: 059QF0KO0R)				
<b>Product Characteristics</b>				
<b>Color</b>	pink (Opaque baby pink)		<b>Score</b>	no score
<b>Shape</b>	OVAL (Oblong)		<b>Size</b>	20mm
<b>Flavor</b>			<b>Imprint Code</b>	SD30
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84324-028-01	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/25/2025	
<b>Marketing Information</b>				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M008	06/25/2025	

**Labeler** - NUVICARE LLC (119257565)

Revised: 6/2025

NUVICARE LLC