

IBUPROFEN- ibuprofen tablet, film coated
Meijer Distribution Inc

Meijer Distribution, Inc. Ibuprofen Tablets, 200 mg Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> • take 1 caplet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 caplet, 2 caplets may be used • do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> • ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

VALUE 2 PACK

meijer®

Compare to Advil® active ingredient

ibuprofen Tablets, 200 mg

SEE REVISED WARNING

Pain Reliever | Fever Reducer (NSAID)

100 Caplets**

**capsule shaped tablets

actual size

A5102 0E C7



7 19283 31159 1 7

FID 861200

Drug Facts (continued)

do not be more than directed
 the smallest effective dose should be used
 children
 4 to 6 hours while
 take 1 caplet every
 4 to 6 hours as needed
 if pain or fever does
 not respond to 1 caplet,
 2 caplets may be used
 do not exceed 6 caplets
 in 24 hours unless
 directed by a doctor
 children under
 12 years
 ask a doctor

Drug Facts (continued)

Directions

do not be more than directed
 the smallest effective dose should be used
 children
 4 to 6 hours while
 take 1 caplet every
 4 to 6 hours as needed
 if pain or fever does
 not respond to 1 caplet,
 2 caplets may be used
 do not exceed 6 caplets
 in 24 hours unless
 directed by a doctor
 children under
 12 years
 ask a doctor

Questions or comments?
 1-800-719-9050

Drug Facts (continued)

Some headache relief may be obtained from this product
 contains an NSAID, which may cause severe
 stomach bleeding. The chance is higher if you
 are age 60 or older
 have had stomach ulcers or bleeding
 prod. bms
 take a blood thinning (anticoagulant) or
 steroid drug
 take other drug containing prescription or
 nonprescription NSAID (aspirin, ibuprofen,
 naproxen, or others)
 have 3 or more alcoholic drinks every day
 while using this product
 take more or for a longer time than directed
 Heart attack and stroke warning: NSAIDs,
 except aspirin, increase the risk of heart attack.
 The risk is higher if you use more than directed
 or for longer than directed
 Do not use
 if you have seen an allergic reaction to
 a product or any other pain reliever
 right before or after heart surgery
 Ask a doctor before use if
 you have problems or serious side effects
 from taking pain relievers or fever reducers
 the stomach bleeding warning applies to you
 you have a history of stomach problems,
 such as heartburn
 you have high blood pressure, heart disease,
 liver disease, kidney disease, asthma, or
 had a stroke
 you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if
 you are
 taking aspirin for heart attack or stroke,
 because it may increase the risk of
 aspirin
 under a doctor's care for an existing condition
 taking any other drug
 when using this product
 take with food or milk if stomach upset occurs
 Stop use and ask a doctor if
 you experience any of the following signs of
 stomach bleeding:
 vomit blood
 have bloody or black stools
 have stomach pain that does not get better
 you have symptoms of an existing condition
 or trouble breathing
 chest pain
 weak knees in one part or side of body
 blurred speech
 big swelling
 pain gets worse or lasts more than 10 days
 fever gets worse or lasts more than 3 days
 redness or swelling is present in the painful area
 any new symptoms appear
 if pregnant or breast-feeding, ask a health
 professional before use, especially
 if pregnant or breast-feeding at 2 weeks or
 later in pregnancy unless directed to
 do so by your doctor because of a specific
 problem in the unborn child or complications
 during delivery
 Keep out of reach of children. In case of
 overdose, get medical help or contact Poison
 Control Center right away (1-800-222-1222).

Drug Facts

Active ingredient
 (in each caplet)
 Ibuprofen 200 mg (NSAID) ...
 other inactive ingredients

Purposes
 temporarily reduces fever
 temporarily reduces pain
 temporarily reduces inflammation

Warnings
 Allergic reactions to ibuprofen may cause a severe
 allergic reaction, especially in people allergic to
 aspirin. Symptoms may include:
 hives
 facial swelling
 asthma (wheezing)
 shock
 skin redness
 rash
 blisters
 If an allergic reaction occurs, stop use and seek
 medical help right away.

Drug Facts (continued)

Warnings
 Allergic reactions to ibuprofen may cause a severe
 allergic reaction, especially in people allergic to
 aspirin. Symptoms may include:
 hives
 facial swelling
 asthma (wheezing)
 shock
 skin redness
 rash
 blisters
 If an allergic reaction occurs, stop use and seek
 medical help right away.

DO NOT USE IF PPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Important: Read all product information on using.
 Keep this box or insert in front of you.

By GlaxoSmithKline Consumer Healthcare Holdings
 (US) LLC, owner of the registered trademark Advil®.
 ***This product is not manufactured or distributed

GF
 GLUTEN FREE
 OUR QUALITY
 GUARANTEED

www.mw.us
 DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49504

www.mw.us
 1-800-368-7777

www.mw.us
 1-800-368-7777

Drug Facts (continued)

Warnings
 Allergic reactions to ibuprofen may cause a severe
 allergic reaction, especially in people allergic to
 aspirin. Symptoms may include:
 hives
 facial swelling
 asthma (wheezing)
 shock
 skin redness
 rash
 blisters
 If an allergic reaction occurs, stop use and seek
 medical help right away.

82-647-4129141 NDC

VALUE 2 PACK

meijer Compare to Advil® active ingredient***

ibuprofen Tablets, 200 mg
 Pain Reliever | Fever Reducer (NSAID)

SEE REVISED WARNING

100 Caplets**
 **capsule shaped tablets

 actual size

meijer Compare to Advil® active ingredient***



IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-647
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BROWN	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	I;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:41250-647-71	1 in 1 CARTON	11/15/1988	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-647-78	1 in 1 CARTON	09/20/1991	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-647-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/07/1994	
4	NDC:41250-647-82	2 in 1 CARTON	09/02/2016	
4		100 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:41250-647-76	1 in 1 CARTON	04/17/1992	06/26/2017
5		120 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	11/15/1988	

Labeler - Meijer Distribution Inc (006959555)

Revised: 9/2023

Meijer Distribution Inc