

ANTACID MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, simethicone suspension

Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Antacid Drug Facts

Active ingredients (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80

Purpose

Antacid

Antigas

Uses

for the relief of

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms
- pressure and bloating commonly referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

do not take more than 40 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician

Keep out of reach of children.

Directions

- shake well before using
- only use the dose cup provided
- adults and children 12 years and older: take 10 mL to 20 mL two times a day or as directed by a physician
- do not take more than 40 mL in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: consult a physician

Other information

- **each 10 mL contains:** magnesium 350 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- protect from freezing

Inactive ingredients

butylparaben, flavor, glycerin, hydroxyethyl cellulose, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol, sorbitol solution

Questions or comments?

1-800-719-9260

Principal Display Panel

ADVANCED MAXIMUM STRENGTH

antacid

Antacid | Anti-Gas

FAST RELIEF OF: Heartburn, Acid Indigestion, Pressure & Bloating

Cherry Flavor

ALCOHOL 0.1%

12 FL OZ (355 mL)

meijer

NDC 41250-588-40

ADVANCED
MAXIMUM STRENGTH

antacid

Antacid | Anti-Gas

FAST RELIEF OF: Heartburn, Acid
Indigestion, Pressure & Bloating

Cherry Flavor

ALCOHOL 0.1%

12 FL OZ (355 mL)

: 58840 6E F7



Drug Facts

Active ingredients (in each 10 mL)	Purpose
Aluminum hydroxide (equiv. to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg.....	Antigas

Uses for the relief of ■ acid indigestion ■ heartburn ■ sour stomach
■ upset stomach associated with these symptoms
■ pressure and bloating commonly referred to as gas

Warnings

Ask a doctor before use if you have

■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product do not take more than 40 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician

Keep out of reach of children.

Directions

■ shake well before using ■ only use the dose cup provided ■ adults and children 12 years and older: take 10 mL to 20 mL two times a day or as directed by a physician ■ do not take more than 40 mL in 24 hours or use the maximum dosage for more than 2 weeks ■ children under 12 years: consult a physician

Other information ■ each 10 mL contains: magnesium 350 mg and sodium 3 mg ■ does not meet USP requirements for preservative effectiveness ■ store at 20-25°C (68-77°F) ■ protect from freezing

Inactive ingredients butylparaben, flavor, glycerin, hydroxyethyl cellulose, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol, sorbitol solution ▶

Drug Facts
(continued)

Questions or
comments?
1-800-719-9260

DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



WWW.MEIJER.COM/
SATISFACTION

: 58840 6E B7

DO NOT USE IF
PRINTED NECKBAND
IS BROKEN OR
MISSING



PID 3103042



7

ANTACID MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-588
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLPARABEN (UNII: 3QP1I3FV8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-588-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/19/2005	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part332	10/19/2005	

