

**PEPTO-BISMOL- bismuth subsalicylate tablet**  
**Procter & Gamble Manufactura S de RL de CV**

-----  
**Pepto BISMOL® Caplets**

**Drug Facts**

**Active ingredient (in each caplet)**

Bismuth subsalicylate 262 mg

**Purpose**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

**Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if** you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- swallow with water, do not chew
- adults and children 12 years and over:
  - 2 caplets (1 dose) every ½ hour or 4 caplets (2 doses) every hour as needed for diarrhea
  - 2 caplets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information**

- **each caplet contains:** calcium 27 mg
- salicylate 99 mg
- low sodium
- avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients**

calcium carbonate, D&C red No. 27 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, povidone, silicon dioxide, sodium starch glycolate

**Questions?**

**1-800-717-3786**

**TAMPER EVIDENT: Do not use if inner printed seal on bottle is broken or missing.**

**Made in Mexico**

**DIST. BY PROCTER & GAMBLE,  
CINCINNATI OH 45202**

**PRINCIPAL DISPLAY PANEL - 24 Caplet Carton**

**Pepto  
Bismol®**

Bismuth Subsalicylate

Upset Stomach Reliever/

Antidiarrheal

**5 SYMPTOM RELIEF**

NAUSEA

HEARTBURN

INDIGESTION

UPSET STOMACH

DIARRHEA

**24 CAPLETS**



**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C RED NO. 27 ALUMINUM LAKE</b> (UNII: ZK64F7XSTX)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	OVAL (Caplet)	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	PeptoBismol
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-476-01	1 in 1 POUCH	06/30/2004	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:58933-476-02	1 in 1 CARTON	06/30/2004	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:58933-476-03	1 in 1 CARTON	06/30/2004	
3		40 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:58933-476-12	17852 in 1 BAG; Type 0: Not a Combination Product	06/30/2004	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	06/30/2004	

---

Revised: 10/2025

Procter & Gamble Manufactura S de RL de CV