

**DIMAPHEN DM- brompheniramine maleate, dextromethorphan hbr,  
phenylephrine hcl solution**  
**Major Pharmaceuticals**

-----  
**Major Pharmaceuticals Children's Dimaphen DM Drug Facts**

**Active ingredients (in each 10 mL)**

Brompheniramine maleate, USP 2 mg

Dextromethorphan HBr, USP 10 mg

Phenylephrine HCl, USP 5 mg

**Purposes**

Antihistamine

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

**Ask a doctor before use if you have**

- heart disease

- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers

**When using this product**

- **do not use more than directed**
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

## Other information

- **each 10 mL contains:** sodium 4 mg
- store at 20°-25°C (68°-77°F)

## Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

## Questions or comments?

**1-800-616-2471**

## Principal Display Panel

Children's

Dimaphen™ DM

ANTIHISTAMINE, Brompheniramine Maleate

COUGH SUPPRESSANT, Dextromethorphan HBr

NASAL DECONGESTANT, Phenylephrine HCl

Relieves – Nasal Symptoms

Stuffy Nose – Runny Nose

Sneezing

Plus Other Symptoms

Itchy, Watery Eyes

Cough

GRAPE FLAVOR

Alcohol-Free

6 yrs. and older

COMPARE TO the active ingredients of DIMETAPP® COLD & COUGH

4 FL. OZ. (118 mL.)



## DIMAPHEN DM

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0904-6463
<b>Route of Administration</b>	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>BROMPHENIRAMINE MALEATE</b> (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

Product Characteristics			
<b>Color</b>	PURPLE (Clear bluish-red)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6463-20	1 in 1 CARTON	09/11/2006	11/30/2026
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/11/2006	11/30/2026

**Labeler** - Major Pharmaceuticals (191427277)