CETIRIZINE HYDROCHLORIDE- cetirizine hydrochloride solution Taro Pharmaceuticals U.S.A., Inc.

Children's Cetirizine Hydrochloride Oral Solution

Drug Facts

Active ingredient (in each 5 mL)

Cetirizine HCl 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

(1-800-222-1222)

Directions

- use only with enclosed dosing cup
- find right dose on chart below

• mL = milliliter

adults and children 6 years and over	5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours.
adults 65 years and over	5 mL once daily; do not take more than 5 mL in 24 hours.
children 2 to under 6 years of age	2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every 12 hours. Do not give more than 5 mL in 24 hours.
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- do not use if carton is opened or if imprinted safety seal is broken or missing
- see bottom panel for lot number and expiration date
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

bubble gum artificial flavor, glacial acetic acid, glycerin, methylparaben, propylene glycol, propylparaben, purified water, sodium acetate anhydrous, sucralose

Questions?

Call 1-866-923-4914

Distributed by: **Taro Pharmaceuticals U.S.A., Inc.** Hawthorne, NY 10532

PRINCIPAL DISPLAY PANEL - 120 mL Bottle Carton

Compare to the active ingredient in Children's Zyrtec®*

NDC 51672-2106-8

Children's
Cetirizine
Hydrochloride
Oral Solution
1 mg/mL ALLERGY
Antihistamine

Indoor & Outdoor Allergies

SUGAR FREE

24 Hour

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

2 Yrs. & older

Bubble Gum

Flavor

Dosing Cup Included

4 FL OZ

(120 mL)



079

SEALED WITH PRINTED NECKBAND

Children's Cetirizine **Hydrochloride Oral Solution** 1 mg/mL ALLERGY

Antihistamine

Indoor & Outdoor Allergies

- Alcohol Free
- Dye Free Sugar Free



Dosing Cup Included

Dosing cup should be washed and left to air dry after each use.

*This product is not manufactured or distributed by UCB Pharma, S.A. CORPORATION BELGIUM, owner of the registered trademark Children's Zyrtece

stributed by: Taro Pharmaceuticals U.S.A., Inc. awthorne, NY 10532 Made in Israel

NDC 51672-2106-8

active ingredient in Children's Zyrtec8*

Children's Cetirizine **Hydrochloride Oral Solution**

1 mg/mL ALLERGY

Antihistamine Indoor & Outdoor Allergies

SUGAR FREE

24 Relief of: Sneezing

- Runny NoseItchy, Watery Eyes
- Itchy Throat or Nose

2 Yrs. & older



2-2110-77002

Drug Facts

Active ingredient Purpose (in each 5 mL) Cetirizine HCl 5 mg.....

.Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezina
- itchy, watery eyes
 itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may
- increase drowsiness be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- · if breast-feeding: not recommended if pregnant: ask a health professional
- before use.

 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)

Directions

- use only with enclosed dosing cup
 find right dose on chart below
 mL = milliliter

adults and children 6 years and over	5 mL or 10 mL once daily depending upon severity of symptoms; do not take more than 10 mL in 24 hours.		
adults 65 years and over	5 mL once daily; do not take more than 5 mL in 24 hours.		
children 2 to under 6 years of age	2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every 12 hours. Do not give more than 5 mL in 24 hours.		
children under 2 years of age	ask a doctor		
consumers with liver or kidney disease	ask a doctor		

Other information

- do not use if carton is opened or if imprinted safety seal is broken or missing see bottom panel for lot number and
- expiration date
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients
bubble gum artificial flavor, glacial acetic acid, glycerin, methylparaben, propylene glycol, propylparaben, purified water, sodium acetate anhydrous, sucralose

Questions? Call 1-866-923-4914



CETIRIZINE HYDROCHLORIDE

cetirizine hydrochloride solution

Product Information

Product Type HUMAN OTC DRUG Item Code (Source)

Route of Administration ORAL NDC:51672-2106

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Cetirizine Hydrochloride (UNII: 64O047KTOA) (Cetirizine - UNII:YO7261ME24)	Cetirizine Hydrochloride	1 mg in 1 mL	

Inactive Ingredients			
Ingredient Name	Strength		
acetic acid (UNII: Q40Q9N063P)			
glycerin (UNII: PDC6A3C0OX)			
methylparaben (UNII: A2I8C7HI9T)			
propylene glycol (UNII: 6DC9Q167V3)			
propylparaben (UNII: Z8IX2SC1OH)			
water (UNII: 059QF0KO0R)			
sodium acetate anhydrous (UNII: NVG71ZZ7P0)			
sucralose (UNII: 96K6UQ3ZD4)			

Product Characteristics				
Color	YELLOW (colorless to slightly yellow)	Score		
Shape		Size		
Flavor	BUBBLE GUM	Imprint Code		
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:51672-2106-8	1 in 1 CARTON	05/20/2011			
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product				
2	NDC:51672-2106-1	1 in 1 CARTON	05/20/2011			
2		240 mL in 1 BOTTLE; Type 0: Not a Combination Product				
3	NDC:51672-2106-4	1 in 1 CARTON	05/20/2011			
3		60 mL in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA201546	05/20/2011		

Labeler - Taro Pharmaceuticals U.S.A., Inc. (145186370)

Establishment				
Name	Address	ID/FEI	Business Operations	
Taro Pharmaceutical Industries, Ltd.		600072078	MANUFACTURE(51672-2106)	

Revised: 1/2020 Taro Pharmaceuticals U.S.A., Inc.