

TEXACLEAR PAIN AND CHEST CONGESTION RELIEF- choline salicylate, guaifenesin liquid
GM Pharmaceuticals, INC

TexaClear Pain & Chest Congestion Relief

TexaClear Pain & Chest Congestion Relief

(Choline Salicylate, Guaifenesin)

NDC: 58809-218-08

GM Pharmaceuticals, Inc.

Drug Facts

Active ingredients (in each 30 mL)

Choline Salicylate (NSAID)* 870 mg

Guaifenesin 200 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever / fever reducer

Expectorant

Uses

- temporarily relieves
 - headache
 - sore throat
 - minor aches and pains
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Salicylates (NSAIDs) may cause a severe allergic reaction which may include:

- hives

- rash
- shock
- skin reddening
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to salicylates (including aspirin) or any other pain reliever/ fever reducer.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn or if you have ulcers or bleeding problems
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, diabetes or thyroid disease
- you have a cough with excessive phlegm (mucus)
- you have a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- you are taking a diuretic
- have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for anticoagulation (thinning the blood) diabetes, gout, or arthritis.
- under a doctor's care for any serious condition

When using this product do not use more than directed

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use salicylates (NSAID) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not exceed recommended dosage.**
- do not take more than 6 doses in any 24-hour period.
- use enclosed dose cup and keep with product
- mL= milliliter

Age	Dose
adults and children 12 years of age and over:	30 mL every 4 hours
children under 12 years of age:	do not use.

Other information

- each 30mL contains: **Sodium 17 mg**
- read all product information before using
- store at 68° to 86°F (20° to 30°C)

Inactive ingredients

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol

Questions or comments?

Call 1-888-535-0305 9 a.m.-5 p.m. CST

PRINCIPAL DISPLAY PANEL

NDC 58809-218-08

TexaClear Pain & Chest Congestion Relief

8 fl oz (237 mL)

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP IS BROKEN OR MISSING

NDC 58809-218-08

TEXACLEAR

PAIN & CHEST CONGESTION RELIEF

**FAST ACTING
FREE & CLEAR
FORMULA**

- Dye Free
- Sugar Free
- Gluten Free
- Alcohol Free
- Acetaminophen Free

Choline Salicylate 870 mg - Pain Reliever/Fever Reducer (**NSAID**)
Guaifenesin 200 mg - Expectorant

Headache • Sore Throat
Thins Mucus • Reduces Fever
Chest Congestion

FAST ACTING

• Ages: 12+
• 30 mL every 4 hours

LIFT HERE FOR COMPLETE DRUG FACTS

Distributed by:
GM Pharmaceuticals, Inc.
Arlington, TX 76018
www.texaclear.com
201086-01 Rev. 0925

8 fl oz (237 mL)

U.S. Pat: 10,987,365

3 58809 21808 0

Drug Facts

Active ingredients (in each 30 mL) Purpose

Choline Salicylate (NSAID)* 870 mg.....Pain reliever/fever reducer
Guaifenesin 200 mg.....Expectorant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves
 - headache
 - sore throat
 - minor aches and pains
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Salicylates (NSAIDs) may cause a severe allergic reaction which may include:

- hives
- rash
- shock
- skin reddening
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to salicylates (including aspirin) or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn or if you have ulcers or bleeding problems
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, diabetes or thyroid disease
- you have a cough with excessive phlegm (mucus)
- you have a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout, or arthritis.
- under a doctor's care for any serious condition

When using this product do not use more than directed ▶

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- pain, cough, or congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use salicylates (NSAID) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage.
- do not take more than 6 doses in any 24-hour period.
- use enclosed dose cup and keep with product
- mL= milliliter

Age	Dose
adults and children 12 years of age and over	30 mL every 4 hours
children under 12 years of age	do not use

Other information

- each 30mL contains: **Sodium 17 mg**
- read all product information before using
- store at 20° to 30°C (68° to 86°F)

Inactive ingredients

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol

Questions or comments?

Call 1-888-535-0305 9 a.m. - 5 p.m. CST

TEXACLEAR PAIN AND CHEST CONGESTION RELIEF

choline salicylate, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58809-218
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 30 mL
CHOLINE SALICYLATE (UNII: KD510K1IQW) (SALICYLIC ACID - UNII:O414PZ4LPZ)	CHOLINE SALICYLATE	870 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58809-218-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/25/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/25/2023	

Labeler - GM Pharmaceuticals, INC (793000860)

Revised: 10/2025

GM Pharmaceuticals, INC