

OPIUM TINCTURE DEODORIZED- morphine tincture solution
ANI Pharmaceuticals, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Opium Tincture, USP (Deodorized) CII
Rx Only

DESCRIPTION

Opium Tincture, USP (Deodorized), is for oral administration. It is freed from unpleasant odor or nauseating substances by “denarcotization” with a petroleum distillate. Opium tincture is a clear, reddish-brown hydroalcoholic solution.

Each 100 mL contains 1 g of anhydrous morphine (represents the equivalent of 10 g of Powdered Opium, USP), alcohol, 19%, and water.

Opium has a very characteristic odor and a very bitter taste. The opiates are the most effective and prompt-acting nonspecific antidiarrheal agents.

CLINICAL PHARMACOLOGY

The opiates act by enhancing tone in the long segments of longitudinal muscle and by inhibiting propulsive contraction of both circular and longitudinal muscles.

Clinical evidence indicates that the analgesic activity of opium preparations is due to their morphine content. Relatively small doses that do not produce significant analgesia are effective in controlling diarrhea. The papaverine content of the mixed alkaloids is too small to have demonstrable spasmolytic activity.

Following oral administration, morphine is well absorbed from the gastrointestinal tract. It is rapidly metabolized following oral administration, however, and plasma levels of unconjugated morphine are lower than those achieved after parenteral administration. Like other narcotic analgesics, opium preparations are metabolized in the liver. Morphine undergoes conjugation with glucuronic acid at the 3-hydroxyl group. Secondary conjugation may also occur at the 6-hydroxyl group to form the 3,6-diglucuronide. Morphine is excreted in the urine mainly as morphine-3-glucuronide and smaller amounts of morphine-3,6-diglucuronide and unchanged drug. Approximately 75% of a dose of morphine is excreted in the urine within 48 hours.

INDICATIONS AND USAGE

Opium tincture is useful for the treatment of diarrhea.

CONTRAINDICATIONS

This preparation is not recommended for use in children. It should not be used in diarrhea caused by poisoning until the toxic material is eliminated from the gastrointestinal tract.

WARNINGS

Addiction can result from opium usage. Opium preparations should be given in the smallest effective dose and as infrequently as possible to minimize the development of tolerance and physical dependence.

PRECAUTIONS

General – Opium should be used with caution in the elderly, in debilitated individuals, and in patients with increased intracranial pressure, cerebral arteriosclerosis, hepatic cirrhosis or liver insufficiency, gastrointestinal hemorrhage, myxedema, emphysema, and bronchial asthma.

Drug Interactions – When preparations containing opium are administered in combination with other drugs, the cautions applicable to each ingredient should be borne in mind. Reduced dosage is indicated in poor-risk patients, in the very young or very old patient, and in those who are receiving other central-nervous system depressants.

Usage in Pregnancy – Animal reproduction studies have not been conducted with Opium Tincture, USP (Deodorized). It is also not known whether Opium Tincture, USP (Deodorized), can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Opium Tincture, USP (Deodorized), should be given to a pregnant woman only if clearly needed.

Nursing Mothers – Caution should be exercised when Opium Tincture, USP (Deodorized), is administered to a nursing woman.

Usage in Children – Safety and effectiveness in children have not been established. See **CONTRAINDICATIONS**.

ADVERSE REACTIONS

Constipation, nausea, and vomiting may occur in some patients. Pruritus and urticaria have been observed.

To report SUSPECTED ADVERSE REACTIONS, contact ANI Pharmaceuticals, Inc. at 1-855-204-1431 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG ABUSE AND DEPENDENCE

Controlled Substance – Opium Tincture, USP (Deodorized), is a Schedule II drug. See **WARNINGS**.

OVERDOSAGE

Signs and Symptoms – Symptoms of toxicity are those of morphine and alcohol, such as nausea, vomiting, miosis, cool and clammy skin, respiratory and CNS depression, bradycardia, hypotension, and skeletal muscle flaccidity. Noncardiogenic pulmonary edema may develop after opioid overdose and monitoring of heart filling pressure may be helpful. Ethanol has been demonstrated to cause hypoglycemia in children or adults with limited glycogen stores. In severe overdosage, apnea, circulatory collapse, cardiac arrest, and death may occur.

Treatment – To obtain up-to-date information about the treatment of overdose, a good resource is your certified Regional Poison Control Center. Telephone numbers of certified poison control centers are listed in the *Physicians' Desk Reference (PDR)*. In managing overdosage, consider the possibility of multiple drug overdoses, interaction among drugs, and unusual drug kinetics in your patient.

Initial management of opioid overdose should emphasize establishment of a secure airway and support of ventilation and perfusion. Meticulously monitor and maintain, within acceptable limits, the patient's vital signs, blood gases, serum electrolytes, blood glucose, etc. Naloxone antagonizes most effects of opioids. Protect the airway as naloxone may induce vomiting. Since naloxone has a shorter duration of action than opioids, repeated doses may be needed. In patients who abuse opioids chronically, a withdrawal syndrome may be manifest on administration of naloxone. This may include yawning, tearing, restlessness, sweating, dilated pupils, piloerection, vomiting, diarrhea, and abdominal cramps. This syndrome usually abates quickly as the effect of naloxone dissipates.

Absorption of drugs from the gastrointestinal tract may be decreased by giving activated charcoal, which, in many cases, is more effective than emesis or lavage; consider charcoal instead of or in addition to gastric emptying. Repeated doses of charcoal over time may hasten elimination of some drugs that have been absorbed. Since opium tincture can delay gastric emptying, evacuation of the stomach may be beneficial. Safeguard the patient's airway when employing gastric emptying or charcoal.

Monitor for and treat hypoglycemia.

The use of forced diuresis, peritoneal dialysis, hemodialysis or charcoal hemoperfusion has not been established to be beneficial.

DOSAGE AND ADMINISTRATION

Usual Adult Dose – 0.6 mL orally 4 times a day.

HOW SUPPLIED

Opium Tincture, USP (Deodorized), is supplied as:
NDC 62559-153-04

Bottles of 4 fl oz

Dispense in tightly closed containers.

Store at controlled room temperature, 68° to 77°F (20° to 25°C) (see USP).

Distributed by:

ANI Pharmaceuticals, Inc.

Baudette, MN 56623



9424 Rev 01/25

Package/Label Display Panel

Opium Tincture USP (Deodorized), 10 mg/mL of anhydrous morphine, CII
NDC 62559-153-04

Rx only

118 mL (4 fl. oz.)

NDC 62559-153-04

Opium **CII**
Tincture USP
(Deodorized) **POISON**

10 mg/mL
of anhydrous morphine

Each 100 mL contains: 1 g of anhydrous morphine (represents the equivalent of 10 g of Powdered Opium USP), and alcohol, 19%.

Rx only
118 mL (4 fl. oz.)

Dispense in tight, light-resistant container as defined in the USP.

Usual Adult Dosage:
0.6 mL orally 4 times a day.
See attached insert for full prescribing information.
Keep tightly closed.

Store at controlled room temperature between 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

Manufactured by:
ANI Pharmaceuticals, Inc.
Baudette, MN 56623
9423 Rev 12/16

GTIN: 00362559000000
EXP: MMM 0000
LOT: 123456
SN: A1234567890123



N 3 62559-153-04 1

OPIUM TINCTURE DEODORIZED

morphine tincture solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62559-153
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MORPHINE (UNII: 76I7G6D29C) (MORPHINE - UNII:76I7G6D29C)	MORPHINE	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	BROWN (Reddish-Brown)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62559-153-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/25/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		02/25/2011	

Labeler - ANI Pharmaceuticals, Inc. (145588013)

Registrant - ANI Pharmaceuticals, Inc. (145588013)

Revised: 9/2025

ANI Pharmaceuticals, Inc.