

**TOPCARE NITE TIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution
Topco Associates LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Nite Time Cold & Flu Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- headache
- fever
- minor aches and pains
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur

if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 42 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

TopCare® health

COMPARE TO VICKS® NYQUIL® SEVERE +
VAPOCOOL™ ACTIVE INGREDIENTS

VAPOR ICE®

Nite Time Cold & Flu

SEVERE

PAIN RELIEVER-FEVER REDUCER - ACETAMINOPHEN

NASAL DECONGESTANT - PHENYLEPHRINE HCl

ANTIHISTAMINE - DOXYLAMINE SUCCINATE

COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

ALCOHOL 10%

12 FL OZ (355 mL)

Maximum Strength

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are
 ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 42 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCARE ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007 ©19P00 PERAD022
 topcare@topcare.com www.topcareand.com
 Visit here for more information:
 http://topcarebrands.com/490110

GLUTEN FREE **QUALITY GUARANTEED**

*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe+ VapoCool™.

NDC 36800-594-40

TopCare
health

COMPARE TO VICKS® NYQUIL® SEVERE+ VAPOCOOL™ ACTIVE INGREDIENTS*

VAPOR ICE®

Nite Time

Cold & Flu

SEVERE

PAIN RELIEVER-FEVER REDUCER - ACETAMINOPHEN
 NASAL DECONGESTANT - PHENYLEPHRINE HCl
 ANTIHISTAMINE - DOXYLAMINE SUCCINATE
 COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr

- Minor Aches & Pains, Fever
- Nasal Congestion & Sinus Pressure
- Sneezing, Runny Nose
- Cough

ALCOHOL 10%

12 FL OZ (355 mL) **Maximum Strength**

Drug Facts **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:
 ■ nasal congestion ■ sinus congestion and pressure ■ headache ■ fever
 ■ minor aches and pains ■ sore throat ■ runny nose and sneezing
 ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep
 ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage
 ■ temporarily restores freer breathing through the nose

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 ■ a sodium-restricted diet

CODE AREA
 * no varnish * no color
 : 59B40 88 F3

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-594
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL	
Inactive Ingredients			
Ingredient Name	Strength		
ALCOHOL (UNII: 3K9958V90M)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
EDETATE DISODIUM (UNII: 7FLD91C86K)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			

GLYCERIN (UNII: PDC6A3C00X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-594-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/01/2019	

Labeler - Topco Associates LLC (006935977)

Revised: 10/2022

Topco Associates LLC