

**DAYTIME COLD FLU NON DROWSY- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled  
Harris Teeter, LLC**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches & pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - sinus pressure
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

## Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

## Other information

- store between 15–30°C (59–86°F)
- avoid excessive heat

## Inactive ingredients

FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

## Questions or comments?

Call **1-877-753-3935 Monday- Friday 9AM- 5PM EST**

## Principal Display Panel

†Compare to the active ingredients in Vicks® DayQuil™ Cold & Flu LiquiCaps™

Daytime

Cold & Flu Relief

MULTI-SYMPTOM RELIEF

**ACETAMINOPHEN ,**

DEXTROMETHORPHAN HBr,

PHENYLEPHRINE HCl

- ACHES, FEVER, SORE THROAT
- COUGH
- NASAL CONGESTION
- SINUS PRESSURE

Alcohol-Free

Antihistamine-Free

Softgels\*\*

(\*\*Liquid Filled Capsules)

†This product is not manufactured or distributed by the Procter & Gamble Company. Vicks® , DayQuil™ and Liquicaps™ are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**PROUDLY DISTRIBUTED BY:**

**HARRIS TEETER, LLC**

**MATTHEWS, NC 28105**

**Product Label**

<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each softgel)</b></p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer  Dextromethorphan HBr 10 mg.....Cough suppressant  Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>■ temporarily relieves common cold and flu symptoms: <ul style="list-style-type: none"> <li>■ minor aches and pains</li> <li>■ headache</li> <li>■ sore throat</li> <li>■ nasal congestion</li> <li>■ fever</li> <li>■ sinus pressure</li> <li>■ cough due to minor throat and bronchial irritation</li> </ul> </li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> <li>■ more than 4,000 mg of acetaminophen in 24 hours</li> <li>■ with other drugs containing acetaminophen</li> <li>■ 3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> <li>■ skin reddening</li> <li>■ blisters</li> <li>■ rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> <ul style="list-style-type: none"> <li>■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p><b>Ask a doctor before use if you have</b> <ul style="list-style-type: none"> <li>■ liver disease</li> <li>■ diabetes</li> <li>■ heart disease</li> <li>■ thyroid disease</li> <li>■ high blood pressure</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>■ cough that occurs with too much phlegm (mucus)</li> </ul> </p></p></p></p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</b></p> <p><b>When using this product, do not exceed recommended dosage.</b></p> <p><b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ pain, cough, or nasal congestion gets worse or lasts more than 7 days</li> <li>■ nervousness, dizziness, or sleeplessness occur</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ redness or swelling is present</li> <li>■ new symptoms occur</li> <li>■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b></p> <p><b>Keep out of reach of children.</b></p> <p><b>Overdose warning:</b> Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b> <ul style="list-style-type: none"> <li>■ do not take more than directed (see Overdose warning)</li> <li>■ do not take more than 4 doses in 24 hours</li> <li>■ adults and children 12 years and over: take 2 softgels with water every 4 hours</li> <li>■ swallow whole; do not crush, chew, or dissolve</li> <li>■ children under 12 years: do not use</li> <li>■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</li> </ul> <p><b>Other information</b> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> FD&amp;C red #40, FD&amp;C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b>  Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p> </p></p></p>
--	--

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PLD-G584C  
FC009599

Lot No.:  
Exp. Date:

**Harris Teeter Daytime Cold and Flu Relief**

**DAYTIME COLD FLU NON DROWSY**  
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69256-898
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K55787QVP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-898-24	24 in 1 CARTON	11/29/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/29/2019	

**Labeler** - Harris Teeter, LLC (048463103)

Revised: 9/2025

Harris Teeter, LLC