

WEAKNESS- iridium pellet
Natural Health Supply

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Indications

Indications: To be used for acute self-limiting conditions according to standard homeopathic indications

Active Ingredient

IRIDIUM METALLICUM

Directions

Take at onset of symptoms. Repeat every 2 hours or as needed until relieved. If condition persists or worsens discontinue use and consult a practitioner.

Adults: dissolve 5-10 pellets in 1 oz. of filtered water or take dry by mouth. Children and infants: 1-5 pellets.

Keep Out of Reach of Children

Keep these and all medications out of the reach of children.

Warning

If pregnant or nursing, consult a practitioner before using.

Inactive Ingredients

Inactive Ingredients - Lactose, Sucrose

Purpose

Purpose: WEAKNESS

Package Label

Est. No. 1724217

Natural Health Supply

505-474-9175

Homeopathic Medicated Pellets

Do not use if clear bottle seal is broken. Do not use if perforated cap seal is broken.

IRIDIUM METALLICUM 30CC

Lot # _____

Est. No. 1724217
Natural Health Supply
 505-474-9175

Homeopathic Medicated Pellets

Indications: to be used for acute, self-limiting conditions according to standard homeopathic indications.
Directions: Take at onset of symptoms. Repeat every 2 hours or as needed until relieved. If condition persists or worsens discontinue use and consult a practitioner. Adults: dissolve 5-10 pellets in 1 oz. of filtered water or take dry by mouth; Children and Infants: 1-5 pellets.
Warning: Keep these and all medicines out of the reach of children; if pregnant or nursing, consult a practitioner before using.
 Inactive Ingredients: Sucrose/Lactose

7.5ML (1/4 OZ) LOT #

Do not use if clear bottle seal is broken.
IRIDIUM METALLICUM
30C
Weakness

Est. No. 1724217

Natural Health Supply
 505-474-9175
 Homeopathic Medicated Pellets

Indications: to be used for acute, self-limiting conditions according to standard homeopathic indications.

Directions: Take at onset of symptoms. Repeat every 2 hours or as needed until relieved. If condition persists or worsens discontinue use and consult a practitioner. Adults: dissolve 5-10 pellets in 1 oz. of filtered water or take dry by mouth; Children and Infants: 1-5 pellets.

Warning: Keep these and all medicines out of the reach of children; if pregnant or nursing, consult a practitioner before using.

Inactive Ingredients: Sucrose/Lactose

1.875 ml (1/16 oz.)

LOT #

IRIDIUM METALLICUM
30C
Weakness

Do not use if perforated cap seal is broken.

res

WEAKNESS

iridium pellet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:64117-240
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IRIDIUM (UNII: 44448S9773) (IRIDIUM - UNII:44448S9773)	IRIDIUM	30 [hp_C] in 1 mL


Inactive Ingredients

Ingredient Name	Strength
LACTOSE (UNII: J2B2A4N98G)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64117-240-01	1.875 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	06/04/1998	

2	NDC:64117-240-02	7.5 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	06/04/1998
---	------------------	--	------------



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		06/04/1998	



Labeler - Natural Health Supply (018504618)

Registrant - Natural Health Supply (018504618)

Establishment

Name	Address	ID/FEI	Business Operations
Natural Health Supply		018504618	manufacture(64117-240)

Revised: 1/2019

Natural Health Supply