MULTI-SYMPTOM COLD SEVERE, NON-DROWSY, DAYTIME- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated Valu Merchandisers Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Best Choice 44-503A

Active ingredients (in each caplet)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Guaifenesin 200 mg Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer Cough suppressant Expectorant Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sore throat
 - cough
 - nasal congestion
 - headache
 - minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- heart disease
- liver disease
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, D&C yellow #10 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

Best Choice®

Non-Drowsy Severe Daytime Multi-Symptom Cold

ACETAMINOPHEN - Pain Reliever/Fever Reducer

Dextromethorphan HBr - Cough Suppressant Guaifenesin - Expectorant Phenylephrine HCl - Nasal Decongestant

Actual Size

Relieves Fever/Headache, Sore Throat, Nasal Congestion, Cough, Chest Congestion **Pseudoephedrine Free**

24 Cool Caplets with Cool Blast Flavor

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0718B50308

PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106

Best Choice®
100% Guaranteed
www.bestchoicebrand.com

 children under 12 years: ask a doctor ■ heart disease
■ liver disease
■ diabetes ■ do not take more than 10 caplets in 24 hours Ask a doctor before use if you have swallow whole - do not crush, chew, or dissolve product or any of its ingredients ∎ rake 2 caplets every 4 hours if you have ever had an allergic reaction to this ■ adults and children 12 years and over or pharmacist before taking this product. Directions a do not take more than directed prescription drug contains an MAOI, ask a doctor the MAOI drug. If you do not know if your children even if you do not notice any signs or Parkinson's disease), or for 2 weeks after stopping medical attention is critical for adults as well as for depression, psychiatric or emotional conditions, or oxidase inhibitor (MAM) (certain drugs for Control Center (1-800-222-1222) right away. Prompt ■ if you are now taking a prescription monoamine overdose, get medical help or contact a Poison Keep out of reach of children. In case of accidental doctor or pharmacist. professional before use. ance whether a drug contains acetaminophen, ask a If pregnant or breast-feeding, ask a health (brescription or nonprescription). If you are not with any other drug containing acetaminophen serious condition. asu jou of headache that lasts. These could be signs of a достог рготриу. condit comes back or occurs with rash or redness or swelling is present fever, headache, rash, nausea, or vomiting, consult a Tever gets worse or lasts more than 3 days for more than 2 days, is accompanied or followed by lasts more than 7 days anew symptoms occur Sore throat warning: If sore throat is severe, persists psin, nasal congestion, or cough gets worse or help nght away. It a skin reaction occurs, stop use and seek medical uervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if mskin reddening mblisters mrash reactions, Symptoms may include: recommended dosage. Allergy aled: Acetaminophen may cause severe skin When using this product do not exceed ruis broance taking the blood thinning drug wartarin. ■3 or more alcoholic drinks every day while using Ask a doctor or pharmacist before use if you are ■ with other drugs containing acetaminophen prostate gland Thigh blood pressure amore than 4,000 mg of acstaminophen in 24 hours difficulty in unnation due to enlargement of the Severe liver damage may occur if you take condy that occurs with too much phiegm (mucus) Liver warning: This product contains acetaminophen.

pemporarily reduces fever

Drug Facts (continued)

Drug Facts

vgninseW

B-2115-503A-08CMSR REV0718B50308

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

secretions to make coughs more productive ■ pelps loosen phiegm (mucus) and thin bronchial minor aches and pains ■ uesej coudespou ■ µesqscµe

smoking, asthma, chronic bronchitis, or

persistent or chronic cough such as occurs with

emphysema

nund racts (continued)

smotomys uit bns ■ soce throat ■ cough

■ thyroid disease

Nses I temporarily relieves these common cold **Drug Facts** (continued) Masal decongestant Phenylephrine HCI 5 mg Expectorant Guaifenesin 200 mg Dextromethorphan HBr 10 mg.....Cough suppressant Acetaminophen 325 mg Pain reliever/fever reducer ezoqru9 (felqes daes ni) stneibergni evitaA

COMPLETE PRODUCT INFORMATION

KEEP OUTER PACKAGE FOR

CO BY Best Choice. 100% Guaranteed

PROUDLY DISTRIBUTED BY: VALU MERCHANDISERS, CO. 5000 KANSAS AVE KANSAS CITY, KS 66106

No print/No varnish Lot & Exp date

Drug Facts (continued Questions or comment

Best Choice® **Non-Drowsy** Severe Daytime Multi-Symptom Cold

Actual Size

ACETAMINOPHEN - Pain Reliever/Fever Reducer Dextromethorphan HBr - Cough Suppressant Guaifenesin - Expectorant



REV0718850300



Best Choice 44-503A

MULTI-SYMPTOM COLD SEVERE, NON-DROWSY, DAYTIME

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-503
Route of Administration	ORAL		

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg			
DEXTRO METHO RPHAN HYDRO BRO MIDE (UNII: 9 D2RTI9 KYH) (DEXTRO METHO RPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg			
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg			
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg			

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	

$\textbf{SODIUM STARCH GLYCOLATE TYPE A POTATO} \hspace{0.1cm} \textbf{(UNII: } 5856 \textbf{)} 3G2 A2 \textbf{)}$

Product Characteristics				
Color	YELLOW	Score	no score	
Shape	OVAL	Size	19 mm	
Flavor	MINT	Imprint Code	44;503	
Contains				

ı	Packag	ging			
ı	# Ite	em Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:6	3941-503-08	2 in 1 CARTON	08/04/2005	
ı	1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH FINAL	part341	08/04/2005		

Labeler - Valu Merchandisers Company (868703513)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		832867894	MANUFACTURE(63941-503)	

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		832867837	PACK(63941-503)	

Revised: 10/2019 Valu Merchandisers Company