SODIUM IODIDE 20%- sodium iodide injection, solution Clipper

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sodium Iodide 20% Injection

INDICATIONS:

For use as an aid in the treatment of actinomycosis (lumpy jaw), actinobacillosis (wooden tongue), and necrotic stomatitis in cattle.

CONTRAINDICATIONS:

The use of sodium iodide is contraindicated in pregnancy and hyperthyroidism.

DOSAGE AND ADMINISTRATION:

Using aseptic procedures, administer slowly by intravenous injection. Inject carefully to avoid deposition outside of the vein. The usual dose is 30 mg per pound of body weight (15 mL/100 lb). May be repeated at weekly intervals, if necessary.

WARNING:

Not for use in lactating dairy cows.

CAUTION:

Animals vary in their susceptibility of iodides. Administer with caution until the animal's tolerance is determined. Discontinue treatment and consult veterinarian if adverse reactions occur.

18-819-25

Iss. 04-12

RMS 92-1042

Lot No.

Exp. Date

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

COMPOSITION:

Each 100 mL of sterile aqueous solution contains:

Sodium Iodide......20 grams

Water for injection......q.s.

Store between 15 degrees C and 30 degrees C (59 degrees F and 86 degrees F).

TAKE TIME OBSERVE LABEL DIRECTIONS

Trademarks are property of Clipper Distributing Company, LLC

Manufactured by Nova-Tech Inc. Grand Island, NE 68801

Net Contents: 250 mL

INDICATIONS: For use as an aid in the treatment of actinomycosis (lumpy jaw), actinobacillosis (wooden tongue), and necrotic stomatitis in cattle.

CONTRAINDICATIONS: The use of sodium iodide is contraindicated in pregnancy and hyperthyroidism.

DOSAGE AND ADMINISTRATION: Using aseptic procedures, administer slowly by intravenous injection. Inject carefully to avoid deposition outside of the vein. The usual dose is 30 mg per pound of body weight (15 mL/100 lb). May be repeated at weekly intervals, if necessary.

WARNING: Not for use in lactating dairy cows.

CAUTION: Animals vary in their susceptibility of iodides. Administer with caution until the animal's tolerance is determined. Discontinue treatment and consult veterinarian if adverse reactions occur.

18-819-25

Lot No.

Iss. 04-12

RMS 92-1042

Exp. Date

Sodium lodide 20% Injection

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Net Contents: 250 mL



Manufactured for: Clipper Distributing Company, LLC St. Joseph, MO 64507 **COMPOSITION:** Each 100 mL of sterile aqueous solution contains: Sodium lodide 20 grams Water for injection q.s.

Store between 15°C and 30°C (59°F and 86°F).



Trademarks are property of Clipper Distributing Company, LLC

Manufactured by Nova-Tech Inc. Grand Island, NE 68801



SODIUM IODIDE 20%

sodium iodide injection, solution

Product Information

 Product Type
 PRESCRIPTION ANIMAL DRUG
 Item Code (Source)
 NDC:57319-601

Route of Administration INTRAVENOUS

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

SODIUM IODIDE (UNII: F5WR8 N145C) (IODIDE ION - UNII:09 G4I6 V86Q)

IODIDE ION

20 g in 250 mL

P	ac	kag	g in	g
_			,	0

;	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC ·57319-601-06	250 mL in 1 BOTTLE PLASTIC		

Marketing Information

Marketing CategoryApplication Number or Monograph CitationMarketing Start DateMarketing End Dateunapproved drug other07/01/2012

Labeler - Clipper (150711039)

Registrant - Clipper (150711039)

Establishment						
Name	Address	ID/FEI	Business Operations			
Nova-Tech, Inc		196078976	manufacture, api manufacture			

Revised: 5/2014 Clipper