

BENADRYL ALLERGY PLUS CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride tablet, film coated
Kenvue Brands LLC

Benadryl Allergy Plus Congestion

Drug Facts

<i>Active ingredients (in each tablet)</i>	<i>Purposes</i>
Diphenhydramine HCl 25 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
 - nasal congestion
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dose**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 1 tablet every 4 hours▪ do not take more than 6 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F).
- **do not use if carton is opened or if blister unit is broken**

Inactive ingredients

carnauba wax, FD&C blue no. 1 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-415-01

Benadryl®

***ALLERGY PLUS
CONGESTION***

**Diphenhydramine HCl 25mg | Antihistamine
Phenylephrine HCl 10mg | Nasal Decongestant**

- ✓ **Sneezing**
- ✓ **Runny Nose**
- ✓ **Sinus Congestion**
- ✓ **Itchy, Watery Eyes**
- ✓ **Itchy Throat**
- ✓ **Sinus Pressure**

ULTRATABS®*

*small tablet size

actual size

24 TABLETS

Benadryl®

ALLERGY PLUS
CONGESTION

EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT!®



3 0045-0557-261

Made in Italy
Distributed by: J&JCI
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016

ALLERGY PLUS
CONGESTION

Benadryl®

Questions or comments?
call 1-888-217-2171 (toll-free) or 215-273-8755 (collect)

Inactive ingredients carnauba wax, FD&C blue no. 1 aluminum lake, glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Other information
■ do not use if carton is opened or if blister unit is broken
■ store between 20-25°C (68-77°F)
■ do not use if carton is opened or if blister unit is broken

Directions
adults and children ■ take 1 tablet every 4 hours ■ do not take more than 6 tablets in 24 hours
children under 12 years ask a doctor

Drug Facts (continued)
Stop use and ask a doctor if:
■ nervousness, dizziness, or sleepiness occur
■ symptoms do not improve within 7 days or occur with a fever
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

When using this product ■ do not exceed recommended dose
■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
■ excitability may occur, especially in children

Warnings
Do not use
■ to make a child sleepy
■ with any other product containing diphenhydramine, even one used on skin
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)
■ certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ heart disease
■ thyroid disease
■ diabetes
■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

Uses
■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ sneezing ■ itchy, watery eyes
■ itching of the nose or throat ■ nasal congestion
■ temporarily relieves these symptoms due to the common cold:
■ runny nose ■ sneezing ■ nasal congestion

Drug Facts
Active ingredients (in each tablet)
Diphenhydramine HCl 25 mg
Phenylephrine HCl 10 mg
Purposes
Antihistamine
Nasal decongestant

Benadryl®

ALLERGY PLUS
CONGESTION

EFFECTIVE ALLERGY RELIEF WHEN YOU NEED IT!®

Benadryl®

ALLERGY PLUS
CONGESTION

Diphenhydramine HCl 25mg | Antihistamine
Phenylephrine HCl 10mg | Nasal Decongestant


- ✓ Sneezing
- ✓ Runny Nose
- ✓ Sinus Congestion
- ✓ Itchy, Watery Eyes
- ✓ Itchy Throat
- ✓ Sinus Pressure

NDC 50580-415-01

116786

116786

30037630/116786



BENADRYL ALLERGY PLUS CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-415
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	PE;WL95
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-415-01	2 in 1 CARTON	06/19/2017	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/19/2017	

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC