

**HABITROL NICOTINE POLACRILEX 2 MG COATED CINNAMON- nicotine
polacrilex gum, chewing
Dr. Reddy's Laboratories Inc.**

Nicotine Polacrilex 2 mg Cinnamon Coated USP Gum

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex (equal to 2 mg nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- oral blistering occurs
- you get symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up**, use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up**, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get best results
- chew the gum slowly until it tingles. then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent craving, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider

Other information

- each piece contains: calcium 115 mg, sodium 11 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, butylated hydroxytoluene, carnauba wax, FD&C red #40 lake, gum base, hydroxypropyl cellulose, hypromellose, natural and artificial cinnamon flavors, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

Questions or comments?

Call toll-free 1-**800-585-8682** weekdays (8:00 am - 8:00pm EST)

Principal Display Panel

*Compare to the active ingredient in Nicorette® Gum

Habitrol

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

Gum

2 mg

Sugar-free

Coated Cinnamon Flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Habitrol 4 mg Gum.

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivate to quit
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks
3. **Use long enough** - Use Nicotine Polacrilex Gum for the full 12 weeks
4. **Use with a Support Program** as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- not for sale to those under 18 years of age
- proof of age required

- not for sale in vending machines or from any source where proof of age cannot be verified

*This product is not manufactured or distributed by Haleon group of companies distributor of Nicorette® Gum

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

DISTRIBUTED BY:

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

Package Label

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

NDC 55741-004-80

*Compare to the active ingredient in Nicorette® Gum

Habitrol®

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

Gum

2 mg

Sugar-free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Habitrol 4 mg Gum.

**80
PIECES
2 mg EACH**



Habitrol®

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID



Habitrol, sodium citrate, sorbitol, sucrose, talc, titanium dioxide, xylitol
 (Questions or comments?)
 Call 1-800-545-8822 weekdays (9:00 am - 5:00 pm EST)
 *This product is not manufactured or distributed by Hesse group of companies.
 distributor of Nicorette® Gum.

If you get symptoms or notice an overdose such as nausea, vomiting, dizziness, drowsiness, and rapid heartbeat
 or oral irritation occurs
 if you have symptoms of an allergic reaction (such as difficulty breathing or rash)
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. They need pieces of gum in paper and throw away in the trash in case of overdose. Get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions
 If you are under 18 years of age, ask a doctor before use
 or before using this product. Read the enclosed User's Guide for complete directions and other important information
 in light using the gum on your quit day
 If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum

**80
PIECES
2 mg EACH**



Color	pink (Light Pink)	Score	no score
Shape	SQUARE	Size	14mm
Flavor	CINNAMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55741-004-15	15 in 1 CARTON	07/28/2025	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55741-004-80	8 in 1 CARTON	07/28/2025	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:55741-004-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	07/28/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079216	07/28/2025	

Labeler - Dr. Reddy's Laboratories Inc. (802315887)

Revised: 12/2025

Dr. Reddy's Laboratories Inc.