### COLTALIN- acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet FORTUNE PHARMACAL COMPANY LIMITED

\_\_\_\_\_

## **Drug Facts**

Active ingredients (in each tablet) Acetaminophen 325 mg Chlorpheniramine maleate 2 mg Phenylephrine hydrochloride 5 mg

### Purpose

Pain reliever-fever reducer Antihistamine Nasal decongestant

### Uses

■ temporarily relieves these symptoms due to a cold, the flu, or hay fever:

- minor aches and pains
- sore throat
- headache
- runny nose
- sinus congestion and pressure
- nasal congestion due to a cold or hay fever
- runny nose

■ sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever

- temporarily reduces fever
- promotes nasal and/or sinus drainage

## Warnings

**Liver warning**: This product contains acetaminophen. Severe liver damage may occur if:

 $\blacksquare$  adult takes more than 12 tablets in 24 hours, which the maximum daily amount for this product

- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

### Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or

for 2 weeks after stopping the MAOI drug. If you do not know whether a prescription

drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## Ask a doctor before use if the user has

- heart disease
- high blood pressure
- diabetes
- glaucoma
- liver disease
- thyroid disease
- a high fever
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

### Ask a doctor or pharmacist before use if the user is

- taking tranquilizers or sedatives
- taking the blood thinning drug warfarin

### When using this product

#### do not exceed recommended dosage

- may cause excitability especially in children
- use caution when driving or operating machinery
- avoid alcoholic beverages
- may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness affect

drowsiness effect

#### Stop use and ask a doctor if

■ pain or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults)

- sore throat persists for more than 2 days
- fever gets worse or lasts more than 3 days
- nasal congestion is accompanied by fever
- nervousness, dizziness, or sleeplessness occur
- redness or swelling is present
- new symptoms occur
- any of the following occurs (these could be signs of a serious condition):
  - severe sore throat

■ sore throat is accompanied or followed by fever, headache, rash, nausea or vomiting

Do not give to children under 6 years of age unless directed by a doctor.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

## Directions

■ adults and children 12 years of age and over: 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses (12 tablets) in 24 hours, or as directed by a doctor

children 6 to under 12 years of age: 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor
children under 6 of age: consult a doctor

children under 6 of age : consult a doctor

## Other information

■ keep tightly sealed
■ store between below 25 C (77 F) in a dry place, away from direct sunlight

### Inactive ingredients

colloidal silicon dioxide, corn starch, FD C yellow no. 6 aluminum lake, magnesium stearate, povidone K30, sodium starch glycolate

## Questions or comments? (888) 221-3496 M-F 9 am to 5 pm

■ you may also report serious side effects to this phone number

COLTALIN

Safely Sealed

NDC 51467-011-01

Acetaminophen - Pain Reliever-Fever Reducer Chlorpheniramine maleate - Antihistamine Phenylephrine hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief

- Minor Aches & Pain, Sore Throat
- Headache, Fever
- Nasal Congestion, Sinus Pressure/Drainage
- Sneezing, Runny Nose
- Itchy, Watery Eyes

24 Tablets

In Individually Sealed Strip

Aspirin Free



COLTALIN

Safely Sealed

NDC 51467-011-02

Acetaminophen - Pain Reliever-Fever Reducer Chlorpheniramine maleate - Antihistamine Phenylephrine hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief


- Minor Aches & Pain, Sore Throat
- Headache, Fever
- Nasal Congestion, Sinus Pressure/Drainage
- Sneezing, Runny Nose
- Itchy, Watery Eyes

36 Tablets

In Individually Sealed Strip

Aspirin Free





# COLTALIN

acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet

Dreduct Turne	HUMAN OTC DRUG	Item Code (Source) NDC:51467-0			7 011
Product Type	HUMAN OTC DRUG	Item Code (Source) NDC:514			07-011
Route of Administration	ORAL				
Active Ingredient/Active	Moiety				
Ingredient Name			<b>Basis of Strength</b>		Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)			ACETAMINOPHEN		325 mg
			CHLORPHENIRAMINE MALEATE		2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) PHENYLEPHRINE				E	5 mg
Inactive Ingredients					
Inactive Ingredients	Ingredient Name			S	trength
Inactive Ingredients SILICON DIOXIDE (UNII: ETJ7Z6XE	•			S	trength
-	3U4)			S	trength
SILICON DIOXIDE (UNII: ETJ7Z6XE	3U4) 5J)			S	trength
SILICON DIOXIDE (UNII: ETJ7Z6XE STARCH, CORN (UNII: 08232NY35	3U4) 5J) /EI93A8)			S	trength
SILICON DIOXIDE (UNII: ETJ7Z6XE STARCH, CORN (UNII: 08232NY35 FD&C YELLOW NO. 6 (UNII: H77)	3U4) 5J) /EI93A8) 097M6I30)			S	trength

Product Characteristics						
Color	orange	Score	no score			
Shape	ROUND	Size	12mm			
Flavor		Imprint Code	SHIELDTM			
Contains						

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:51467- 011-01	2 in 1 BOX	12/26/2024			
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product				
2	NDC:51467- 011-02	3 in 1 BOX	12/26/2024			
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product				
Marketing Information						
M	larketing	Information				
M	larketing Marketing Category	nformation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		

Labeler - FORTUNE PHARMACAL COMPANY LIMITED (686280561)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
FORTUNE PHARMACAL COMPANY LIMITED		686280561	manufacture(51467-011)	

Revised: 12/2024

FORTUNE PHARMACAL COMPANY LIMITED