


MECLIZINE HYDROCHLORIDE - meclizine hydrochloride tablet
STAT Rx USA LLC

DESCRIPTION

Meclizine hydrochloride, an oral antiemetic, is a white, slightly yellowish, crystalline powder which has a slight odor and is tasteless. It has the following structural formula:



C₂₅H₂₇ClN₂•2HCl•H₂O M.W. 481.89

The chemical name is 1-(*p*-chloro- α -phenylbenzyl)-4-(*m*-methyl-benzyl) - piperazine dihydrochloride monohydrate.

Meclizine Hydrochloride Tablets are available in 12.5 mg, and *25 mg strengths for oral administration.

*Contains FD&C Yellow #5 (see PRECAUTIONS).

Each tablet contains the following inactive ingredients: colloidal silicon dioxide, lactose, magnesium stearate, microcrystalline cellulose, sodium starch glycolate, starch, and stearic acid. In addition, the 12.5 mg tablet contains FD&C Blue #1; and the 25 mg tablet contains D&C Yellow #10 and FD&C Yellow #5.

CLINICAL PHARMACOLOGY

Meclizine hydrochloride is an antihistamine which shows marked protective activity against nebulized histamine and lethal doses of intravenously injected histamine in guinea pigs. It has a marked effect in blocking the vasodepressor response to histamine, but only a slight blocking action against acetylcholine. Its activity is relatively weak in inhibiting the spasmogenic action of histamine on isolated guinea pig ileum.

INDICATIONS AND USAGE

For the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

CONTRAINDICATIONS

Meclizine hydrochloride is contraindicated in individuals who have shown a previous hypersensitivity to it.

WARNINGS

Since drowsiness may, on occasion, occur with use of this drug, patients should be warned of this

possibility and cautioned against driving a car or operating dangerous machinery.

Patients should avoid alcoholic beverages while taking the drug. Due to its potential anticholinergic action, this drug should be used with caution in patients with asthma, glaucoma, or enlargement of the prostate gland. Do not give to children under 12 years of age unless directed by a doctor.

PRECAUTIONS

The Meclizine Hydrochloride Tablets, 25 mg contain FD&C Yellow #5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible individuals. Although the overall incidence of FD&C Yellow #5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

Usage in Children: Clinical studies establishing safety and effectiveness in children have not been done; therefore, usage is not recommended under 12 years of age.

Usage in Pregnancy: *Pregnancy Category B.* Reproduction studies in rats have shown cleft palates at 25-50 times the human dose. Epidemiological studies in pregnant women, however, do not indicate that meclizine hydrochloride increases the risk of abnormalities when administered during pregnancy.

Despite the animal findings, it would appear that the possibility of fetal harm is remote. Nevertheless, meclizine hydrochloride, or any other medication should be used during pregnancy only if clearly necessary.

ADVERSE REACTIONS

Drowsiness, dry mouth, and on rare occasions, blurred vision have been reported.

DOSAGE AND ADMINISTRATION

Motion Sickness: The initial dose of 25 to 50 mg meclizine hydrochloride, should be taken one hour prior to travel for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

HOW SUPPLIED

Meclizine Hydrochloride Tablets, USP 12.5 mg - blue, oval tablets debossed with "034" on one side and "par" on the other side. Tablets may contain characteristic dye spots. They are supplied in:

Bottles of 30 NDC 16590-301-30

Bottles of 60 NDC 16590-301-60

Bottles of 90 NDC 16590-301-90

Meclizine Hydrochloride Tablets, USP 25 mg - yellow, oval tablets debossed with "035" on one side and "par" on the other side. They are supplied in:

Bottles of 14 NDC 16590-146-14

Bottles of 15 NDC 16590-146-15

Bottles of 20 NDC 16590-146-20

Bottles of 30 NDC 16590-146-30

Bottles of 40 NDC 16590-146-40

Bottles of 60 NDC 16590-146-60

Bottles of 90 NDC 16590-146-90

Dispense in tight, light-resistant containers as defined in the USP.

Store at controlled room temperature 15°-30°C (59°-86°F).

Manufactured by:

PAR PHARMACEUTICAL COMPANIES, INC.

Spring Valley, NY 10977

Revised: 11/09

OS034-01-1-11

Relabeling and Repackaging by:

STAT Rx USA LLC

Gainesville, GA

PACKAGE LABEL - MECLIZINE 12.5 MG TABLETS

S Packaged and Distributed By: **STAT Rx USA** Gainesville, GA 30501

Meclizine
12.5mg **30 Tabs**

Generic For: **Antivert**

NDC 16590-301-30 Prod# 301-30
Each Tablet Contains: Meclizine Hcl., USP. Lot# SAMPLE

12.5mg

Mfg By: Par Pharmaceutical Co., Inc.
Spring Valley, NY 10977 NDC 49884-034-01

Mfg Lot: SAMPLE
Discard After: 06/13 BW 9/18/2012 SAMPLE

CH RDA
Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.

May cause DROWSINESS. ALCOHOL may INTENSIFY this effect. Use care when operating dangerous machinery.

Dosage: See package insert
Store between 68-77 degrees F

RX ONLY-KEEP OUT OF REACH OF CHILDREN

PACKAGE LABEL - MECLIZINE 25 MG TABLETS

S Packaged and distributed by: **STAT Rx USA** LLC Gainesville, GA 30501

25mg Meclizine 30 Tabs

Generic For: **Antivert**

NDC 16590-146-30 Prod# 146-30 Lot# SAMPLE

Each Tablet Contains: Meclizine Hydrochloride USP, 25mg

Mfg By: Par Pharmaceutical Co., Inc. Spring Valley, NY 10977 NDC 49884-035-20

Mfg Lot: 2373956 Discard After: 10/13 BW 6/6/2012 SAMPLE

RX ONLY-KEEP OUT OF REACH OF CHILDREN

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed. May cause DROWSINESS. ALCOHOL may INTENSIFY this effect. Use care when operating dangerous machinery.

0

MECLIZINE HYDROCHLORIDE

meclizine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16590-301(NDC:49884-034)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	12.5 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
WATER (UNII: 059QF0K00R)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	5mm
Flavor		Imprint Code	Par;034
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16590-301-30	30 in 1 BOTTLE		
2	NDC:16590-301-60	60 in 1 BOTTLE		
3	NDC:16590-301-90	90 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA087127	06/03/1981	

MECLIZINE HYDROCHLORIDE

meclizine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16590-146(NDC:49884-035)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
WATER (UNII: 059QF0KO0R)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	6mm
Flavor		Imprint Code	Par;035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16590-146-14	14 in 1 BOTTLE		
2	NDC:16590-146-15	15 in 1 BOTTLE		
3	NDC:16590-146-20	20 in 1 BOTTLE		
4	NDC:16590-146-30	30 in 1 BOTTLE		
5	NDC:16590-146-40	40 in 1 BOTTLE		
6	NDC:16590-146-60	60 in 1 BOTTLE		
7	NDC:16590-146-90	90 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA087128	06/03/1981	

Labeler - STAT Rx USA LLC (786036330)**Registrant** - PSS World Medical Inc. (101822682)**Establishment**

Name	Address	ID/FEI	Business Operations
STAT Rx USA LLC		786036330	relabel(16590-301, 16590-146) , repack(16590-301, 16590-146)

Revised: 9/2012

STAT Rx USA LLC