

# **NITROFURANTOIN- nitrofurantoin suspension**

## **Biocon Pharma Inc**

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use NITROFURANTOIN ORAL SUSPENSION safely and effectively. See full prescribing information for NITROFURANTOIN ORAL SUSPENSION.**

**NITROFURANTOIN oral suspension**  
**Initial U.S. Approval: 1953**

### **RECENT MAJOR CHANGES**

Dosage and Administration (2) 1/2024

### **INDICATIONS AND USAGE**

Nitrofurantoin oral suspension is a nitrofuran antibacterial indicated in adults and pediatric patients 1 month of age and older for the treatment of urinary tract infections caused by susceptible bacteria. (1)

#### Limitations of Use

Nitrofurantoin oral suspension is not indicated for the treatment of pyelonephritis or perinephric abscesses. (1)

#### Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of nitrofurantoin oral suspension and other antibacterial drugs nitrofurantoin oral suspension should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. (1)

### **DOSAGE AND ADMINISTRATION**

- Adult Patients: 50 mg to 100 mg four times a day - the lower dosage level is recommended for uncomplicated urinary tract infections. (2.2)
- Pediatric Patients: 5 mg/kg to 7 mg/kg of body weight per 24 hours, given in four divided doses (contraindicated under one month of age). (2.3)

### **DOSAGE FORMS AND STRENGTHS**

Oral Suspension: 25 mg/5 mL. (3)

### **CONTRAINDICATIONS**

- Known hypersensitivity to nitrofurantoin. (4)
- History of cholestatic jaundice/hepatic dysfunction associated with nitrofurantoin. (4)
- Patients who have anuria, oliguria, or significant impairment of renal function (creatinine clearance under 60 mL per minute or clinically significant elevated serum creatinine). (4)
- Pregnant patients at term (38 weeks to 42 weeks gestation), during labor and delivery, or when the onset of labor is imminent. (4)
- Pediatric Patients under one month of age. (4)

### **WARNINGS AND PRECAUTIONS**

- Hypersensitivity Reactions: Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving nitrofurantoin oral suspension. If signs and symptoms of a hypersensitivity reaction occurs, immediately discontinue nitrofurantoin oral suspension. (5.1)
- Pulmonary Reactions: Discontinue if sign and symptoms of pulmonary reactions occur and take appropriate measures. (5.2)
- Hepatotoxicity: Discontinue if signs/symptoms of hepatitis occur. Monitor liver function tests. (5.3)
- Neuropathy: Peripheral neuropathy has occurred. (5.4)
- Hemolytic Anemia: Discontinue if sign and symptoms of hemolysis occur. (5.5)
- *Clostridioides difficile*-associated diarrhea: Evaluate patients if diarrhea occurs. (5.6)

### **ADVERSE REACTIONS**

The most common adverse reactions occurring in patients are nausea, vomiting, anorexia, vertigo, nystagmus, dizziness, headache, angioedema, rash, urticaria, pulmonary hypersensitivity reactions, hepatic reactions, peripheral neuropathy, increased aspartate aminotransferase, increased alanine aminotransferase, decreased hemoglobin, increased serum phosphorus, and eosinophilia (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Biocon Pharma Inc., at 1-866-924-6266 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## -----**DRUG INTERACTIONS**-----

- Antacids: Decreased absorption of nitrofurantoin. (7.1)
- Uricosuric drugs: Inhibit renal tubular secretion of nitrofurantoin. (7.2)

## -----**USE IN SPECIFIC POPULATIONS**-----

- Lactation: Breastfeeding is not recommended in infants under one month of age or with glucose-6-phosphate dehydrogenase (G6PD) deficiency. (8.2)
- Pediatric Use: Nitrofurantoin oral suspension is contraindicated in pediatric patients under one month of age. (8.4)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 10/2024**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Recommended Dosage and Administration in Adult Patients

2.2 Recommended Dosage and Administration in Pediatric Patients (1 month of age and older)

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Hypersensitivity Reactions

5.2 Pulmonary Reactions

5.3 Hepatotoxicity

5.4 Neuropathy

5.5 Hemolytic Anemia

5.6 *Clostridioides difficile*-associated Diarrhea

5.7 Persistence or Reappearance of Bacteriuria

### **6 ADVERSE REACTIONS**

### **7 DRUG INTERACTIONS**

7.1 Antacids

7.2 Uricosuric Drugs

7.3 Drug Interference with Laboratory Tests

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Nitrofurantoin oral suspension is indicated in adults and pediatric patients 1 month of age and older for the treatment of urinary tract infections due to susceptible strains of *Escherichia coli*, *Enterococcus* species, *Staphylococcus aureus*, *Klebsiella* species and *Enterobacter* species.

#### Limitations of Use

Nitrofurantoin oral suspension is not indicated for the treatment of pyelonephritis or perinephric abscesses [see *Warnings and Precautions (5.7)*].

#### Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of nitrofurantoin oral suspension and other antibacterial drugs, nitrofurantoin oral suspension should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Recommended Dosage and Administration in Adult Patients**

The recommended dosage is 50 mg to 100 mg of nitrofurantoin oral suspension four times a day.

For long-term suppressive therapy in adults, a reduction of dosage to 50 mg to 100 mg at bedtime may be adequate. The benefits of long-term suppressive therapy should be balanced against the increased potential for systemic toxicity and for the development of antibacterial resistance [see *Warnings and Precautions (5.2, 5.4, 5.6)*].

Administer nitrofurantoin oral suspension with food to improve drug absorption [see *Clinical Pharmacology (12.3)*] and, in some patients, tolerance.

#### **2.2 Recommended Dosage and Administration in Pediatric Patients (1 month of age and older)**

The recommended dosage of nitrofurantoin oral suspension is 5 mg/kg to 7 mg/kg of body weight per 24 hours, given in four divided doses in pediatric patients aged 1 month and older. Administer nitrofurantoin oral suspension with food to improve drug absorption [see *Clinical Pharmacology (12.3)*] and, in some patients, tolerance.

Table 1 lists individual dosage volumes for two different strengths of nitrofurantoin oral suspension (25 mg/5 mL and 50 mg/5 mL) based on body weight for pediatric patients.

**Table 1: Nitrofurantoin Oral Suspension Pediatric Dosing Table for Pediatric Patients 1 month of Age and Older**

Weight in Kilograms (kg)	Pediatric Doses (mL), Frequency: Four times Daily	
	25 mg/5 mL oral suspension	50 mg/5 mL oral suspension
4 kg or lower	1	0.5
Greater than 4 kg to 5 kg	1.4	0.7
Greater than 5 kg to 7 kg	1.8	0.9
Greater than 7 kg to 10 kg	2.5	1.25
Greater than 10 kg to 14 kg	3.5	1.75
Greater than 14 kg to 20 kg	5	2.5
Greater than 20 kg to 25 kg	7	3.5
Greater than 25 kg to 30 kg	8.5	4.25
Greater than 30 kg to 40 kg	10	5
40 kg or greater	See Adult Dose	See Adult Dose

To measure the correct pediatric doses, it is important to administer nitrofurantoin oral suspension with an appropriate size oral dosing syringe with graduations that align with the volume prescribed in Table 1 above.

Continue therapy for one week or for at least 3 days after sterility of the urine is obtained. Continued infection indicates the need for reevaluation.

For long-term suppressive therapy in pediatric patients, doses as low as 1 mg/kg per 24 hours, given in a single dose or in two divided doses, may be adequate [see *Warnings and Precautions* (5.2, 5.4, 5.6)].

### 3 DOSAGE FORMS AND STRENGTHS

Nitrofurantoin oral suspension, USP is available as an opaque, yellow liquid oral suspension containing 25 mg/5 mL of nitrofurantoin.

### 4 CONTRAINDICATIONS

Nitrofurantoin oral suspension is contraindicated in:

- patients with known hypersensitivity to nitrofurantoin [see *Warnings and Precautions* (5.1)].
- patients with a previous history of cholestatic jaundice/hepatic dysfunction associated

with nitrofurantoin [see *Warnings and Precautions (5.3)*].

- patients who have anuria, oliguria, or significant impairment of renal function (creatinine clearance under 60 mL per minute or clinically significant elevated serum creatinine) due to an increased risk of toxicity resulting from impaired excretion of the drug [see *Warnings and Precautions (5.4)*].
- pregnant patients at term (38 weeks to 42 weeks gestation), during labor and delivery, or when the onset of labor is imminent and in pediatric patients younger than 1 month of age because of the possibility of hemolytic anemia due to immature erythrocyte enzyme systems (glutathione instability) [see *Warnings and Precautions (5.5)* and *Use in Specific Populations (8.1 and 8.4)*].

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypersensitivity Reactions**

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving nitrofurantoin oral suspension [see *Adverse Reactions (6)*]. If signs and symptoms of a hypersensitivity reaction occurs, immediately discontinue nitrofurantoin oral suspension and initiate appropriate medications and/or supportive care. Nitrofurantoin oral suspension is contraindicated in patients with known hypersensitivity to nitrofurantoin.

### **5.2 Pulmonary Reactions**

Acute, subacute, or chronic pulmonary reactions have been reported in patients treated with nitrofurantoin oral suspension. If these reactions occur, discontinue nitrofurantoin oral suspension and take appropriate measures. Reports have cited pulmonary reactions as a contributing cause of death.

Chronic pulmonary reactions (diffuse interstitial pneumonitis or pulmonary fibrosis, or both) can develop insidiously. These reactions occur generally in patients receiving therapy for six months or longer. Close monitoring of the pulmonary condition of patients receiving long-term therapy is warranted and requires that the benefits of therapy be weighed against potential risks [see *Adverse Reactions (6)*].

### **5.3 Hepatotoxicity**

Hepatic reactions, including hepatitis, cholestatic jaundice, chronic active hepatitis, and hepatic necrosis, has occurred. Fatalities have been reported. The onset of chronic active hepatitis may be insidious. Monitor patients periodically for changes in biochemical tests that would indicate liver injury. If hepatitis occurs, discontinue nitrofurantoin oral suspension immediately, and take appropriate measures.

### **5.4 Neuropathy**

Peripheral neuropathy, which may become severe or irreversible, has occurred. Fatalities have been reported. Conditions such as renal impairment (creatinine clearance under 60 mL per minute or clinically significant elevated serum creatinine), anemia, diabetes mellitus, electrolyte imbalance, vitamin B deficiency, and debilitating disease may enhance the occurrence of peripheral neuropathy. Monitor patients receiving long-term therapy periodically for changes in renal function.

Optic neuritis has been reported with nitrofurantoin formulations [see *Adverse Reactions (6)*].

### **5.5 Hemolytic Anemia**

Cases of hemolytic anemia of the primaquine-sensitivity type have been induced by nitrofurantoin. Hemolysis appears to be linked to a glucose-6-phosphate dehydrogenase deficiency in the red blood cells of the affected patients. This deficiency is found in 10 percent of Blacks and a small percentage of ethnic groups of Mediterranean and Near-Eastern origin. If hemolysis occurs, discontinue nitrofurantoin oral suspension immediately; hemolysis ceases when the drug is withdrawn.

### **5.6 *Clostridioides difficile*-associated Diarrhea**

*Clostridioides difficile*-associated Diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including nitrofurantoin, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antibacterial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued.

Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and institute surgical evaluation as clinically indicated.

### **5.7 Persistence or Reappearance of Bacteriuria**

Nitrofurantoin lacks the broader tissue distribution of other therapeutic agents approved for urinary tract infections. Consequently, many patients who are treated with nitrofurantoin oral suspension are predisposed to persistence or reappearance of bacteriuria. If persistence or reappearance of bacteriuria occurs with symptoms of urinary tract infection, after treatment with nitrofurantoin oral suspension, other therapeutic agents with broader tissue distribution should be selected. In considering the use of nitrofurantoin oral suspension, lower eradication rates should be balanced against the increased potential for systemic toxicity and for the development of antibacterial resistance when agents with broader tissue distribution are utilized.

## **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are discussed in more detail in other sections of the labeling:

- Hypersensitivity Reactions [see *Warnings and Precautions (5.1)*]
- Pulmonary Reactions [see *Warnings and Precautions (5.2)*]
- Hepatotoxicity [see *Warnings and Precautions (5.3)*]
- Neuropathy [see *Warnings and Precautions (5.4)*]

- Hemolytic anemia [see *Warnings and Precautions (5.5)*]
- *Clostridioides difficile*-associated Diarrhea [see *Warnings and Precautions (5.6)*]
- Persistence or Reappearance of Bacteriuria [see *Warnings and Precautions (5.7)*]

The following adverse reactions associated with the use of nitrofurantoin formulations, including, nitrofurantoin oral suspension were identified in clinical studies or post-marketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Respiratory:* chronic, subacute, or acute pulmonary hypersensitivity reactions have occurred.

Chronic pulmonary reactions have occurred generally in patients who have received continuous treatment for six months or longer. Malaise, dyspnea on exertion, cough, and altered pulmonary function are common manifestations which can occur insidiously. Radiologic and histologic findings of diffuse interstitial pneumonitis or fibrosis, or both, are also common manifestations of the chronic pulmonary reaction. Fever is prominent.

The severity of chronic pulmonary reactions and their degrees of resolution appear to be related to the duration of therapy after the first clinical signs appear. Pulmonary function may be impaired permanently, even after cessation of therapy. The risk is greater when chronic pulmonary reactions are not recognized early.

In subacute pulmonary reactions, fever and eosinophilia occur less often than in the acute form. Upon cessation of therapy, recovery may require several months. If the symptoms are not recognized as being drug-related and nitrofurantoin therapy is not stopped, the symptoms may become more severe.

Acute pulmonary reactions are commonly manifested by fever, chills, cough, chest pain, dyspnea, pulmonary infiltration with consolidation of pleural effusion on x-ray, and eosinophilia. Acute reactions usually occur within the first week of treatment and are reversible with cessation of therapy. Resolution often is dramatic.

Changes in EKG (e.g., non-specific ST/T wave changes, bundle branch block) have been reported in association with pulmonary reactions.

Cyanosis has been reported.

*Hepatic:* Hepatic reactions, including hepatitis, cholestatic jaundice, chronic active hepatitis, and hepatic necrosis, have occurred.

*Neurologic:* Peripheral neuropathy, which may become severe or irreversible, has occurred. Fatalities have been reported. Conditions such as renal impairment (creatinine clearance under 60 mL per minute or clinically significant elevated serum creatinine), anemia, diabetes mellitus, electrolyte imbalance, vitamin B deficiency, and debilitating diseases may increase the possibility of peripheral neuropathy.

Asthenia, vertigo, nystagmus, dizziness, headache, and drowsiness also have been reported with the use of nitrofurantoin.

Benign intracranial hypertension (pseudotumor cerebri), confusion, depression, optic neuritis, and psychotic reactions have been reported. Bulging fontanels, as a sign of benign intracranial hypertension in infants, have been reported rarely.

*Dermatologic:* Exfoliative dermatitis and erythema multiforme (including Stevens-Johnson

syndrome) have been reported. Alopecia also has been reported.

*Allergic:* A lupus-like syndrome associated with pulmonary reactions to nitrofurantoin has been reported. Also, angioedema; maculopapular, erythematous, or eczematous eruptions; pruritus; urticaria; anaphylaxis; arthralgia; myalgia; drug fever; and vasculitis (sometimes associated with pulmonary reactions) have been reported. Hypersensitivity reactions present the most frequent spontaneously-reported adverse reactions in worldwide post marketing experience with nitrofurantoin formulations, including nitrofurantoin oral suspension.

*Gastrointestinal:* Nausea, emesis, and anorexia occur most often. Abdominal pain and diarrhea are less common gastrointestinal reactions. These dose-related reactions can be minimized by reduction of dosage. Sialadenitis and pancreatitis have been reported. There have been sporadic reports of pseudomembranous colitis with the use of nitrofurantoin. The onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment.

*Hematologic:* Cyanosis secondary to methemoglobinemia has been reported.

*Miscellaneous:* As with other antibacterial agents, superinfections caused by resistant organisms, e.g., *Pseudomonas* species or *Candida* species, can occur. There are sporadic reports of *Clostridioides difficile* superinfections, or pseudomembranous colitis, with the use of nitrofurantoin, including nitrofurantoin oral suspension.

#### *Laboratory Adverse Reactions*

The following laboratory adverse reactions have been reported with the use of nitrofurantoin formulations, including nitrofurantoin oral suspension; increased AST (SGOT), increased ALT (SGPT), decreased hemoglobin, increased serum phosphorus, eosinophilia, glucose-6-phosphate dehydrogenase deficiency anemia, agranulocytosis, leukopenia, granulocytopenia, hemolytic anemia, thrombocytopenia, megaloblastic anemia. In most cases, these hematologic abnormalities resolved following cessation of therapy. Aplastic anemia has been reported.

## **7 DRUG INTERACTIONS**

### **7.1 Antacids**

Antacids containing magnesium trisilicate, when administered concomitantly with nitrofurantoin, reduce both the rate and extent of absorption. The mechanism for this interaction probably is adsorption of nitrofurantoin onto the surface of magnesium trisilicate. If coadministration of nitrofurantoin with antacids containing magnesium trisilicate cannot be avoided, monitor for lack of efficacy [see *Clinical Pharmacology (12.3)*].

### **7.2 Uricosuric Drugs**

Uricosuric drugs, such as probenecid and sulfinpyrazone, can inhibit renal tubular secretion of nitrofurantoin. The resulting increase in nitrofurantoin serum levels may increase toxicity, and the decreased urinary levels could lessen its efficacy as a urinary tract antibacterial. Therefore, monitor for nitrofurantoin adverse reactions when co-administering nitrofurantoin with uricosuric drugs.



### 7.3 Drug Interference with Laboratory Tests

The presence of nitrofurantoin can cause a false-positive reaction for glucose in the urine when using Benedict's or Fehling's copper reduction reaction to determine the amount of reducing substances like glucose in the urine. Use glucose tests based on enzymatic glucose oxidase reactions to detect glucose in the urine.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Nitrofurantoin is contraindicated in pregnant women at term (38 weeks to 42 weeks gestation), during labor and delivery, or when the onset of labor is imminent because of the possibility of hemolytic anemia in the infant (*see Clinical Considerations*).

Published epidemiological studies, including cohort studies and case control studies, have reported inconsistent findings related to nitrofurantoin use during the first trimester and risk of major birth defects. These studies cannot definitively establish the presence or absence of risk due to several methodological limitations. The limited number of epidemiological studies available have not identified drug-associated risks of major birth defects, miscarriage or other adverse maternal or fetal outcomes.

In animal reproduction studies, no fetotoxicity was observed when nitrofurantoin was administered orally to pregnant rats and rabbits, during organogenesis, at up to 6 times the recommended human dose (*see Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

#### Clinical Considerations

##### *Fetal/Neonatal Adverse Reactions*

Nitrofurantoin is not recommended in pregnant women at term (38 weeks to 42 weeks gestation), during labor and delivery, or when the onset of labor is imminent because of the possibility of hemolytic anemia due to immature erythrocyte enzyme systems (glutathione instability) [*see Contraindications (4) and Warnings and Precautions (5.4)*].

#### Data

##### *Animal Data*

Several reproduction studies have been performed in rabbits and rats at doses up to six times the human dose and have revealed no harm to the fetus due to nitrofurantoin. In a single published study conducted in mice at 68 times the human dose (based on mg/kg administered to the dam), growth retardation and a low incidence of minor and common malformations were observed. However, at 25 times the human dose, fetal malformations were not observed.

Nitrofurantoin has been shown in one published transplacental carcinogenicity study to

induce lung papillary adenomas in the F1 generation mice at doses 19 times the human dose on a mg/kg basis. The relationship of this finding to potential human carcinogenesis is presently unknown.

## **8.2 Lactation**

### Risk Summary

Nitrofurantoin is present in human breast milk following oral administration. There are insufficient data on the effect of nitrofurantoin on milk production.

Infants less than one month of age or who have glucose-6-phosphate dehydrogenase (G6PD) deficiency are at risk for hemolytic anemia from nitrofurantoin exposure. Therefore, breastfeeding is not recommended in these infants.

In a breastfed infant over the age of one month and with normal G6PD, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for nitrofurantoin and any potential adverse effects on the breastfed infant from nitrofurantoin or from the underlying maternal condition. Monitor these infants for vomiting, diarrhea, and rash.

## **8.3 Females and Males of Reproductive Potential**

### Infertility

#### *Male*

Based on findings from a fertility study conducted in 36 healthy male volunteers and an animal fertility study, nitrofurantoin may reversibly decrease spermatogenesis [see *Nonclinical Toxicology (13.1)*]. Nitrofurantoin doses of 10 mg/kg/day or greater in healthy human males may produce a slight to moderate spermatogenic arrest with a decrease in sperm count. The effect on fertility is unknown.

## **8.4 Pediatric Use**

Nitrofurantoin oral suspension is contraindicated in pediatric patients younger than 1 month of age because of the possibility of hemolytic anemia due to immature erythrocyte enzyme systems (glutathione instability). [see *Contraindications (4) and Warnings and Precautions (5.4)*].


The safety and effectiveness of nitrofurantoin oral suspension in pediatric patients aged 1 month of age and older for the treatment of urinary tract infections have been established.

## **10 OVERDOSAGE**

Incidents of acute overdosage of nitrofurantoin oral suspension have resulted in symptoms such as vomiting. Induction of emesis is recommended. There is no specific antidote, but a high fluid intake should be maintained to promote urinary excretion of the drug. Nitrofurantoin is dialyzable.

## **11 DESCRIPTION**

Nitrofurantoin Oral Suspension, USP contains nitrofurantoin, a synthetic nitrofuran antibacterial agent specific for urinary tract infections. The chemical name is 1-[[[(5-nitro-2-furanyl)methylene]amino]-2,4imidazolidinedione monohydrate. The molecular formula is  $C_8H_6N_4O_5 \cdot H_2O$  and the molecular weight is 256.17. The structural formula is:



Nitrofurantoin is a stable, yellow, crystalline compound.

Nitrofurantoin oral suspension is available as an opaque, yellow liquid suspension for oral administration containing 25 mg/5 ml of nitrofurantoin. The suspension also contains the following inactive ingredients: citric acid anhydrous, fruit flavor, glycerin, magnesium aluminum silicate, methyl paraben, propyl paraben, purified water, sodium carboxymethyl cellulose, sorbitol solution, tri-sodium citrate dihydrate powder, xanthan gum.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Nitrofurantoin is an antibacterial drug [see *Microbiology (12.4)*].

### 12.2 Pharmacodynamics

Pharmacodynamic effects of nitrofurantoin oral suspension are unknown.

### 12.3 Pharmacokinetics

#### Absorption

Orally administered nitrofurantoin oral suspension is readily absorbed. Blood concentrations at therapeutic dosage are usually low.

#### Effect of Food

The presence of food or agents delaying gastric emptying can increase the bioavailability of nitrofurantoin oral suspension.

#### Distribution

Nitrofurantoin is highly soluble in urine. Nitrofurantoin lacks the broader tissue

distribution of other therapeutic agents approved for urinary tract infections.

### Elimination

#### *Excretion*

Nitrofurantoin is rapidly excreted in urine, to which it may impart a brown color.

Following a dose regimen of 100 mg four times a day for 7 days, average urinary drug recoveries (0 hour to 24 hours) on Day 1 and Day 7 were 42.7% and 43.6%.

Nitrofurantoin is dialyzable.

### Specific Populations

The pharmacokinetics of nitrofurantoin oral suspension are unknown for the geriatric patients, pediatric patients, patients with hepatic impairment, or pregnant women. Differences between male and female patients, and racial or ethnic groups are unknown.

#### *Patients with Renal Impairment*

The reported urinary recoveries of nitrofurantoin after a single 100 mg oral dose and 100 mg four times a day in patients with varying renal function showed approximate linear relationship between the amount of nitrofurantoin recovered and creatinine clearance.

### Drug Interaction Studies

#### *Antacids such as magnesium trisilicate*

In a cross-over study, the *in vivo* absorption characteristics of nitrofurantoin and nitrofurantoin-magnesium trisilicate combination were evaluated in 6 healthy males. The administration of 5 g magnesium trisilicate with 100 mg nitrofurantoin reduced the rate and extent of nitrofurantoin excretion reflecting decrease in both rate and extent of its absorption [see *Drug Interaction (7.1)*].

#### *Propantheline*

In a crossover study, the *in vivo* absorption characteristics of nitrofurantoin and propantheline (an anticholinergic agent) combination were evaluated in six subjects. Administration of 30 mg of propantheline 45 min prior to nitrofurantoin 100 mg administration resulted in a statistically significant increase in nitrofurantoin excretion, however, the clinical significance of this effect is unknown.

## **12.4 Microbiology**

### Mechanism of Action

Nitrofurantoin is reduced by a wide range of enzymes including bacterial flavoproteins to reactive intermediates which are damaging to macromolecules such as DNA and proteins.

### Resistance

Nitrofurantoin is not active against most strains of *Proteus* species or *Serratia* species. It has no activity against *Pseudomonas* species. Although cross-resistance with other antibacterials may occur, cross resistance with sulfonamides has not been observed.

### Interaction with Other Antibacterials

Antagonism has been demonstrated *in vitro* between nitrofurantoin and quinolones.

### Antibacterial Activity

Nitrofurantoin has been shown to be active against most of the following bacteria both *in vitro* and in clinical infections:

#### **Aerobic bacteria**

##### **Gram-positive bacteria**

*Staphylococcus aureus*

*Enterococcus* species

##### **Gram-negative bacteria**

*Escherichia coli*

*Enterobacter* species

*Klebsiella* species

The following *in vitro* data are available, but their clinical significance is unknown. Nitrofurantoin exhibits *in vitro* activity against the following bacteria; however, the safety and efficacy of nitrofurantoin in treating clinical infections due to these bacteria have not been established in adequate and well-controlled clinical trials.

#### **Aerobic bacteria**

##### **Gram-positive bacteria**

Coagulase-negative staphylococci (including *Staphylococcus epidermidis* and *Staphylococcus saprophyticus*)

*Streptococcus agalactiae*

Viridans group streptococci

##### **Gram-negative bacteria**

*Citrobacter koseri*

*Citrobacter freundii*

*Klebsiella oxytoca*

### Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria, and associated test methods and quality control standards recognized by FDA for this drug, please see: <http://www.fda.gov/STIC>.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Nitrofurantoin was not carcinogenic when fed to female Holtzman rats for 44.5 weeks

or to female Sprague-Dawley rats for 75 weeks. Two chronic rodent bioassays utilizing male and female Sprague-Dawley rats and two chronic bioassays in Swiss mice and in BDF<sub>1</sub> mice revealed no evidence of carcinogenicity.

Nitrofurantoin presented evidence of carcinogenic activity in female B6C3F<sub>1</sub> mice as shown by increased incidences of tubular adenomas, benign mixed tumors, and granulosa cell tumors of the ovary. In male F344/N rats, there were increased incidences of uncommon kidney tubular cell neoplasms, osteosarcomas of the bone, and neoplasms of the subcutaneous tissue.

The significance of carcinogenicity findings relative to the therapeutic use of nitrofurantoin in humans is unknown.

### Mutagenesis

Nitrofurantoin has been shown to induce point mutations in certain strains of *Salmonella typhimurium* and forward mutations in L5178Y mouse lymphoma cells. Nitrofurantoin induced increased numbers of sister chromatid exchanges and chromosomal aberrations in Chinese hamster ovary cells but not in human cells in culture. Results of the sex-linked recessive lethal assay in *Drosophila* were negative after administration of nitrofurantoin by feeding or by injection. Nitrofurantoin did not induce heritable mutation in the rodent models examined.

The significance of mutagenicity findings relative to the therapeutic use of nitrofurantoin in humans is unknown.

### Impairment of Fertility

The administration of high doses of nitrofurantoin to rats causes temporary spermatogenic arrest; this is reversible on discontinuing the drug.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### How Supplied

Nitrofurantoin oral suspension, USP (25 mg/5 mL) is available in:

**NDC 70377-118-31** PET amber bottle of 230 mL

### Storage and Handling

Avoid exposure to strong light which may darken the drug. It is stable when stored between 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature]. Protect from freezing. Shake vigorously. Dispense in tight, light-resistant, plastic (PET) or glass container.

Use within 30 days.

## **17 PATIENT COUNSELING INFORMATION**

### Administration Instructions

Instruct the patients to:

- complete the full course of therapy; however, advise the patients or their caregiver to contact their physician if any unusual symptoms occur during therapy.

- take daily doses of nitrofurantoin oral suspension with food, as food will further enhance tolerance and improve the absorption of the drug.
- shake nitrofurantoin oral suspension vigorously before use each time.
- use an oral dosing syringe to correctly measure the prescribed amount of nitrofurantoin oral suspension. Inform patients or caregivers that an appropriate size oral dosing syringe with graduations that align with the volume prescribed may be obtained from their pharmacy.

### Serious Allergic Reactions

Advise patients that allergic reactions, including serious allergic reactions, could occur with nitrofurantoin oral suspension and that serious reactions require immediate treatment. Advise patients to discontinue nitrofurantoin oral suspension at the first sign of allergic reactions [see *Warnings and Precautions (5.1)*].

### Diarrhea

Advise patients that diarrhea is a common problem caused by antibacterial drugs, including nitrofurantoin oral suspension which usually ends when the antibacterial drug is discontinued. Sometimes frequent watery and bloody diarrhea (with or without stomach cramps and fever) may occur even as late as two or more months after having taken the last dose of the antibacterial drug. If this occurs, advise patients to contact their physician as soon as possible [see *Warnings and Precautions (5.6)*].

### Drug Interactions with Antacids

Advise patients not to use antacid preparations containing magnesium trisilicate while taking nitrofurantoin oral suspension [see *Drug Interactions (7.1)*].

### Lactation

Advise women not to breastfeed infants who are less than one month of age or who have G6PD deficiency during treatment with nitrofurantoin oral suspension [see *Use in Specific Populations (8.2)*].

### Antibacterial Resistance

Patients should be counseled that antibacterial drugs including nitrofurantoin oral suspension should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When nitrofurantoin oral suspension is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of the therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by nitrofurantoin oral suspension or other antibacterial drugs in the future.

### Change in Urine Color

Inform patients that nitrofurantoin oral suspension may impart a brown color to their urine [see *Clinical Pharmacology (12.3)*].

Manufactured for:

**Biacon Pharma Inc.,**

Iselin, New Jersey, 08830-3009 USA

Manufactured by:  
Carnegie Pharmaceuticals LLC  
Delran, NJ 08075, USA  
Revised: 10/2024

## PRINCIPAL DISPLAY PANEL

NDC 70377-**118**-31  
Nitrofurantoin Oral Suspension USP, 25 mg/5 mL  
**230 mL**  
**Rx only**

NDC 70377-**118**-31

**Nitrofurantoin Oral Suspension, USP**

**25 mg/5 mL**

Urinary Tract Antibacterial

230 mL

 **Rx only**

SHAKE VIGOROUSLY.  
Dispense in tight, light-resistant, plastic (PET) or glass container.

**For Patients:** Date of first opening \_\_\_/\_\_\_/\_\_\_.  
Discard unused portion 30 days after first opening.

**Recommended Dosage:** See Prescribing Information.  
Each 5 mL contains 25 mg of nitrofurantoin.

**Store between 20°C-25°C (68°F-77°F); excursions permitted to 15°C - 30°C (59°F - 86°F). [See USP].**

**Avoid exposure to strong light which may darken the drug.**

**Protect from freezing.**

**Keep out of reach of children.**

Manufactured for: **Biocon Pharma Inc.**,  
Iselin, New Jersey, 08830-3009 USA


Manufactured by: Carnegie Pharmaceuticals LLC  
Delran, NJ 08075, USA

Revised: 10/2024  
BPCM0272/01

9  
7037711831  
N3

Unvarnished Area  
0.75" x 2.0"





# NITROFURANTOIN

nitrofurantoin suspension

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70377-118
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>NITROFURANTOIN</b> (UNII: 927AH8112L) (NITROFURANTOIN - UNII:927AH8112L)	NITROFURANTOIN	25 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>MAGNESIUM ALUMINUM SILICATE</b> (UNII: 6M3P64V0NC)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KOOR)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70377-118-31	1 in 1 CARTON	11/15/2024	
1		230 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA218346	11/15/2024	

**Labeler** - Biocon Pharma Inc (080000063)**Registrant** - Biocon Pharma Limited (871412155)**Establishment**

Name	Address	ID/FEI	Business Operations
Carnegie Pharmaceuticals LLC		079839141	manufacture(70377-118)

