

STAFAC 20- virginiamycin powder
Phibro Animal Health

Stafac® 20
(virginiamycin)
TYPE A MEDICATED ARTICLE

Active Drug Ingredient

Virginiamycin 4.4%

(Contains 20 g virginiamycin activity per lb)

CAUTION:

Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Ingredients:

Processed grain by-products, roughage products, calcium carbonate, carboxymethylcellulose, mineral oil

(20 g virginiamycin activity per lb)

For use in swine and poultry feeds as specified below

Directions for Use

Mixing Directions

Stafac 20 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of Stafac 20 required from the table below. Dilute Stafac 20 with a portion of one of the feed ingredients (enough to make a quantity equal to 10 lb per ton) before final mixing.

Swine	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 20 per ton of Type C Medicated Feed (complete feed)	Amount of virginiamycin activity provided to animal per lb of body weight per day
Treatment of swine dysentery in nonbreeding swine weighing over 120 lb	100 g for 2 weeks	5.00 lb	2.27 mg
Treatment and control of swine dysentery in swine weighing up to 120 lb	100 g for 2 weeks followed by 50g	5.00 lb 2.50 lb	2.27 mg 1.14 mg

Aid in control of swine dysentery in swine weighing up to 120 lb For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred	25 g	1.25 lb	0.57 mg
CAUTION: Not for use in breeding swine over 120 lbs.			
Poultry	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 20 per ton of Type C Medicated Feed (complete feed)	
Prevention of necrotic enteritis caused by <i>Clostridium perfringens</i> susceptible to virginiamycin Not for Use in Laying Chickens	20 g	1.00 lb	

Feed Continuously as Sole Ration

Store at Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Close Container After Use

For further manufacture only.

Follow Directions on Back Panel

Stafac is a registered trademark of Phibro Animal Health Corporation

Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.6 kg)

Approved by FDA under NADA # 091-467

9907000

101-9007-12

Stafac 20

Stafac 20



Stafac[®] 20 (virginiamycin)

TYPE A
MEDICATED ARTICLE

(20 g virginiamycin activity per lb)
For use in swine and poultry feeds as specified below

Directions for Use

Mixing Directions

Stafac 20 Type A Medicated Article must be thoroughly mixed in the feed to ensure even distribution. Refer to the user specific inserts of Stafac 20 for all other feed instructions. Please consult the user specific insert of use of the feed ingredients through to make a quantity equal to 10 lb per ton (2000 lb) of feed.

Name	Required level of virginiamycin activity (g per ton of Type 2 Medicated Feed (complete feed))	Required amount of Stafac 20 (g per ton of Type 2 Medicated Feed (complete feed))	Amount of virginiamycin activity (g) per lb of dry weight per ton
Treatment of swine (quantity in complete feed weighing up to 100 lb)	20 g	1.00 lb	1.00 mg
Treatment and control of swine (quantity in complete feed weighing up to 100 lb)	100 g	5.00 lb	1.00 mg
Add to control of swine (quantity in complete feed weighing up to 100 lb)	20 g	1.00 lb	1.00 mg

CAUTION: Not for use in feeding swine to or 100 lb.

Name	Required level of virginiamycin activity (g per ton of Type 2 Medicated Feed (complete feed))	Required amount of Stafac 20 (g per ton of Type 2 Medicated Feed (complete feed))	Amount of virginiamycin activity (g) per lb of dry weight per ton
Treatment of swine (quantity in complete feed weighing up to 100 lb)	20 g	1.00 lb	1.00 mg

For use in swine or swine with a history of severe diarrhoea, reduce the amount of feed to 100 lb.

06/2020
01-400-11A

Developed by
Phibro Animal Health Corporation
Teaneck, NJ 07666

Stafac[®] 20
(virginiamycin)
Phibro

Stafac[®] 20 (virginiamycin)

TYPE A
MEDICATED ARTICLE

Active Drug Ingredient

Virginiamycin 4.4%
(Contains 20 g virginiamycin activity per lb)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Ingredients:

Processed grain by-products, roughage products, calcium carbonate, carboxymethylcellulose, mineral oil

For further manufacture only.

Follow Directions on Back Panel

Stafac is a registered trademark of Phibro Animal Health Corporation
Phibro Animal Health Corporation, Teaneck, NJ 07666

NADA 401-402, Approved by FDA
09/2003
101-1002-11A

Net Weight 50 lb (22.6 kg)

Phibro
ANIMAL HEALTH CORPORATION

Phibro
Stafac[®] 20
(virginiamycin)

STAFAC 20 virginiamycin powder

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66104-1220
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VIRGINIAMYCIN (UNII: C49WS9N75L) (VIRGINIAMYCIN - UNII:C49WS9N75L)	VIRGINIAMYCIN	20 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
RICE BRAN (UNII: R60QEP13IC)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1 NDC:66104-1220-0 22.6 kg in 1 BAG

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA091467	03/24/2010	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 10/2025

Phibro Animal Health