

## **NIGHT TIME COLD AND COUGH CHILDRENS- diphenhydramine hcl, phenylephrine hcl liquid**

### **Best Choice (Valu Merchandisers Company)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredients (in each 5 mL)**

Diphenhydramine HCl 6.25 mg

Phenylephrine HCl 2.5 mg

### **Purposes**

Antihistamine/Cough suppressant

Nasal decongestant

### **Uses**

temporarily relieves:

- sneezing
- runny nose
- itchy nose or throat
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

### **Warnings**

#### **Do not use**

- in a child under 4 years of age
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other product containing diphenhydramine, even one used on skin
- for the purpose of making your child sleepy

#### **Ask a doctor before use if the child has**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma
- a breathing problem such as chronic bronchitis

#### **Ask a doctor or pharmacist before use if the child is**

taking sedatives or tranquilizers

### When using this product

- **do not exceed recommended dosage**
- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

### Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache.

These could be signs of a serious condition.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

- Do not give more than 6 doses in 24 hours unless directed by a doctor.
- mL = milliliter
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	do not use unless directed by a doctor
<b>children under 4 years of age</b>	<b>do not use</b>

### Other information

- **each 5 mL contains:** sodium 3 mg
- store between 20-25° (68-77°F). Do not refrigerate.
- protect from light

### Inactive ingredients

acesulfame potassium, citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### Principal Display Panel

FOR AGES 6 to 11 YEARS

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TRIAMINIC® NIGHTTIME COLD & COUGH SYRUP

**Children's**

**Night Time**

**Cold and Cough Syrup**

Diphenhydramine HCl

- Antihistamine/ Cough suppressant

Phenylephrine HCl

- Nasal decongestant

GRAPE FLAVOR

Dosing Cup Included

FL OZ (mL)

\*This product is not manufactured or distributed by Novartis distributors of Children's Triaminic® Night Time Cold & Cough.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

PROUDLY DISTRIBUTED BY:

VALU MERCHANDISERS, CO.

5000 KANSAS AVE

KANSAS CITY, KS 66106

[www.bestchoicebrand.com](http://www.bestchoicebrand.com)

**Package Label**

**Drug Facts (continued)**

**Directions**

- do not give more than 6 doses in 24 hours unless directed by a doctor
- ml = milliliter
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	do not use unless directed by a doctor
children under 4 years of age	do not use

**Other information**

- each 5 mL contains: sodium 3 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

**Inactive ingredients** acesulfame potassium, citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by Noveris, distributor of Children's Triaminic® Night Time Cold & Cough.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.



Dosing Cup Included



PROUDLY DISTRIBUTED BY:  
WALLU MERCHANTISERS, CO.  
5000 KANSAS AVE  
KANSAS CITY, KS 66106



FOR AGES 6 TO 11 YEARS

**Best Choice®**

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TRIAMINIC® NIGHT TIME COLD & COUGH\*

**Children's NIGHT TIME COLD & COUGH SYRUP**

**Diphenhydramine HCl**  
• Antihistamine/Cough Suppressant

**Phenylephrine HCl**  
• Nasal Decongestant

**GRAPE FLAVOR**

4 FL OZ (118 mL)

FOR AGES 6 TO 11 YEARS

**Best Choice®**

COMPARE TO THE ACTIVE INGREDIENTS IN CHILDREN'S TRIAMINIC® NIGHT TIME COLD & COUGH\*

**Children's NIGHT TIME COLD & COUGH SYRUP**

**Diphenhydramine HCl**  
• Antihistamine/Cough Suppressant

**Phenylephrine HCl**  
• Nasal Decongestant

**GRAPE FLAVOR**

4 FL OZ (118 mL)

**Drug Facts**

**Active ingredients (in each 5 mL)**

Diphenhydramine HCl 6.25 mg.....Antihistamine/Cough suppressant  
Phenylephrine HCl 2.5 mg.....Nasal decongestant

**Purposes**

**Uses**

- temporarily relieves
- sneezing
- runny nose
- itchy nose or throat
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritations as may occur with a cold

**Warnings**

**Do not use**

- in a child under 4 years of age
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other product containing diphenhydramine, even one used on skin
- for the purpose of making your child sleepy

**Ask a doctor before use if the child has**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma
- a breathing problem such as chronic bronchitis

**Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.**

**When using this product**

- do not exceed recommended dosage
- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Best Choice Children's NightTime Cold & Cough Syrup Grape Liquid**

**NIGHT TIME COLD AND COUGH CHILDRENS**

diphenhydramine hcl, phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63941-079
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)</b>	DIPHENHYDRAMINE HYDROCHLORIDE	6.25 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)</b>	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MALTITOL (UNII: D65DG142WK)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

**Product Characteristics**

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-079-04	1 in 1 BOX	10/31/2014	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	10/31/2014	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)

Revised: 7/2016

Best Choice (Valu Merchandisers Company)