

**BACTERIOSTATIC WATER- water injection, solution**  
**Medical Purchasing Solutions, LLC**

-----  
**BACTERIOSTATIC WATER Rx only**

**for Injection, USP**

Multiple-dose Plastic Vial

<p><b>WARNING:</b> <b>NOT FOR USE IN NEONATES.</b></p>
--

***DESCRIPTION***

The following preparation is designed solely for parenteral use only after addition of drugs that require dilution or must be dissolved in an aqueous vehicle prior to injection.

Bacteriostatic Water for Injection, USP is a sterile, nonpyrogenic preparation of water for injection containing 0.9% (9 mg/mL) of benzyl alcohol added as a bacteriostatic preservative. It is supplied in a multiple-dose container from which repeated withdrawals may be made to dilute or dissolve drugs for injection. The pH is 5.7 (4.5 to 7.0).

Water for Injection, USP is chemically designated H<sub>2</sub>O.

The semi-rigid vial is fabricated from a specially formulated polyolefin. It is a copolymer of ethylene and propylene. The safety of the plastic has been confirmed by tests in animals according to USP biological standards for plastic containers. The container requires no vapor barrier to maintain the proper labeled volume.

***CLINICAL PHARMACOLOGY***

Water is an essential constituent of all body tissues and accounts for approximately 70% of total body weight. Average normal adult daily requirement ranges from two to three liters (1.0 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water distribution depends primarily on the concentration of electrolytes in the body compartments and sodium (Na<sup>+</sup>) plays a major role in maintaining physiologic equilibrium.

The small volume of fluid provided by Bacteriostatic Water for Injection, USP when used only as a pharmaceutical aid for diluting or dissolving drugs for parenteral injection, is unlikely to exert a significant effect on fluid balance except possibly in very small infants.

***INDICATIONS AND USAGE***

This parenteral preparation is indicated only for diluting or dissolving drugs for intravenous, intramuscular or subcutaneous injection, according to instructions of the manufacturer of the drug to be administered.

## **CONTRAINDICATIONS**

Due to the potential toxicity of benzyl alcohol in neonates, solutions containing benzyl alcohol must not be used in this patient population.

Parenteral preparations with benzyl alcohol should not be used for fluid replacement.

Parenteral preparations containing benzyl alcohol should not be used in epidural or spinal anesthesia procedures.

Bacteriostatic Water for Injection, USP must be made approximately isotonic prior to use.

## **WARNINGS**

Benzyl alcohol, a preservative in Bacteriostatic Water for Injection, USP has been associated with toxicity in neonates. Data are unavailable on the toxicity of other preservatives in this age group. Where water is required for preparing or diluting medications for use in neonates, only preservative-free Sterile Water for Injection should be used.

Intravenous administration of Bacteriostatic Water for Injection without a solute may result in hemolysis.

## **PRECAUTIONS**

Do not use Bacteriostatic Water for Injection, USP for intravenous injection unless the osmolar concentration of additives results in an approximate isotonic admixture.

Consult the manufacturer's instructions for choice of vehicle, appropriate dilution or volume for dissolving the drugs to be injected, including the route and rate of injection.

Inspect reconstituted (diluted or dissolved) drugs for clarity (if soluble) and freedom from unexpected precipitation or discoloration prior to administration.

*Pregnancy Category C.* Animal reproduction studies have not been conducted with Bacteriostatic Water for Injection. It is also not known whether Bacteriostatic Water for Injection containing additives can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Bacteriostatic Water for Injection, USP containing additives should be given to a pregnant woman only if clearly needed.

## **Pediatric Use**

The safety and effectiveness of Bacteriostatic Water for Injection, USP have not been established in pediatric patients. Due to the potential for toxicity, solutions containing benzyl alcohol should not be used in neonates.

## **Drug Interactions**

Some drugs for injection may be incompatible in a given vehicle, or when combined in the same vehicle or in a vehicle containing benzyl alcohol. Consult with pharmacist, if available.

Use aseptic technique for single or multiple entry and withdrawal from all containers.

When diluting or dissolving drugs, mix thoroughly and use promptly.

Do not store reconstituted solutions of drugs for injection unless otherwise directed by the manufacturer of the solute.

Do not use unless the solution is clear and seal intact.

### **ADVERSE REACTIONS**

Reactions which may occur because of this solution, added drugs or the technique of reconstitution or administration include febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasation.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate countermeasures, and if possible, retrieve and save the remainder of the unused vehicle for examination.

Although adverse reactions to intravenous, intramuscular or subcutaneous injection of 0.9% benzyl alcohol are not known to occur in man, experimental studies of small volume parenteral preparations containing 0.9% benzyl alcohol in several species of animals have indicated that an estimated intravenous dose up to 30 mL may be safely given to an adult without toxic effects. Administration of an estimated 9 mL to a 6 kg infant or neonate is potentially capable of producing blood pressure changes.

### **OVERDOSAGE**

Use only as a diluent or solvent. This parenteral preparation is unlikely to pose a threat of fluid overload except possibly in very small infants. In the event these should occur, re-evaluate the patient and institute appropriate corrective measures. See WARNINGS, PRECAUTIONS and ADVERSE REACTIONS.

### **DOSAGE AND ADMINISTRATION**

The volume of the preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration as recommended by the manufacturer.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. See PRECAUTIONS.

### **HOW SUPPLIED**

<b>Unit of Sale</b>	<b>Benzyl Alcohol Preservative Concentration</b>	<b>Each</b>
<b>NDC 0409-3977-03</b> Tray of 25	9 mg/mL	NDC 0409-3977-01 Multiple-dose 30 mL plastic flip-top vial

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Revised: 10/2013

EN-3371

Hospira, Inc., Lake Forest, IL 60045 USA



**PRINCIPAL DISPLAY PANEL - 30 mL Vial Label**

30 mL Multiple-dose

**BACTERIOSTATIC**

**WATER**

for Injection, USP

**NOT FOR USE IN NEONATES**

30 mL Multiple-dose

**BACTERIOSTATIC  
WATER**  
for Injection, USP

**NOT FOR USE IN NEONATES**

***For drug diluent use only.***

0.9% benzyl alcohol added. Sterile,  
nonpyrogenic. Use only if clear  
and seal is intact and undamaged.  
Cleanse stopper with antiseptic.

RL-4365

Hospira, Inc.  
Lake Forest, IL 60045 USA

Rx only  
  
Hospira



**OUTER PACKAGE LABEL**

NDC 71872-7129-1

Rx Only

Bacteriostatic Water for Injection, USP

1 x 30mL Multiple-dose Vial

**FOR DRUG DILUENT USE ONLY**



Rx only



(01) 0 0971872 71291 0  
(17) 000101  
(10) XXXX-XXXXXX  
(21) 7129400000

**Bacteriostatic Water for Injection, USP**  
**30 mL Multiple-dose Vial**

Qty: 1 vial      Lot# XXXX-XXXXXX      Exp: 01/01/1900

FOR DRUG DILUENT USE ONLY.

WARNING: NOT FOR USE IN NEONATES.

Each mL contains 0.9% benzyl alcohol added as a preservative.

Sterile, nonpyrogenic.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

USE ASEPTIC TECHNIQUE. Use only if clear and seal is intact and undamaged. Cleanse stopper with antiseptic.

Manufactured by: Hospira, Inc. Lake Forest, IL 60045 USA  
Manufacturer NDC: 00409-3977-03

Repackaged  
& Distributed By:



Medical Purchasing Solutions  
Scottsdale, AZ 85260  
www.medicalpurchasingsolutions.com

## BACTERIOSTATIC WATER

water injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:71872-7129(NDC:0409-3977)
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS, SUBCUTANEOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>WATER</b> (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	1 mL in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	9 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71872-7129-1	1 in 1 BAG	03/30/2018	
1		30 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018802	04/30/2005	

**Labeler** - Medical Purchasing Solutions, LLC (601458529)

## Establishment

Name	Address	ID/FEI	Business Operations
Medical Purchasing Solutions, LLC		601458529	repack(71872-7129)

Revised: 5/2023

Medical Purchasing Solutions, LLC