

HEPAFLEX- virginiamycin powder
ELANCO US. INC

HepafleX™
(virginiamycin)
Type A Medicated Article

To be mixed in cattle feed

Active Drug Ingredient

Virginiamycin: 50%

(Contains 227 g virginiamycin activity per lb)

CAUTION:

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Inert ingredients:

Calcium carbonate, carboxymethylcellulose, mineral oil

Important: Must be diluted in feed before use

Follow Directions For Use

Approved by FDA under NADA # 140-998

102-8409-1

HepafleX, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

NET WEIGHT: 55 lb (25kg)

Distributed by:

Elanco US Inc.

Indianapolis, IN 46221

Made in Canada

For use in complete feeds for cattle fed in confinement for slaughter
Directions for Use

Cattle fed in confinement for slaughter	Virginiamycin required mg/hd/day	Virginiamycin (g/tons) of complete feed (90% dry matter basis)
Reduction of incidence of liver abscesses	85-240	13.5-16.0

Caution: Not for use in

animals intended for breeding.

Mixing Directions

Type B medicated feed for cattle fed in confinement for slaughter -

Thoroughly mix 4.4 lb of Hepaflex Type A Medicated Article per ton of Type B medicated feed to provide 1000 g/ton virginiamycin.

An intermediate blending step, consistent with the mixing equipment specifications, should be performed to ensure adequate mixing.

Type C medicated feed

The Type B medicated feed must be diluted to a Type C medicated feed before being fed. Prepare a Type B medicated feed as described above. Thoroughly mix the Type B medicated feed to make 1 ton of Type C medicated feed to provide 13.5–16.0g of virginiamycin per ton of complete feed on a 90% dry matter basis using the examples provided in Table 1.

Table 1. Type C medicated feed

lb of a 1,000 g/ton Type B medicated feed per ton of complete feed	Virginiamycin concentration in complete feed 90% dry matter basis (g/ton)
27	13.5
32	16.0

Feed continuously as sole ration

Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Close container after use

SKU:

LOT No.:

DATE OF EXPIRATION:

Distributed by:

Elanco US Inc.

Indianapolis, IN 46221

Made in Canada.

Not For Human Use

Restricted Drug (California) – Use Only as Directed

Hepaflex

Label

Elanco™

Hepaflex™

NET WEIGHT: 55 lb
(25 kg)

(virginiamycin)

Type A Medicated Article
To be mixed in cattle feed

Active Drug Ingredient

Virginiamycin: 50%

(Contains 227 g virginiamycin activity per lb)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Inert ingredients:

Calcium carbonate, carboxymethylcellulose, mineral oil

Important: Must be diluted in feed before use
Follow Directions For Use

Approved by FDA under NADA # 140-998

102-8400-01

Hepaflex, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

For use in complete feeds for cattle fed in confinement for slaughter

Directions for Use

Cattle fed in confinement for slaughter	Virginiamycin required mg/hd/day	Virginiamycin (g/ton) of complete feed (90% dry matter basis)
Reduction of incidence of liver abscesses	85-240	13.5-16.0

Caution: Not for use in animals intended for breeding.

Mixing Directions

Type B medicated feed for cattle fed in confinement for slaughter - Thoroughly mix 4.4 lb of Hepaflex Type A Medicated Article per ton of Type B medicated feed to provide 1000 g/ton virginiamycin. An intermediate blending step, consistent with the mixing equipment specifications, should be performed to ensure adequate mixing.

Type C medicated feed

The Type B medicated feed must be diluted to a Type C medicated feed before being fed. Prepare a Type B medicated feed as described above. Thoroughly mix the Type B medicated feed to make 1 ton of Type C medicated feed to provide 13.5-16.0 g of virginiamycin per ton of complete feed on a 90% dry matter basis using the examples provided in Table 1.

Table 1. Type C medicated feed

lb of a 1,000 g/ton Type B medicated feed per ton of complete feed	Virginiamycin concentration in complete feed 90% dry matter basis (g/ton)
27	13.5
32	16.0

Feed continuously as sole ration

Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Close container after use

Not For Human Use

Restricted Drug (California) - Use Only as Directed

SKU:

LOT No.:

DATE OF EXPIRATION:

Distributed by:

Elanco US Inc.
Indianapolis, IN 46221

Made in Canada.



PMS 001 C



Black

276 x 105 mm

Hepaflex (Elanco) 250G Label 276x105 102-8400-01 12500-010.indd 1

8/8/25 9:17 AM

Elanco™

Hepaflex™

NET WEIGHT: 55 lb
(25 kg)

(virginiamycin)

Type A Medicated Article
To be mixed in cattle feed

Active Drug Ingredient

Virginiamycin: 50%

(Contains 227 g virginiamycin activity per lb)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Inert Ingredients:

Calcium carbonate, carboxymethylcellulose, mineral oil

Important: Must be diluted in feed before use
Follow Directions For Use

Approved by FDA under NADA # 140-998

102-8409-01

Hepaflex, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

For use in complete feeds for cattle fed in confinement for slaughter

Directions for Use

Cattle fed in confinement for slaughter	Virginiamycin required mg/hd/day	Virginiamycin (g/ton) of complete feed (90% dry matter basis)
Reduction of incidence of liver abscesses	85-240	13.5-16.0

Caution: Not for use in animals intended for breeding.

Mixing Directions

Type B medicated feed for cattle fed in confinement for slaughter - Thoroughly mix 4.4 lb of Hepaflex Type A Medicated Article per ton of Type B medicated feed to provide 1000 g/ton virginiamycin. An intermediate blending step, consistent with the mixing equipment specifications, should be performed to ensure adequate mixing.

Type C medicated feed

The Type B medicated feed must be diluted to a Type C medicated feed before being fed. Prepare a Type B medicated feed as described above. Thoroughly mix the Type B medicated feed to make 1 ton of Type C medicated feed to provide 13.5-16.0 g of virginiamycin per ton of complete feed on a 90% dry matter basis using the examples provided in Table 1.

Table 1. Type C medicated feed

lb of a 1,000 g/ton Type B medicated feed per ton of complete feed	Virginiamycin concentration in complete feed 90% dry matter basis (g/ton)
27	13.5
32	16.0

Feed continuously as sole ration

Store At Or Below 25°C (77°F), Excursions Permitted Up To 40°C (104°F)

Close container after use

Not For Human Use

Restricted Drug (California) - Use Only as Directed

SKU:

LOT No.:

DATE OF EXPIRATION:

Distributed by:

Elanco US Inc.
Indianapolis, IN 46221

Made in Canada.



8/8/25 9:17 AM

HEPAFLEX

virginiamycin powder

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:58198-1504
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VIRGINIAMYCIN (UNII: C49WS9N75L) (VIRGINIAMYCIN - UNII:C49WS9N75L)	VIRGINIAMYCIN	227 g in 0.45 kg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
MINERAL OIL (UNII: T5L8T28FGP)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
SOYBEAN OIL (UNII: 241ATL177A)	
RICE BRAN (UNII: R60QEP13IC)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-1504-1	25 kg in 1 BOX		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA140998	10/15/2025	

Labeler - ELANCO US. INC (966985624)

Registrant - ELANCO US. INC (966985624)

Establishment

Name	Address	ID/FEI	Business Operations
Phibro Saude Animal Internacional Ltda		907147578	api manufacture

Revised: 9/2025

ELANCO US. INC